

# Electromagnetic follow-ups in the era of forecasting gamma-ray bursts

Subtitle here if needed

Sarp Akcay<sup>1,2</sup>, Antonio Martin-Carrillo<sup>3</sup>, and Morgan Fraser<sup>3</sup>

<sup>1</sup> Theoretisch-Physikalisches Institut, Friedrich-Schiller-Universität Jena, 07743, Jena, Germany

<sup>2</sup> School of Mathematics & Statistics, University College Dublin, Belfield, Dublin 4, Ireland

<sup>3</sup> School of Physics, University College Dublin, Belfield, Dublin 4, Ireland

October 12, 2018

## ABSTRACT

The detection of gravitational waves from the binary neutron star inspiral-merger event GW170817 and the subsequent extended electromagnetic follow-up observations of the resulting kilonova gave us a small taste of multi-messenger astronomy across the spectra of *two* fundamentally different kinds of radiation. The opportunities to conduct such multi-disciplinary study will increase by two orders of magnitude in the 2030s with Einstein Telescope, LIGO’s European successor. Due to its extreme sensitivity in the 1 – 10 Hz regime, the Einstein Telescope’s C configuration (ET-C) will be capable of detecting inspiralling binary neutron star systems out to luminosity distances of 1 Gpc. For inspirals within half of this distance ET-C will accumulate signal-to-noise ratios of  $\geq 15$  with more than an hour left to merger. However, the localization of ET alone is rather poor: within  $z = 0.1$  we expect to have  $\sim 5$  binary neutron stars to be localized to  $\Delta\Omega \lesssim 10 \text{ deg}^2$ . On the other hand, a second less sensitive gravitational-wave detector (such as future KAGRA) would increase the number of well-localized sources to  $O(100)$ . Thus it is imperative to have at least one companion detector to ET with significantly improved seismic isolation in the 2030s. Having numerous GW sources localized to  $\sim 10 \text{ deg}^2$  opens the possibility of doing detailed follow-up observations of the resulting kilonovae with ATHENA, LSST, BlackGEM ... Here we explore this intriguing possibility... Thus, this letter is an appeal/plea(?) to the astronomy community to have in place ...

**Key words.** gravitational waves – gamma-ray bursts – kilonovae

## 1. Introduction

Gravitational waves offer a unique insight into some of the most extreme physical processes in the Universe - including the merger of black holes (BH) and neutron stars (NS), and the first seconds of core-collapse supernovae explosions.

With the first direct detection of gravitational waves (GWs) in 2015 by the Advanced Laser Interferometer Gravitational-Wave Observatory (Advanced LIGO; Abbott et al. 2016c), gravitational wave astronomy moved from prospect to reality. The first GW source observed by Advanced LIGO, GW150914, matched the signal predicted for the merger of two black holes with masses 36 and 29  $M_{\odot}$ . Along with being the first direct detection of GWs, GW150914 was also the first detection of such heavy black holes, which had significantly larger masses compared to those measured for Galactic high mass x-ray binaries. Such massive black holes provide an interesting constraint on stellar evolutionary channels at low metallicity (e.g. Abbott et al. 2016a; Belczynski et al. 2016). While no electromagnetic counterpart is generally expected to accompany the merger of two black holes, an intensive multi-wavelength search of the probable location of GW150914 was carried out (Abbott et al. 2016b). Despite yielding a null result, this effort served as a rehearsal in preparation for searches for counterparts to GW sources that *are* expected to be accompanied by an electromagnetic (EM) source.

Only two years after the first detection of merging black holes by Advanced LIGO, both the Advanced LIGO and Virgo gravitational wave observatories detected GW170817, with a

waveform consistent with the merger of two neutron stars (Abbott et al. 2017c). A spatially and temporally coincident short Gamma Ray Burst (GRB) was also seen by the *Fermi* and *INTEGRAL* satellites (Abbott et al. 2017b). This discovery sparked a global effort to find the counterpart of GW170817 at optical wavelengths, which resulted in the identification of AT2017gfo less than 11 hours later (Abbott et al. 2017d). AT2017gfo faded exceptionally rapidly, and displayed cool temperatures and lines from unusual r-process elements at exceptionally high velocities (Smartt et al. 2017; Arcavi et al. 2017; Pian et al. 2017; Coulter et al. 2017; Kilpatrick et al. 2017). These characteristics marked AT2017gfo as a kilonova; a transient powered by the radioactive decay of short-lived nuclides formed in the merger of two neutron stars.

The detection of electromagnetic counterparts to gravitational waves from merging neutron stars is of exceptional significance for astrophysics. Kilonovae are the predominant site for r-process nucleosynthesis, and so play a critical role in the chemical evolution of galaxies. If a kilonova can be identified and associated with a host galaxy of known redshift, then the degeneracy between inclination angle and distance inherent to a GW signal can be broken. This in turn allows the opening angle of the GRB jet to be constrained, something that has only been done for a handful of GRBs to date (Jin et al. 2018). Gravitational wave sources can also be used to independently determine the Hubble constant  $H_0$  (Abbott et al. 2017a). This of particular interest given the disagreement between measurements of  $H_0$

from Type Ia SNe and from the Cosmic Microwave Background (e.g. Bernal et al. 2016)

The identification of AT2017gfo as the counterpart to GW170817 was realised by the ability of Advanced LIGO-Virgo to localise the GW signal to  $\sim 30 \text{ deg}^2$ . In addition, at only 40 Mpc, GW170817 was exceptionally close. This enabled the EM counterpart to be identified through targeted observations of galaxies which were at this distance within the GW localisation region (Coulter et al. 2017). Unfortunately such a strategy is only feasible for the nearest GW sources, and rapidly becomes unfeasible beyond  $\sim 100 - 200 \text{ Mpc}$ , both as the number of galaxies within the search volume increases, and as the fraction of galaxies with reliable redshifts decreases. This embarrassment of riches is a serious obstacle for identifying EM counterparts to GW transients in the 2030s with Einstein Telescope (Abernathy et al. 2011).

Einstein Telescope (ET) will be sensitive enough to “pick up” GW sources at a few Hz thanks to its cryogenic design and underground housing which will shield it from low-frequency contaminants such as seismic and gravity-gradient noises. Moreover, ET will consist of three V-shaped interferometers which eliminate blind spots and further allow it to construct a null stream (Sathyaprakash et al. 2012) which can be used to veto spurious events (Wen & Schutz 2005). Additionally, ET will be a xylophone (Hild et al. 2010), i.e., a multi-band detector capable of delivering high sensitivities both at low frequencies ( $\sim 5 \text{ Hz}$ ) and high frequencies ( $\sim 100 \text{ Hz}$ ). Here, we focus on the C configuration (ET-C) which offers the highest low-frequency sensitivity as shown in Fig. 1. ET-C will detect  $\gtrsim O(10^3)$  binary neutron star inspirals per year out to 1000Mpc with SNRs  $\gtrsim 30$  (Akca 2018). A subset of these sources will be close enough that they will be detected a few hours before their respective mergers (Akca 2018), hence opening up the possibility of alerting EM observatories to conduct follow-up observations *before, during* and *after* the prompt gamma-ray bursts. Additionally, ET-C will forecast a few potential tidal disruption events per year, in which a neutron star gets tidally torn by a  $\sim 5M_\odot$ , high-spin black hole companion.

To fully exploit the prospect of multi-messenger astronomy, a number of wide-field survey telescopes are either operational, in commissioning, or under construction. Foremost among these is the Large Synoptic Survey Telescope (LSST; LSST Science Collaboration et al. 2009), which has an 8.4 m primary mirror, and will image  $3.5 \text{ deg}^2$  in a single pointing. Construction of LSST is well underway, and the telescope is expected to begin full survey operations at the start of 2023. Apart from LSST, the majority of current and next-generation survey telescopes have a relatively small mirror, but a large camera, and are designed to observe  $\sim 10 - 50 \text{ sq degrees}$  in a single pointing to a limiting magnitude of  $\sim 20 - 22$ . ZTF (Bellm 2014), GOTO (Dyer et al. 2018) and ATLAS (Tonry 2011) are all currently operational at present, while BlackGEM is currently under construction (Bloemen et al. 2016).

There has been a considerable amount of discussion in the literature as to the optimal strategy to identify an EM counterpart to future GW transients (Gehrels et al. 2016; Coward et al. 2011; Ghosh et al. 2016; Chan et al. 2017; Siellez et al. 2014; Antolini & Heyl 2016). In most cases however, a large number of candidates will be found within the search region, for which further spectroscopic followup observations will be required. This spectroscopic classification bottleneck will remain a problem

Our aim here is to demonstrate exciting EM follow-up studies that can be done by taking advantage of the early GW warning capability of ET. More specifically we consider binary neu-

tron star inspirals out to luminosity distances of  $\sim 600 \text{ Mpc}$ , the expected range of LSST. Within this range, ET-C will be able to detect the inspiral GWs a few hours before a given merger thus provide a window of opportunity for mobilizing the EM observatories in time to witness the birth of the associated kilonova. Furthermore, a pre-GW warning allows us to obtain reference images with EM telescopes immediately prior to a kilonova, reducing the number of unrelated transients that will be found within a given search region. However, in order to fully benefit from ET’s early warnings, several issues must be addressed: (i) ET’s poor localisation by itself, (ii) large number of supernovae creating a confusion background, (iii) the slow response time of certain EM observatories essential to follow-up. Here, we consider each of these setbacks and suggest solutions which require support from the global astronomy community.

This letter is organized as follows: Sec. 2 provides more details on ET, Sec. 3 investigates the implications of optical follow-up. Sec. 4 ... We use  $f$  to denote the quadrupole GW frequency in the detector frame.  $c$  is the speed of light and  $G$  is Newton’s constant.

## 2. Einstein Telescope

In this section, we compute advance warning times ( $T_{\text{AW}}$ ) ET will provide. The computational details are provided in Akca (2018). To this end, consider a binary neutron star (BNS) system with component masses  $m_1, m_2$  inspiraling at a luminosity distance  $D$  with a corresponding redshift  $z$ . For GW frequencies of interest to us here ( $f \lesssim 10 \text{ Hz}$ ), the binary undergoes an adiabatic inspiral dominated by the emission of leading-order (quadrupole) gravitational radiation. By balancing the power emission in GWs to the rate of change of binding energy, we obtain the frequency evolution of the GW frequency

$$\dot{f} = \frac{96}{5} \pi^{8/3} \frac{(GM_c)^{5/3}}{c^5} f^{11/3}, \quad (1)$$

where  $M_c = (m_1 m_2)^{3/5} (m_1 + m_2)^{-1/5}$  is the chirp mass. After fixing an integration constant, Eq. (1) can be integrated to yield the time left to merger at a given frequency, usually called the inspiral time

$$\begin{aligned} \tau_{\text{insp}}(f) &= \frac{5}{256\pi} \frac{c^5}{(\pi G M_c)^{5/3}} f^{-8/3} \\ &= 16.72 \text{ minutes} \left( \frac{1.219 M_\odot}{M_c} \right)^{5/3} \left( \frac{10 \text{ Hz}}{f} \right)^{8/3}. \end{aligned} \quad (2)$$

This result can be supplemented with a post-Newtonian series up to  $O(c^{-7})$  (Blanchet 2014), but the resulting expressions are rather ungainly and only change  $\tau_{\text{insp}}$  by  $\lesssim 2\%$ .

To obtain  $T_{\text{AW}}$  we must choose  $f$  which necessitates a brief discussion on the effects GWs have on interferometers (IFOs). It is a well known result in general relativity that passing GWs induce a scalar response in a given IFO known as the GW strain which is a function of GW polarization amplitudes and IFO antenna pattern functions. In frequency domain, the norm of the GW strain is given by  $|\tilde{h}(f)| = A h_0 f^{-7/6} |Q|$ , where  $A = \pi^{-2/3} (5/24)^{1/2}$ ,  $h_0 = c(1+z)^{-1} \tilde{M}^{5/6} / D$  with  $\tilde{M} = G(1+z)M_c c^{-3}$  and  $Q$  is the IFO quality factor which is a function of source sky location angles  $(\theta, \phi)$ , its inclination  $\iota$ , and the relative detector-source polarization angle  $\psi$ .

The IFO response to a GW strain is quantified in terms of a signal-to-noise ratio (SNR). As we can not a priori know the

angles  $\{\theta, \phi, \iota, \psi\}$ , we use an angle-RMS-averaged SNR, which for a detector with triangular topology like ET, reads

$$\rho_{\text{ET}}(f_1, f_2) = \frac{6}{5} A h_0 (1+z)^{-1/6} \left[ \int_{f_1}^{f_2} df' \frac{f'^{-7/3}}{S_n(f')} \right]^{1/2}, \quad (3)$$


where  $\sqrt{S_n(f)}$  is the *amplitude spectral density* (ASD) of the detector (also called detector noise) and the factor of  $6/5$  in Eq. (3) is due to RMS-averaging. For each BNS source, this factor may vary by  $\approx \pm 30\%$ , but will always be  $> 0$  thanks to ET not having any blind spots.

We now use Eq. (3) to provide a precise definition of  $T_{\text{AW}}$ . Keep in mind that the advance warning time is the time interval between the moment of detection and the merger, therefore we must first define the former. We do this as follows: let  $f_0$  be the frequency at which the GW strain equals the detector noise, i.e.,  $\sqrt{S_n(f_0)} = 2\sqrt{f_0 \tilde{H}_{\text{ET}}(f_0)}$  where  $\tilde{H}_{\text{ET}}(f) = 3h_0 f^{-7/6}/5$ . The moment of detection is given by  $\bar{f} > f_0$  such that  $\rho_{\text{ET}}(f_0, \bar{f}) = 15$ . Then we immediately have  $T_{\text{AW}} = \tau_{\text{insp}}(\bar{f})$  and the total accumulated SNR is given by  $\rho_{\text{tot}} = \rho_{\text{ET}}(f_0, f_{\text{ISCO}})$ , where  $f_{\text{ISCO}}$  is the frequency at which the inspiral transitions to plunge. Here, we use the standard approximation from general relativity:  $f_{\text{ISCO}} \approx \frac{c^3}{6^{3/2}\pi G(m_1+m_2)} \simeq 1571 \left( \frac{2.8M_\odot}{m_1+m_2} \right)$  Hz. Our chosen threshold SNR of 15 is not set in stone; other common choices are 8 and 12, but we choose to be more conservative.

The only free variable left to determine  $T_{\text{AW}}$  is the luminosity distance  $D$  ( $z$  is obtained from it and vice versa). In Fig. 1 we display the GW strain for four canonical ( $m_1 = m_2 = 1.4M_\odot$ ) BNS inspirals at  $D = 100, 200, 400, 600$  Mpc. The corresponding ASDs scale as  $\sqrt{f} f^{-7/6} = f^{-2/3}$ , thus are straight lines with slopes of  $-2/3$  in the figure. ET-C's noise is the thick, red curve with highest sensitivity for  $f \lesssim 30$  Hz. We can clearly see the frequencies  $f_0$  where the straight lines intersect ET-C's noise curve. We list the advance warning times along with the total SNRs for these four sources in Table 1. We can see that ET-C is capable of providing up to five hours of early warning before merger, thus offering a tremendous opportunity to electromagnetically observe the merger-GRB-kilonova with available resources in the 2030s. But we should first see if ET can forecast enough BNS events and localise them to make a follow-up campaign worthwhile.

**Table 1.** Forecasting capabilities of the C configuration of Einstein Telescope summarized in terms of luminosity distance  $D$  and the corresponding redshift  $z$ .  $\bar{f}$  is the threshold frequency at which ET-C accumulates SNR of 15.  $T_{\text{AW}}$  is the inspiral time from the instant when the GW frequency equals  $\bar{f}$ .  $\rho_{\text{tot}}$  is the total accumulated SNR of each inspiral. Last column lists the event rates in the format  $\text{av}_{\text{min}}^{\text{max}}$  in units of  $D^{-3} \text{ yr}^{-1}$ . The numbers have been rounded when appropriate. **SA: Morgan, should we add apparent magnitude to this table?**

$D$ (Mpc)	$z$	$\bar{f}$ (Hz)	$T_{\text{AW}}$ (hrs)	$\rho_{\text{tot}}$	$R_{\text{av}}^{\text{max}}_{\text{min}}$
100	0.022	3.3	5.3	365	$1.54^{4.7}_{0.32}$
200	0.044	4.1	2.9	182	$12.3^{38.}_{2.6}$
300	0.065	4.66	1.9	121	$41.6^{130}_{8.6}$
400	0.085	5.1	1.5	90.5	$98.6^{300}_{20}$
500	0.10	5.4	1.2	72	$193^{590}_{40}$
600	0.12	5.7	1.03	60	$333^{1020}_{69}$



**Fig. 1.**  $1.4M_\odot - 1.4M_\odot$  inspiralling BNS systems sweeping across the sensitivity band of Einstein Telescope's C configuration (thick red curve). The solid (black), dotted (blue), dashed (green), and dot-dashed lines (gray) lines are the redshift-corrected angle-RMS-averaged GW strains,  $2\sqrt{f}\tilde{H}_{\text{ET}}$ , at luminosity distances of  $D = 100, 200, 400, 600$  Mpc, respectively. The vertical lines with correspondingly identical patterns (colors) mark the redshifted ISCO frequencies  $(1+z)^{-1} f_{\text{ISCO}}$  at which point we terminate each inspiral. As the true ISCO frequency is likely larger than  $f_{\text{ISCO}}$  (Marronetti et al. 2004), the inspirals would continue to nearly 2 kHz indicated by the faded lines in the plot. We also show the strain sensitivity of Cosmic Explorer (thick blue) and KAGRA+ (brown curve). The faint brown curve represents the sensitivity of Advanced LIGO during GW170817. As RMS averaged BNS strains in an L-topology detector (CE, KAGRA+, LIGO) are  $2/3$  of strains in triangular-topology detector (Akcay 2018), we rescale the CE, KAGRA+, LIGO curves by  $3/2$  thus making them effectively less sensitive. Otherwise, we would have to draw four additional BNS strains, each of which scaled down by  $2/3$  in amplitude.

We base our event rate calculations on  $R = 1540^{+3200}_{-1220} \text{ Gpc}^{-3} \text{ yr}^{-1}$  inferred from GW170817 (Abbott et al. 2017c). Assuming flat spatial geometry, this translates roughly to  $330^{+690}_{-260}$  yearly BNS events within 600 Mpc. We partition this volume into concentric spheres with radii equalling integer multiples of 100 Mpc. For each sphere, we compute the corresponding average (av), maximum (max), and minimum (min) event rates using  $R$ . We present these results in Table 1 along with the redshifts for each distance computed using a flat  $\Lambda$ CDM model ( $\Omega_k = 0$ ) with the latest Planck satellite parameters:  $\Omega_\Lambda = 0.6911$ ,  $\Omega_m = 0.3089$ ,  $H_0 = 67.74 \text{ km s}^{-1} \text{ Mpc}^{-1}$  (Adam et al. 2016). It is then straightforward to translate  $D$  to  $z$  [cf. Hogg (1999)].

We can see from Table 1 that an event as near as GW170817 is a once-per-decade occurrence, but within  $z \lesssim 0.1$ , we can expect ET-C to annually detect GWs from 40 to 600 BNS transients. Roughly 7% of these will be within 200 Mpc yielding  $T_{\text{AW}} \gtrsim 3$  hours which means that EM observatories will be alerted 3 - 40 times yearly with the prospects of observing the birth of kilonovae. If we make the bold assumption that future EM observatories would suffice with an hour's warning then all the BNS transients within 600 Mpc become possible candidates for merger-kilonovae observations. In this case, early warnings would be sent by ET at least once a week and at most three times per day. It is not feasible to have a system in place to observe each one of these events so it will make sense to be selective and focus on the best candidates. These will naturally mostly be the nearest sources which will additionally be favoured by ET because of its limitations with localisation that we discuss next.

Though ET will have no blind spots because it consists of three V-IFOs in one plane, each rotated by  $120^\circ$ , it will have poor source localisation because all three IFOs are at the same location. However, as our findings in Table 1 indicate, ET-C will detect BNS inspirals hours before the merger during which time the Earth will have rotated by as much as a few dozen degrees. This means that over this time ET can be treated as a network. It is expected that within 1 Gpc, ET *alone* will localise approximately 20% of the sources to within  $100 \text{ deg}^2$ , 2% to within  $20 \text{ deg}^2$ , and 0.5% to within  $10 \text{ deg}^2$  (Zhao & Wen 2018)<sup>1</sup>.

Let us be more specific and pick  $D = 400 \text{ Mpc}$  as a cut-off distance. Given that the  $10 \text{ deg}^2$ -localisable sources make up 0.5% of the detectable population within 1 Gpc, this percentage increases roughly to  $2.5^3 \times (0.5\%) \approx 7\%$  within 400 Mpc because there are  $(2.5^3 - 1)$  times as many sources within  $400 \text{ Mpc} \leq D \leq 1 \text{ Gpc}$  than  $D \leq 400 \text{ Mpc}$ . Rounding down to 5% to err on the side of caution, we obtain 5 yearly BNS transients with  $\Delta\Omega \lesssim 10 \text{ deg}^2$  and  $\sim 20$  transients with  $\Delta\Omega \lesssim 20 \text{ deg}^2$  within 400 Mpc corresponding to  $T_{\text{AW}} \gtrsim 1.5 \text{ hrs}$ .

Our above approximations on localisation should be taken as the pessimistic case as ET will be not be operating alone. Currently suggested companion detectors are the Japanese cryogenic detector KAGRA (Akutsu et al. 2017, 2018), LIGO’s successor Voyager in the US (Berger et al. 2016) and its “relative” in India (Unnikrishnan 2013), and finally, Cosmic Explorer (CE): the ambitious third generation US detector with 40 km armlength (Abbott et al. 2017). Assuming a three detector network consisting of ET and two CEs with a total network SNR  $> 12$  and two detectors each with SNR  $> 5$ , Mills et al. (2018) find that half of signals will be localised to within  $1 \text{ deg}^2$  out to a redshift of  $z \sim 0.25$ . However, this survey is not concerned with issuing sufficiently early warnings to EM facilities. The problem is that only ET has the extreme low-frequency sensitivity enabling  $T_{\text{AW}} \sim \text{hours}$ . The other detectors will not accumulate any SNR from BNS inspirals in the  $f \lesssim 5 \text{ Hz}$  domain<sup>2</sup>. However, CE will be sensitive enough to accumulate SNR from 5 Hz for BNSs with  $D \lesssim 400 \text{ Mpc}$  (Fig. 1). Given that its sensitivity increases steeply between 5 and 10 Hz, CE will accumulate SNR = 5 with 1.5 hours left to merger and SNR = 15 with 1.25 hours left resulting in a total network SNR,  $\rho_{\text{net}} \equiv (\rho_{\text{ET}}^2 + \rho_{\text{CE}}^2)^{1/2} = \{18.8, 27.4\}$  for  $\tau_{\text{insp}} = \{1.5, 1.25\}$  hours, respectively. As localisation improves with increasing SNR, this means that an initial  $\Delta\Omega$  of  $\sim 100 \text{ deg}^2$  can be reduced by as the BNS inspirals through its second to last hour before the merger.

This region can be further decreased once the BNS enters a third detector’s bandwidth, which will most likely be some future version of KAGRA. We do not have any strain data for a 2030s KAGRA. However, there are plans for a mid 2020s upgrade, KAGRA+, with increased strain sensitivity (brown curve in Fig. 1). Using the same analysis as for CE, we obtain that KAGRA+ will pick up a 400-Mpc inspiral at  $\sim 10 \text{ Hz}$  and will accumulate SNR = 5 within a minute. For a less likely 100-Mpc source, KAGRA+ would accumulate SNR  $> 15$  more than a half hour before the merger. Thus, for nearby sources, even KAGRA+ sensitivity could contribute to pre-merger localisation efforts. **SA: How much time does LSST need before moving to final target?** Once again, this is rather a pes-

simistic estimation as we expect KAGRA’s 3G successor to be more sensitive than KAGRA+.

In short, within 400 Mpc, we can annually expect five BNS transients to be localised to  $10 \text{ deg}^2$  1.5 hours before the merger. We can expect an additional  $\sim 15$  more with initial localisation of  $\sim 20 \text{ deg}^2$  which we expect will be narrowed down to  $\sim 10 \text{ deg}^2$  about one hour before the merger. We envision a three-stage localisation procedure whereby ET conducts the operations alone until  $f \sim 5 \text{ Hz}$  — roughly two hours before the merger — at which point CE joins in and, finally, around  $f \sim 10 \text{ Hz}$  KAGRA+ (or 3G) starts accumulating SNR with  $\gtrsim 15$  minutes left to merger. For sources closer than 400 Mpc these estimations improve.

### 3. Implications for optical followup of GW detections

Identifying an optical or near-infrared (NIR) counterpart to a GW is an observational challenge. If a GW is only localised to tens, or even hundreds of square degrees, then we must survey a large area of the sky to find an EM counterpart. While large format CCDs make taking imaging of an area of  $\sim 100 \text{ sq degrees}$  relatively straightforward, we must identify our EM counterpart of interest among the many unrelated astrophysical transients that we expect by chance within the same area. Thus far, this has relied upon large scale efforts to spectroscopically classify credible candidates that are found within the sky localisation of a GW. As an example, for the BH merger GW151226, Smartt et al. (2016) found 49 candidate transients within  $290 \text{ deg}^2$ , and obtained spectra for 20 of these. While such a survey strategy is the only feasible approach at present, it is clearly an inefficient use of scarce telescope time.

The early warning obtained for future GW events discussed in Sect. 2 offers an alternative approach for finding EM counterparts. In brief, if we can detect a GW with  $\sim 1 \text{ hr}$  advance warning, and can localise it to  $\sim 50 \text{ deg}^2$ , then we can obtain imaging of this area both immediately prior to, and after, the merger happens. Since the merger will be the only thing that has changed over such a short period of time, identifying an EM counterpart in difference imaging becomes straightforward.

**Morgan:** I would suggest repeating this for  $\sim 10 \text{ deg}^2$  localisation as well

#### 3.1. The rates and nature of contaminants


There are broadly three classes of contaminants that we must consider when searching for EM counterparts to GW; stellar variables and flares such as cataclysmic variables (CV); variability in Active Galactic Nuclei (AGN); and supernovae (SNe). The first class of contaminants show a strong dependence on Galactic latitude (Drake et al. 2014), and are concentrated in the disk of the Milky Way. In addition, for at least some CV outbursts ...

AGN can often be identified through their historical lightcurves, which may show previous variability. Given the relatively straightforward removal of stellar and AGN contaminants, we are left with SNe as the dominant contaminant. Three quarters of SNe are SNe Ia in a mag limited survey (cf LOSS). Also, this is borne out by the experience of Smartt et al. (2016), where they found...

<sup>1</sup> Zhao & Wen (2018) perform their calculations for ET-D which in fact has slightly worse sensitivity for  $f \lesssim 5 \text{ Hz}$  than ET-C, cf. Fig. 19 of Pitkin et al. (2011)

<sup>2</sup> There is a prospect for improving LIGO’s low-frequency end called LIGO-LF (Yu et al. 2018).

*Acknowledgements.* S. A. acknowledges support by the EU H2020 under ERC Starting Grant, no. BinGraSp-714626.



**Fig. 2.** More or less a placeholder. Number of contaminant SN Ia within our search region as a function of distance...

## References

- Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2016a, *ApJ*, 818, L22  
 Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2016b, *ApJ*, 826, L13  
 Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2016c, *Physical Review Letters*, 116, 061102  
 Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2017a, *Nature*, 551, 85  
 Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2017b, *ApJ*, 848, L13  
 Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2017c, *Physical Review Letters*, 119, 161101  
 Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2017d, *ApJ*, 848, L12  
 Abbott, B. P. et al. 2017, *Class. Quant. Grav.*, 34, 044001  
 Abernathy, M., Acernese, F., Ajith, P., et al. 2011, *Einstein gravitational wave Telescope Conceptual Design Study*, Tech. rep., European Commission  
 Adam, R. et al. 2016, *Astron. Astrophys.*, 594, A1  
 Akcay, S. 2018, *Annals der Physik*, 441, 1186  
 Akutsu, T., Cadonati, L., Cavaglià, M., et al. 2017, in *15th International Conference on Topics in Astroparticle and Underground Physics (TAUP 2017)* Sudbury, Ontario, Canada, July 24–28, 2017  
 Akutsu, T. et al. 2018, *PTEP*, 2018, 013F01  
 Antolini, E. & Heyl, J. S. 2016, *MNRAS*, 462, 1085  
 Arcavi, I., Hosseinzadeh, G., Howell, D. A., et al. 2017, *Nature*, 551, 64  
 Belczynski, K., Holz, D. E., Bulik, T., & O’Shaughnessy, R. 2016, *Nature*, 534, 512  
 Bellm, E. 2014, in *The Third Hot-wiring the Transient Universe Workshop*, ed. P. R. Wozniak, M. J. Graham, A. A. Mahabal, & R. Seaman, 27–33  
 Berger, B. et al. 2016, *What Comes Next for LIGO? 2016 LIGO-DAWN Workshop II*, <https://wiki.ligo.org/pub/LSC/LIGOWorkshop2016/WebHome/Dawn-II-Report-SecondDraft-v2.pdf>  
 Bernal, J. L., Verde, L., & Riess, A. G. 2016, *J. Cosmology Astropart. Phys.*, 10, 019  
 Blanchet, L. 2014, *Living Rev. Rel.*, 17, 2  
 Bloemen, S., Groot, P., Woudt, P., et al. 2016, in *Proc. SPIE*, Vol. 9906, *Ground-based and Airborne Telescopes VI*, 990664  
 Chan, M. L., Hu, Y.-M., Messenger, C., Hendry, M., & Heng, I. S. 2017, *ApJ*, 834, 84  
 Coulter, D. A., Foley, R. J., Kilpatrick, C. D., et al. 2017, *Science*, 358, 1556  
 Coward, D. M., Gendre, B., Sutton, P. J., et al. 2011, *MNRAS*, 415, L26  
 Drake, A. J., Gänsicke, B. T., Djorgovski, S. G., et al. 2014, *MNRAS*, 441, 1186  
 Dyer, M. J., Dhillon, V. S., Littlefair, S., et al. 2018, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 10704, *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, 107040C  
 Gehrels, N., Cannizzo, J. K., Kanner, J., et al. 2016, *ApJ*, 820, 136  
 Ghosh, S., Bloemen, S., Nelemans, G., Groot, P. J., & Price, L. R. 2016, *A&A*, 592, A82  
 Hild, S., Chelkowski, S., Freise, A., et al. 2010, *Class. Quant. Grav.*, 27, 015003  
 Hogg, D. W. 1999 [arXiv:astro-ph/9905116]  
 Jin, Z.-P., Li, X., Wang, H., et al. 2018, *ApJ*, 857, 128  
 Kilpatrick, C. D., Foley, R. J., Kasen, D., et al. 2017, *Science*, 358, 1583  
 LSST Science Collaboration, Abell, P. A., Allison, J., et al. 2009, *ArXiv e-prints* [arXiv:0912.0201]

- Marronetti, P., Duez, M. D., Shapiro, S. L., & Baumgarte, T. W. 2004, *Phys. Rev. Lett.*, 92, 141101  
 Mills, C., Tiwari, V., & Fairhurst, S. 2018, *Phys. Rev.*, D97, 104064  
 Pian, E., D’Avanzo, P., Benetti, S., et al. 2017, *Nature*, 551, 67  
 Pitkin, M., Reid, S., Rowan, S., & Hough, J. 2011, 14  
 Sathyaprakash, B. et al. 2012, *Class. Quant. Grav.*, 29, 124013, [Erratum: *Class. Quant. Grav.* 30, 079501 (2013)]  
 Siellez, K., Boër, M., & Gendre, B. 2014, *MNRAS*, 437, 649  
 Smartt, S. J., Chambers, K. C., Smith, K. W., et al. 2016, *ApJ*, 827, L40  
 Smartt, S. J., Chen, T.-W., Jerkstrand, A., et al. 2017, *Nature*, 551, 75  
 Tonry, J. L. 2011, *PASP*, 123, 58  
 Unnikrishnan, C. S. 2013, *Int. J. Mod. Phys.*, D22, 1341010  
 Wen, L. & Schutz, B. F. 2005, *Class. Quant. Grav.*, 22, S1321  
 Yu, H. et al. 2018, *Phys. Rev. Lett.*, 120, 141102  
 Zhao, W. & Wen, L. 2018, *Phys. Rev.*, D97, 064031