
Second-line and later treatment, monthly cost (D05) [±25%]
First-line treatment, monthly cost, with pembrolizumab (D03P)
Utility, event-free after treatment (U02)
Utility, distant recurrence (U04)
Utility, locoregional recurrence (U03)
End-of-life care cost (D06) [mirrored]
Follow-up cost per year, year 6 onward (C03) [mirrored]
Locoregional recurrence, initial cost (D01) [±25%]
Pembrolizumab cost per dose (A03) [price]
First-line treatment duration, with pembrolizumab (D04P) [±25%]
Musculoskeletal AEs, any grade, with exercise (E06)
Follow-up cost per year, years 3 to 5 (C02) [mirrored]
Musculoskeletal AEs, any grade, without exercise (E07)
Musculoskeletal AEs, grade 3 or higher, with exercise (E09)
Locoregional recurrence, monthly cost (D02) [±25%]
Utility decrement, grade 3 or higher AE (U05)
Musculoskeletal AE, cost per event (E08) [mirrored]
Musculoskeletal AEs, grade 3 or higher, without exercise (E10)
Follow-up cost per year, years 1 to 2 (C01) [mirrored]
Drug administration cost per visit (A04)
Utility, event-free during treatment (U01)
Input at its lower bound
Input at its upper bound
2,600
2,650
2,700
2,750
2,800
2,850
Maximum affordable program cost (JPY 1,000)