

Recursive Spectral Fibers on Simplicial Cobordisms

A geometric program for quarks, color, fermion statistics, Fock space, and baryons

Tessera - cobordism programme

Implementation is tracked by GitHub epic [#763](#).

Abstract

This paper proposes a single geometric formulation for the particle content already suggested by Tessera's cobordism experiments. A connected, high-modularity simplicial component is treated as one effective vertex at the next resolution. The component's selected localized Hodge eigenspace is its fiber, and the couplings between components induce transport between those fibers. Repeating this operation produces a nested, potentially fractal hierarchy of complexes. No independent gauge field, particle label, or auxiliary lattice is introduced: every state, connection, exchange sign, and observable is read from oriented simplices, complex edge data, edge qubits, and the Hodge/Regge operators already present in the construction.

The proposal has a substantial exact core. Schur reduction proves when a component may be replaced by a vertex without changing boundary quadratic energies. The graded tensor product of chain complexes gives the correct Hodge operator for interacting systems. Three oriented edge qubits form the exact exterior algebra $\Lambda^\bullet \mathbb{C}^3 = \mathbf{1} \oplus \mathbf{3} \oplus \bar{\mathbf{3}} \oplus \mathbf{1}$; the one-occupation sector is a color qutrit, its bilinears close $\mathfrak{su}(3)$, the three-occupation sector is a color singlet, and the grading gives the fermionic exchange sign and canonical anticommutation relations. A connection is derived from overlap of neighboring spectral frames, and Wilson loops are then gauge-invariant observables rather than new degrees of freedom. Successive cobordism interactions generate the finite stages of an inductive-limit Fock space.

The physical identification remains a hypothesis to be tested. A quark is proposed to be a persistent, odd-parity, rank-three spectral fiber localized on a high-modularity component; a proton is three such components bound into one persistent supercomponent, occupying a normalized color wedge, carrying baryon number +1, electric charge +1, and a total-space spin-1/2 holonomy. The paper separates exact identities, results already supported by Tessera, and new falsifiable conjectures. It also states an implementation program in which analytic or structure-exact operations are used before iterative numerical work, so the model can scale without turning its defining observables into uncontrolled approximations.

Contents

1	Epistemic status and design constraint	1
2	Present evidence in Tessera	1
3	The microscopic geometric state	2
4	A component is exactly an effective vertex	3
5	Recursive spectral fibers	3
6	Interactions and the expanding Hilbert space	4
7	A triangle carries the exact color algebra	6
7.1	Geometric normalization	7
7.2	The existing omega phase pattern	7
8	Color transport and Wilson loops without a new gauge field	7
9	Quarks as modular clusters	8
10	Fermion statistics from simplicial orientation	9
10.1	A label-independent exchange experiment	9
11	Fock space as an inductive limit of interactions	10
12	The proton as the maximally informative baryon	11
13	The master recursive construction	11
14	Exactness and performance principles	12
15	Prior art and boundary of novelty	12
16	Falsification program	13
17	Conclusion	14
	Prior-art references	15
	Repository evidence	16

1 Epistemic status and design constraint

Three kinds of statement are deliberately distinguished:

1. **Exact identity** — follows algebraically from the stated finite complex, orientation, and inner product.
2. **Existing Tessera evidence** — measured by an existing experiment, with the scope and residual reported in the repository.
3. **Proposed physical identification** — a new hypothesis with an explicit falsification test.

The governing constraint is parsimony. The ontology is limited to:

- an oriented simplicial complex and its cobordisms;
- a complex squared length and a normalized qubit on each edge;
- incidence, Hodge, and Regge operators derived from that data; and
- tensor composition when systems interact.

Spectral fibers, color frames, connections, Wilson loops, particle sectors, and coarse vertices are *derived views* of that same data. They are not separately sampled fields. This is important both scientifically and computationally: adding a new independent field could fit a desired answer, while deriving every readout from one complex leaves the construction falsifiable.




Figure 1: Concept map for the recursive complex construction. Colors encode epistemic status, not physical sectors: blue is established or exact machinery, green is a derived observable, and amber is a proposed physical identification.

2 Present evidence in Tessera

Two existing results motivate the construction.

First, the state-operation-cobordism experiments show that the Hodge-carried register is an isometry to machine precision and that its spectral value reproduces the quantum transition amplitude for every operation that the tested geometry actually carries. Generic fixed-complexity operations can remain obstructed, and the obstruction is visible both as a residual floor and as leakage from the carried subspace. The claim is therefore not that every finite complex realizes every gate; it is that a realized, isometrically embedded register computes the corresponding amplitude. See [cobordism-results.md](#).

Second, interaction-history complexes exhibit a stable near-four-dimensional spectral regime. The strongest reported measurements approach, but do not yet prove, an exact spectral dimension of

four. The current status and finite-size caveats are recorded in [h_ds4_status.md](#). Diffusion-based spectral dimension on simplicial quantum geometries has important precedent in causal dynamical triangulations [1]; the Tessera evidence is an independent result for a different construction and should be compared at the level of the return-probability estimator and its finite-size window.

The current proton animation adds a third, more preliminary observation. It starts with the phase pattern $\{1, \omega, \omega^2\}$ and evaluates its singlet diagnostics while a joint Regge-Hodge stationarity objective changes the complex. It no longer forces register holes to appear, and holes have not re-emerged in the current construction. That negative result is useful: the proposed quark should therefore not be defined as a hole. It will be sought as a persistent modular spectral cluster, while Betti numbers remain independent topological observables.

3 The microscopic geometric state

Let K be a finite oriented simplicial complex. For every edge e , store

$$x_e = (z_e, q_e), \quad z_e = \rho_e e^{i\theta_e} \in \mathbb{C}, \quad q_e = \alpha_e |0\rangle + \beta_e |1\rangle, \quad |\alpha_e|^2 + |\beta_e|^2 = 1.$$

The complex phase is not a second link field; it is part of the existing complex edge geometry. The qubit is the occupation/amplitude carrier. An implementation may choose a convention in which the relative qubit phase is locked to θ_e , but the mathematics only requires that the convention be explicit and invariant under edge reorientation.

Let

$$\partial_k : C_k(K) \longrightarrow C_{k-1}(K), \quad \partial_{k-1} \partial_k = 0$$

be the oriented boundary maps and let $w_k(z)$ be the metric weight on k -chains. With the weighted adjoint

$$\partial_k^* = W_k^{-1} \partial_k^\dagger W_{k-1},$$

the degree- k Hodge Laplacian is

$$L_k = \partial_{k+1} \partial_{k+1}^* + \partial_k^* \partial_k.$$

In the positive metric regime this is self-adjoint in the w_k inner product. In the signed Lorentzian regime it can be non-normal; then left and right spectral frames and their biorthogonal condition numbers must be reported rather than silently treating L_k as Hermitian.

The geometry evolves toward joint stationary points of the existing Regge and Hodge functionals. In emergence mode, particle-specific observables below are read after optimization and are not inserted as target terms. Controlled synthesis mode may pin a carrier to test realizability, but that is a separate experiment.

This operator stack sits on established foundations: Regge calculus encodes piecewise-flat gravity in simplicial deficit angles [2], discrete exterior calculus supplies metric-dependent chain/cochain

operators [3], and combinatorial Hodge spectra on simplicial complexes have a developed spectral theory [4]. Tessera’s proposal is not a replacement for those constructions; it is a constrained use of them as the sole source of the later particle readouts.

4 A component is exactly an effective vertex

Partition the k -cells of a connected component into interface cells $\mathbf{B}$ and interior cells $\mathbf{I}$, and block its Hodge operator as

$$L = \begin{pmatrix} L_{BB} & L_{BI} \\ L_{IB} & L_{II} \end{pmatrix}.$$

After projecting out incompatible interior zero modes, minimization over the interior has the exact solution

$$x_I^* = -L_{II}^+ L_{IB} x_B,$$

and the exact effective boundary operator

$$L_{\text{eff}} = L_{BB} - L_{BI} L_{II}^+ L_{IB}.$$

Here $+$ denotes the Moore-Penrose inverse on the supported interior subspace. For every compatible boundary value,

$$\min_{x_I} \begin{pmatrix} x_B \\ x_I \end{pmatrix}^\dagger L \begin{pmatrix} x_B \\ x_I \end{pmatrix} = x_B^\dagger L_{\text{eff}} x_B.$$

This is the precise sense in which a connected component can be replaced by a coarse vertex: all external quadratic response is preserved. The effective blocks between two coarse components become the coarse edges. A harmonic interior mode is not discarded; it becomes an explicit fiber coordinate attached to the coarse vertex.

This reduction is exact for the retained interface. Approximation enters only if one additionally truncates the interface or the fiber, and that approximation has a measurable residual.

For graph Laplacians this is the classical Kron reduction by Schur complement [5]. Spectral graph reduction provides related approximation guarantees when additional coarsening or truncation is performed [6]. The extension proposed here is to apply the same response-preserving logic degree by degree to the weighted Hodge blocks and to retain localized zero or near-zero modes as explicit fiber coordinates rather than silently eliminating them.

5 Recursive spectral fibers

Let $P_\ell = \{C_v^\ell\}$ be an intrinsic partition of the complex at scale ℓ into persistent, high-modularity connected components. Define the next complex by

$$K_{\ell+1} = K_\ell / P_\ell,$$

with one vertex for each component and effective couplings supplied by Schur reduction. Within component $\mathbb{C}$, choose an isolated localized spectral band and a weighted orthonormal frame

$$\Phi_C = (\phi_1, \dots, \phi_r), \quad \Phi_C^\dagger W_C \Phi_C = I_r.$$

The derived fiber is

$$E_C = \text{Ran } \Phi_C.$$

It need not be a harmonic space and therefore need not be supported by a hole. What it does require is a spectral gap, localization, and persistence. A candidate component is accepted only if all of the following remain stable across a stated range of scales:

- high modularity and low conductance relative to neighboring cuts;
- a localized spectral projector with stable rank;
- a nonzero band gap separating it from discarded modes;
- overlap with its predecessor and successor components;
- lifetime across multiple cobordism frames; and
- small external transport leakage.

This gives a self-similar hierarchy

$$\cdots \longrightarrow K_2 \longrightarrow K_1 \longrightarrow K_0$$

in which a vertex at one level resolves into a connected complex at the next. A fractal-like pattern is permitted but not required: the measured scaling of module count, volume, boundary size, and spectral gap decides whether the hierarchy is self-similar.

Community objectives supply principled cluster candidates [7], while network renormalization supplies tests for genuine self-similarity rather than visual resemblance [8]. The partition is therefore a measured part of the state: a recursively drawn pattern is not evidence of a fractal unless its scaling observables survive a refinement window.

6 Interactions and the expanding Hilbert space

For chain complexes $\mathbf{A}$ and $\mathbf{B}$, the graded tensor differential is the exact rule

$$d_{A \widehat{\otimes} B}(a \otimes b) = d_A a \otimes b + (-1)^{\deg a} a \otimes d_B b.$$

For a noninteracting product with product metric,

$$L_{A \widehat{\otimes} B} = L_A \otimes I + I \otimes L_B,$$




Figure 2: One recursive step. Persistent connected modules in K_ℓ become vertices of $K_{\ell+1}$, their external quadratic response is preserved by a Schur complement, and selected internal modes remain attached as fibers. A persistent supermodule can then be treated identically at the next scale.

so eigenvalues add and eigenvectors tensor. Connecting simplices contributed by a cobordism add a sparse interaction operator V_{AB} :

$$L_{AB} = L_A \otimes I + I \otimes L_B + V_{AB}.$$

At the fiber level, an interaction grows the carried space as

$$\mathcal{H}_{AB} = E_A \widehat{\otimes} E_B,$$

and a later interaction appends another factor. If J_C embeds an abstract state into the geometric carrier, exact amplitude preservation requires

$$J_C^\dagger W_C J_C = I.$$

Tensor products preserve isometry exactly. If $G = J_C^\dagger W_C J_C$ has Gram defect $\varepsilon = \|G - I\|$, then

$$|a^\dagger G b - a^\dagger b| \leq \varepsilon \|a\| \|b\|,$$

and two tensor factors obey

$$\varepsilon_{AB} \leq \varepsilon_A + \varepsilon_B + \varepsilon_A \varepsilon_B.$$

Thus the amplitude claim has an explicit, composable error budget.

Cobordism composition as a map between boundary state spaces is the organizing idea of topological field theory [9]; the general-boundary program makes the region/boundary assignment explicit

for quantum theory [10], and categorical quantum mechanics formalizes tensor composition and diagrammatic process semantics [11]. Tessera keeps only the parts that can be realized by its finite simplicial carrier and tests the resulting map numerically rather than assuming topological invariance.

7 A triangle carries the exact color algebra

Consider three oriented edge qubits around a triangle and interpret $|1\rangle$ as an occupied edge mode. Their graded tensor product is canonically the exterior algebra

$$(\mathbb{C}^2)^{\widehat{\otimes} 3} \cong \Lambda^\bullet \mathbb{C}^3 = \mathbf{1} \oplus \mathbf{3} \oplus \bar{\mathbf{3}} \oplus \mathbf{1}.$$

The sectors have occupation number $N = 0, 1, 2, 3$:

Sector	Basis dimension	Color interpretation	Fermion parity
$\Lambda^0 \mathbb{C}^3$	1	vacuum	even
$\Lambda^1 \mathbb{C}^3$	3	fundamental color triplet	odd
$\Lambda^2 \mathbb{C}^3$	3	antisymmetric anti-triplet	even
$\Lambda^3 \mathbb{C}^3$	1	color singlet	odd

Let $a_i^\dagger, a_i$ be the exterior creation and contraction operators. They satisfy the canonical anticommutation relations exactly:

$$\{a_i, a_j\} = 0, \quad \{a_i^\dagger, a_j^\dagger\} = 0, \quad \{a_i, a_j^\dagger\} = \delta_{ij}.$$

On the one-occupation sector, the bilinears

$$E_{ij} = a_i^\dagger a_j$$

satisfy

$$[E_{ij}, E_{kl}] = \delta_{jk} E_{il} - \delta_{il} E_{kj}.$$

The six Hermitian off-diagonal combinations together with

$$H_1 = E_{11} - E_{22}, \quad H_2 = \frac{E_{11} + E_{22} - 2E_{33}}{\sqrt{3}}$$

are the eight generators of $\mathfrak{su}(3)$. Thus the triangle does not merely hold three phases: its one-particle edge sector carries the fundamental representation, its two-particle sector carries the dual representation, and its traceless bilinears carry the adjoint octet.

The triplet description of quarks and the three-quark construction of baryons originate with the quark model [12]; the additional color triplet was introduced to resolve the statistics and state-counting problem [13, 14]. The claim here is narrower and new: Tessera's three oriented edge

modes would provide a geometric carrier of the same representation content, not a derivation of QCD from the combinatorics alone.




Figure 3: Exact representation content of three oriented edge qubits. The exterior sectors and their parity are algebraic identities. Interpreting the rank-three odd sector as quark color and the top wedge as a baryon color singlet is the physical hypothesis to be tested.

7.1 Geometric normalization

For triangle edge magnitudes ℓ_i and phases θ_i , define

$$c_i = \frac{\ell_i e^{i\theta_i}}{\sqrt{\ell_1^2 + \ell_2^2 + \ell_3^2}}, \quad |c\rangle = \sum_{i=1}^3 c_i |i\rangle.$$

Then $\langle c|c\rangle = 1$. Constraining the perimeter to one is a valid geometric scale gauge, but it is an L^1 condition and does not replace the L^2 Hilbert normalization. Normalized pure color states form $\mathbb{CP}^2$; $SU(3)$ is the transformation group, not the surface of the triangle itself.

7.2 The existing omega phase pattern

Let $\omega = e^{2\pi i/3}$. The exact Fourier frame

$$F_3 = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \end{pmatrix}$$

is unitary. The existing pattern $(1, \omega, \omega^2)/\sqrt{3}$ is therefore one color basis vector, not by itself the whole color fiber. Its cyclic orbit supplies an exact orthonormal triad.

8 Color transport and Wilson loops without a new gauge field

Let T_{AB} be the chain-level transfer already induced by connecting simplices from component B to component A . In local spectral frames, the raw fiber map is

$$M_{AB} = \Phi_A^\dagger W_A T_{AB} \Phi_B.$$

Its departure from an isometry is a physical leakage diagnostic:

$$\eta_{AB} = \|M_{AB}^\dagger M_{AB} - I\|.$$

When the selected band is isolated and η_{AB} is small, take the polar unitary and remove its determinant phase:

$$V_{AB} = M_{AB}(M_{AB}^\dagger M_{AB})^{-1/2}, \quad U_{AB} = \frac{V_{AB}}{(\det V_{AB})^{1/3}} \in SU(3).$$

Under a change of local spectral frame, $\Phi_A \mapsto \Phi_A g_A$ and $\Phi_B \mapsto \Phi_B g_B$,

$$U_{AB} \mapsto g_A^\dagger U_{AB} g_B.$$

Consequently the Wilson observable for a closed coarse loop γ is exactly gauge-invariant:

$$W(\gamma) = \frac{1}{3} \text{Tr} \prod_{(AB) \in \gamma} U_{AB}.$$

Polar normalization must never conceal a bad fiber assignment: every Wilson value is reported together with η_{AB} , the band gap, and the frame condition number.

This construction is the discrete spectral-frame analogue of Berry transport [15] and its non-Abelian Wilczek-Zee generalization [16]. Overlap matrices are also a standard route to gauge-covariant lattice observables [17]. Wilson loops themselves are foundational lattice gauge observables [18]. What is specific here is that the link matrix is not independently assigned: it is reconstructed from neighboring Hodge frames and accompanied by a leakage certificate.

9 Quarks as modular clusters

A *quark candidate* is proposed to be a component $\mathbf{Q}$ satisfying all of the following derived conditions:

1. $\mathbf{Q}$ is a persistent high-modularity cluster, not a prescribed region.
2. Its selected color fiber has stable rank three.
3. It occupies an odd exterior sector.
4. Its color transport has bounded leakage over its lifetime.
5. Its oriented world tube has one unit of quark-direction flux; reversing the tube yields the dual color representation and an antiquark.
6. Its total spectral fingerprint is stable under refinement and vertex relabeling.

The distinction between an antiquark and the $\Lambda^2 \mathbb{C}^3$ anti-triplet of two quarks is made by oriented world-tube flux and total occupation, not color alone. Assigning $B = +1/3$ to a forward odd quark tube and $B = -1/3$ to its orientation reverse makes pair creation conserve baryon number exactly.

Flavor and electric charge are not assumed as hidden labels. The conservative hypothesis is that two stable subclasses of the same cluster fiber provide an isospin doublet. On such a doublet, the measured orientation flux supplies baryon number and the standard relation

$$Q = I_3 + \frac{B}{2}$$

gives $Q_u = +2/3$ and $Q_d = -1/3$. This is a proposed identification, not yet a derivation: it succeeds only if an unlabeled two-dimensional spectral band emerges, is transported coherently, and its Gauss-flux readout agrees with those values.

10 Fermion statistics from simplicial orientation

The graded interchange law is

$$\tau(a \widehat{\otimes} b) = (-1)^{F_a F_b} b \widehat{\otimes} a.$$

Two odd clusters therefore acquire a minus sign on exchange, while an even composite does not. Parity adds modulo two:

Object	Odd constituents	Composite parity
quark or antiquark	1	odd
meson $q\bar{q}$	2	even
diquark qq	2	even
baryon qqq	3	odd

Pauli exclusion is the exact exterior-algebra identity

$$\|v_1 \wedge \cdots \wedge v_n\|^2 = \det[\langle v_i, v_j \rangle].$$

If two complete one-particle modes coincide, the determinant and the state vanish. The “complete” qualifier prevents double-counting signs: color, spin, flavor, space, and component support are wedged once as one mode. One must not multiply an extra fermion sign by the sign already present in the color epsilon tensor.

10.1 A label-independent exchange experiment

Let Φ_t be an orthonormal frame for the isolated odd subspace at frame t . Define the parallel transport

$$R_t = \text{polar}(\Phi_{t+1}^\dagger W_t \Phi_t), \quad U_\gamma = R_{T-1} \cdots R_0.$$

The determinant line gives the exchange character

$$\chi_F(\gamma) = \det U_\gamma.$$

The proposed fermion test is

$$\chi_F(\text{single exchange}) = -1, \quad \chi_F(\text{double exchange}) = +1.$$

Because the determinant is invariant under conjugation, this test is independent of the arbitrary

frame chosen inside the isolated band. A separate 2π rotation loop must also return -1 before the construction can claim a spin-statistics link.

Configuration-space topology already explains how exchange classes can carry quantum phases [19] and, in two dimensions, more general statistics [20]. The Tessera proposal uses this precedent only as a diagnostic template. The minus sign from the graded exterior algebra is exact; the claim that an actual geometric exchange cobordism realizes the corresponding determinant holonomy remains an experiment.

11 Fock space as an inductive limit of interactions

For M oriented fermionic edge modes,

$$\widehat{\bigotimes}_{m=1}^M \mathbb{C}^2 \cong \Lambda^\bullet \mathbb{C}^M = \bigoplus_{n=0}^M \Lambda^n \mathbb{C}^M,$$

and the dimension identity is exact:

$$2^M = \sum_{n=0}^M \binom{M}{n}.$$

Adding a new noninteracting mode uses the vacuum embedding

$$\iota_M : \mathcal{H}_M \hookrightarrow \mathcal{H}_{M+1}, \quad \iota_M(\psi) = \psi \widehat{\otimes} |0\rangle.$$

The infinite Fock space is the direct limit

$$\mathcal{F} = \varinjlim (\mathcal{H}_M, \iota_M).$$

This makes the infinite expansion precise without ever allocating an infinite array. At every finite simulation time only finitely many modes have interacted. Consistency requires

$$\|\iota_M U_M - U_{M+1} \iota_M\| \rightarrow 0$$

over a refinement sequence.

Bosonic gauge excitations need not add a new local oscillator. The traceless even bilinears

$$a_i^\dagger a_j - \frac{1}{3} \delta_{ij} N$$

span the color octet in $\mathbf{3} \otimes \bar{\mathbf{3}} = \mathbf{1} \oplus \mathbf{8}$ and have even fermion parity. Arbitrarily many such collective excitations are represented by adding more microscopic modes at finer resolution. The finite edge qubit remains unchanged.

This scale-by-scale state growth is adjacent to entanglement renormalization, where local Hilbert data are reorganized across layers before truncation [21]. The distinction is material: Tessera uses

a response-preserving Schur quotient plus an inductive vacuum embedding, and it must certify compatibility between successive finite spaces rather than assume a fixed bond dimension.

12 The proton as the maximally informative baryon

The proton is chosen because, beyond generic baryon structure, it demands a nontrivial flavor pattern, electric charge, spin, and experimentally meaningful form factors.

Let three persistent quark components A, B, C have color frames and normalized color columns c_A, c_B, c_C . The invariant color volume is

$$S_{ABC} = \epsilon_{ijk} c_A^i c_B^j c_C^k = \det[c_A \ c_B \ c_C].$$

Under a common $g \in SU(3)$, $S \mapsto \det(g)S = S$. Its squared magnitude is the Gram determinant

$$|S_{ABC}|^2 = \det(C^\dagger C) \in [0, 1].$$

The value one means the three color directions form an orthonormal frame. Their normalized wedge is then the unique $\Lambda^3 \mathbb{C}^3$ singlet. The proposed proton certificate is the conjunction:

- three persistent odd rank-three quark clusters;
- one persistent bound supercluster containing them;
- normalized color wedge with $|S_{ABC}|^2 \approx 1$ and vanishing net color flux;
- flavor spectrum with the uud occupation pattern;
- oriented baryon flux $B = 1$;
- Gauss flux $Q = +1$;
- total-space spin holonomy $J^2 = 3/4$ and $2\pi \mapsto -1$;
- finite radius and stable spectral mass/form-factor readouts; and
- stability of every dimensionless certificate under refinement.

None of these conditions should be included as an emergence target. The proton is found only if the base geometric optimization produces a component satisfying them. Targeted runs remain valuable as existence and obstruction experiments, but must be labeled as synthesis rather than emergence.

13 The master recursive construction

At every scale:

$$\begin{aligned} K_{\ell+1} &= K_\ell / P_\ell, \\ E_v^{\ell+1} &= \text{isolated localized spectral subspace of } C_v^\ell, \\ L_{\ell+1} &= \text{Schur}(L_\ell; P_\ell), \\ U_{vw}^{\ell+1} &= \text{SU3Polar}((\Phi_v^\ell)^\dagger W T_{vw} \Phi_w^\ell), \\ \mathcal{H}_{\ell+1} &= \widehat{\bigotimes}_v E_v^{\ell+1}. \end{aligned}$$

This one recursion supplies the effective geometry, the internal fiber, the derived connection, and




Figure 4: No-feedback emergence protocol. Only the base geometric objective drives optimization. Cluster, fiber, color, exchange, and baryon observables are computed from accepted snapshots and cannot feed back into the objective in an emergence run. Targeted synthesis is a separate, explicitly labeled mode.

the expanding state space. A high-level vertex can itself contain the same pattern at the next finer level.

14 Exactness and performance principles

The simulation should prefer an exact structural identity over a general dense numerical operation whenever both compute the same object:

- use sparse Schur solves, not explicit dense inverses;
- use Künneth sums for uncoupled product spectra, not diagonalization of the full Kronecker matrix;
- use exterior bit parity for exchange signs, not sampled phases;
- use exact 3×3 determinants and the fixed F_3 frame for color certificates;
- use analytic Regge/Hodge derivatives and Wirtinger gradients, not finite differences;
- use Smith normal form for integer homology and a spectral threshold only as a cross-check;
- use matrix-determinant/Woodbury updates for local cobordism changes;
- cache component factorizations and invalidate only affected stars;
- keep tensor products lazy and block-sparse by occupation/parity; and
- attach residual, gap, leakage, and condition-number certificates to every iterative eigensolve.

The exact route is not only faster. It prevents a numerical tolerance from becoming an undocumented physical postulate.

15 Prior art and boundary of novelty

No single cited work establishes the full recursive spectral-fiber proposal. The construction is a synthesis of several mature ideas, and its novelty should be evaluated at the joins. Table 3 states the boundary explicitly.

Table 3: Established ingredients and the additional Tessera claim.

Topic	Established prior art	Additional claim made here
Simplicial geometry and Hodge spectra	Regge curvature, discrete exterior calculus, and combinatorial Laplace spectra [2–4].	Use one jointly optimized Regge-Hodge complex as the only microscopic carrier for geometry and quantum readouts.
Coarse response and recursive modules	Kron/Schur reduction, certified graph reduction, modular communities, and self-similar network renormalization [5–8].	Treat a persistent component as a coarse vertex while retaining its localized Hodge band as a fiber, and repeat the operation recursively.
Geometric gauge transport	Berry and Wilczek-Zee holonomy, overlap-based lattice links, and Wilson loops [15–18].	Derive every $SU(3)$ link from overlap of rank-three component frames; assign no independent gauge link variable.
Color and fermion structure	Quark/color triplets and topological exchange phases [12–14, 19, 20].	Realize $\mathbf{1} \oplus \mathbf{3} \oplus \bar{\mathbf{3}} \oplus \mathbf{1}$ on three edge qubits, identify quarks with persistent odd modular fibers, and test exchange through determinant-line holonomy.
Scale composition and boundaries	TQFT cobordisms, general-boundary state assignments, categorical tensor composition, and entanglement renormalization [9–11, 21].	Build finite Fock stages from actual interaction cobordisms and require vacuum-embedding compatibility under refinement.
Spectral spacetime	Diffusion spectral dimension on ensembles of simplicial geometries [1].	Test whether many interacting Tessera cobordisms yield a stable four-dimensional spectral window while simultaneously supporting the particle certificates.

16 Falsification program

The formulation fails, or must be narrowed, if any of the following persists under refinement and tighter numerical certification:

1. **No persistent rank-three clusters.** High-modularity components appear, but their localized fiber rank or spectral gap is unstable.
2. **No faithful coarse response.** Schur-reduced components fail to reproduce measured external amplitudes within their stated residual.
3. **No derived gauge covariance.** Wilson values depend on the local spectral frame after leakage is controlled.
4. **No fermion holonomy.** A single cluster exchange does not give -1, or the sign changes under relabeling.
5. **No spinor rotation.** Exchange works but a 2π physical rotation does not give -1.
6. **No inductive compatibility.** Adding vacuum modes changes already-computed amplitudes by a nonvanishing amount.
7. **No unforced baryon.** Targeted synthesis can build the certificates, but the stationary geometric ensemble never produces them without a proton-specific term.




Figure 5: Relationship to prior art. The blue inputs are established research programs; the green center is the proposed Tesseract synthesis; the amber outputs are new physical identifications and must be validated independently. An arrow denotes conceptual inheritance, not a proof of the downstream claim.

8. **No continuum stability.** Dimensionless color, parity, charge, spin, and amplitude certificates drift rather than converge with refinement.

Holes may re-emerge and may correlate with some phases, but no claim in this paper depends on them doing so.

17 Conclusion

The proposed geometry is economical: a quark is a modular spectral component, its color is a three-edge exterior sector, its gauge transport is the overlap of local spectral frames, its fermionic sign is the grading forced by orientation, and its Fock space is the limit of the tensor factors created by interactions. Three such components form a baryon through their normalized color wedge; the proton is the sharpest test because it also demands the correct charge, flavor, and spin.

Much of the algebra works out exactly. What remains genuinely open is the important part: whether Tesseract's unforced Regge-Hodge dynamics actually produces the required persistent clusters and holonomies. That question is now a finite sequence of measurable implementation and validation tasks rather than a metaphor.

Prior-art references

- [1] Jan Ambjørn, Jerzy Jurkiewicz, and Renate Loll. Spectral dimension of the universe. *Physical Review Letters*, 95:171301, 2005. doi: 10.1103/PhysRevLett.95.171301. URL <https://arxiv.org/abs/hep-th/0505113>.
- [2] Tullio Regge. General relativity without coordinates. *Il Nuovo Cimento*, 19:558–571, 1961. doi: 10.1007/BF02733251.
- [3] Mathieu Desbrun, Anil N. Hirani, Melvin Leok, and Jerrold E. Marsden. Discrete exterior calculus, 2005. URL <https://arxiv.org/abs/math/0508341>.
- [4] Danijela Horak and Jürgen Jost. Spectra of combinatorial laplace operators on simplicial complexes. *Advances in Mathematics*, 244:303–336, 2013. doi: 10.1016/j.aim.2013.05.005. URL <https://arxiv.org/abs/1105.2712>.
- [5] Florian Dörfler and Francesco Bullo. Kron reduction of graphs with applications to electrical networks. *IEEE Transactions on Circuits and Systems I: Regular Papers*, 60(1):150–163, 2013. doi: 10.1109/TCSI.2012.2215780. URL <https://arxiv.org/abs/1102.2950>.
- [6] Andreas Loukas. Graph reduction with spectral and cut guarantees. *Journal of Machine Learning Research*, 20(116):1–42, 2019. URL <https://jmlr.org/papers/v20/18-680.html>.
- [7] Jörg Reichardt and Stefan Bornholdt. Statistical mechanics of community detection. *Physical Review E*, 74:016110, 2006. doi: 10.1103/PhysRevE.74.016110.
- [8] Chaoming Song, Shlomo Havlin, and Hernán A. Makse. Self-similarity of complex networks. *Nature*, 433:392–395, 2005. doi: 10.1038/nature03248.
- [9] Michael F. Atiyah. Topological quantum field theories. *Publications Mathématiques de l’IHÉS*, 68:175–186, 1988. doi: 10.1007/BF02698547.
- [10] Robert Oeckl. A “general boundary” formulation for quantum mechanics and quantum gravity. *Physics Letters B*, 575:318–324, 2003. doi: 10.1016/j.physletb.2003.08.043. URL <https://arxiv.org/abs/hep-th/0306025>.
- [11] Samson Abramsky and Bob Coecke. A categorical semantics of quantum protocols. In *Proceedings of the 19th Annual IEEE Symposium on Logic in Computer Science*, pages 415–425, 2004. doi: 10.1109/LICS.2004.1319636. URL <https://arxiv.org/abs/quant-ph/0402130>.
- [12] Murray Gell-Mann. A schematic model of baryons and mesons. *Physics Letters*, 8(3):214–215, 1964. doi: 10.1016/S0031-9163(64)92001-3.
- [13] Moo-Young Han and Yoichiro Nambu. Three-triplet model with double SU(3) symmetry. *Physical Review*, 139:B1006–B1010, 1965. doi: 10.1103/PhysRev.139.B1006.
- [14] O. W. Greenberg. Spin and unitary-spin independence in a paraquark model of baryons and mesons. *Physical Review Letters*, 13:598–602, 1964. doi: 10.1103/PhysRevLett.13.598.
- [15] Michael V. Berry. Quantal phase factors accompanying adiabatic changes. *Proceedings of the Royal Society of London A*, 392(1802):45–57, 1984. doi: 10.1098/rspa.1984.0023.

- [16] Frank Wilczek and A. Zee. Appearance of gauge structure in simple dynamical systems. *Physical Review Letters*, 52:2111–2114, 1984. doi: 10.1103/PhysRevLett.52.2111.
- [17] Takahiro Fukui, Yasuhiro Hatsugai, and Hiroshi Suzuki. Chern numbers in discretized brillouin zone: Efficient method of computing (spin) hall conductances. *Journal of the Physical Society of Japan*, 74(6):1674–1677, 2005. doi: 10.1143/JPSJ.74.1674. URL <https://arxiv.org/abs/cond-mat/0503172>.
- [18] Kenneth G. Wilson. Confinement of quarks. *Physical Review D*, 10:2445–2459, 1974. doi: 10.1103/PhysRevD.10.2445.
- [19] Michael G. G. Laidlaw and Cécile Morette DeWitt. Feynman functional integrals for systems of indistinguishable particles. *Physical Review D*, 3:1375–1378, 1971. doi: 10.1103/PhysRevD.3.1375.
- [20] Jon M. Leinaas and Jan Myrheim. On the theory of identical particles. *Il Nuovo Cimento B*, 37:1–23, 1977. doi: 10.1007/BF02727953.
- [21] Guifré Vidal. Entanglement renormalization. *Physical Review Letters*, 99:220405, 2007. doi: 10.1103/PhysRevLett.99.220405. URL <https://arxiv.org/abs/cond-mat/0512165>.

Repository evidence

- Tessera amplitude and obstruction results, [cobordism-results.md](#).
- Tessera spectral-dimension status, [h_ds4_status.md](#).
- Tessera interaction-history construction, [interaction-history-monte-carlo.md](#).
- Existing total-space spin obstruction, [joint_proton_spin_findings.md](#).
- Existing fixed-partition modularity implementation, [ModularityOptimizer.h](#).
- Current visualization and joint-stationarity experiment, [proton_animation.py](#).