A Multi-Dataset Audit of CARDIODIAG: Nested Evaluation, Cohort Transfer, and Uncertainty on Public Cardiovascular Benchmarks

Akhil Preetham E, Ch Hari Krishna, and G Vennela
Department of Computer Science and Engineering
Prasad V. Potluri Siddhartha Institute of Technology, Vijayawada, India
Author order, individual affiliations, contributions, and declarations require confirmation

Abstract—Public cardiovascular benchmarks can produce optimistic estimates when model selection, preprocessing, or cohort reuse is not controlled. We evaluated the existing CARDIODIAG pipeline on seven binary tasks with six models: Logistic Regression, K-Nearest Neighbors, Support Vector Machine, Random Forest, and the project's unchanged OVEC and OSEC ensembles. Four UCI angiographic coronary artery disease (CAD) cohorts (920 records total) were evaluated separately, with three additional tasks for heart-failure death, myocardial-infarction lethal complications, and SPECT imaging diagnosis. We used nested stratified 3-fold cross-validation with training-only preprocessing and calibration partitions; the official SPECTF split was retained. All 72 directed transfers among the four compatible CAD feature schemas completed with source-only selection. The highest observed within-task ROC-AUC ranged from 0.679 to 0.911 across CAD cohorts and was 0.805 on the SPECTF held-out test set. OVEC and OSEC did not consistently outperform individual baselines. Cross-cohort ROC-AUC ranged from 0.238 to 0.903 across model-transfer pairs. Empirical coverage for nominal 90% conformal sets ranged from 0.791 to 1.000, and source-versus-target feature-score AUROC varied from 0.357 to 0.934. These retrospective public-data experiments do not constitute independent clinical validation: the four UCI cohorts were represented in an earlier project composite, and participant-level overlap cannot be ruled out. Results support a reproducible benchmark audit, not clinical utility or deployment readiness.

Index Terms—cardiovascular prediction, coronary artery disease, nested cross-validation, dataset shift, calibration, conformal prediction, reproducibility

I. Introduction

High internal discrimination is not sufficient evidence that a clinical prediction model is reliable at another site. Reuse of small public datasets, data-dependent preprocessing outside validation folds, and endpoint ambiguity can inflate estimates or weaken their clinical meaning. Reporting guidance and risk-of-bias frameworks therefore emphasize transparent data definitions, prediction timing, validation design, and model evaluation [1], [2]. Nested resampling is needed when the same development data support hyperparameter selection and performance estimation [3], [4].

CARDIODIAG began as a feature-reduction and ensemble approach for heart-disease classification [5]. Previous project audits identified historical result artifacts, data-lineage concerns, and evaluation-boundary issues. This study preserves the established optimized voting classifier (OVEC) and stacking classifier (OSEC), then compares them with individual baselines using a leakage-controlled pipeline. The aims are to: (1) compare models across compatible and task-distinct public datasets without pooling outcomes; (2) assess discrimination, calibration, overfitting diagnostics, conformal coverage, selective risk, and feature-support alerts; and (3) measure source-only transfer among four schema-compatible CAD cohorts.

II. Datasets and Study Design

A. Dataset selection and endpoints

The UCI Heart Disease collection contains four historical cohorts, with 13 shared clinical predictors and the 'num' angiographic disease field. We mapped 'num=0' to negative and 'num=1…4' to positive, following the repository's established binary rule [6]. The four component cohorts total 920 records. They are not independent replication datasets for this project: the same cohorts were represented in the previously examined Heart5 composite. Exact predictor-row matching found no cross-cohort exact duplicates, but no persistent participant identifiers were available to rule out transformed or near-duplicate records.

We also evaluated UCI Heart Failure Clinical Records (299 records), UCI Myocardial Infarction Complications (1,700 records), and UCI SPECTF Heart (267 records) [7], [8], [9]. These were kept as separate tasks. For the heart-failure task, follow-up time was excluded as a predictor and death event was the target; follow-up horizons vary. For MI complications, the target was any lethal outcome and only admission-available variables were retained after excluding identifiers, later variables, and outcome proxies. SPECTF uses 44 SPECT-derived features and an overall normal/abnormal diagnosis; it is not a clinical CAD endpoint. The official 80-record training / 187-record test split was preserved, including 172 positive-coded and 15 negative-coded test records.

All source records remain separate. UCI dataset pages list CC BY 4.0; source URLs, archive checksums, class counts, missingness, and decisions for screened candidates are included in the accompanying dataset manifest and compatibility table. UCI Statlog Heart was excluded as

a new cohort because the project's exact matching audit found all 270 harmonized records match Cleveland [10]. Previously examined cohorts were not presented as new confirmation.

B. Evaluation protocol

For the six tabular cohorts evaluated with resampling, we used stratified three-fold outer validation and stratified three-fold inner model selection (seed 20261108). Within each outer training partition, 20% was reserved for conformal and feature-support threshold calibration. Hyperparameter search used only the remaining development subset. Each outer test fold was held out until scoring. SPECTF used the official test partition; its training set was separated into development and calibration subsets only.

All imputation, scaling, categorical encoding, and outlier replacement were fitted within each pipeline's training data. The existing 'TrainingOnlyCleaner' replaces values outside training-fitted 1st/99th percentiles with training medians; numeric values are median-imputed and scaled; categorical values are most-frequent imputed and one-hot encoded. No SMOTE was applied. The corrected pipeline does not use a feature selector in this study. Consequently, no global feature selection or preprocessing was performed before splitting.

The individual model grids were logistic-regression $C \in \{0.1, 1, 10\}$; KNN $k \in \{3, 5, 7\}$ and uniform/distance weighting; SVM $C \in \{0.1, 1, 10\}$ and linear/RBF kernels; and random-forest square-root/all-feature subsampling and leaf size 1/3. OVEC and OSEC retained their existing definitions. OVEC uses the project's soft-voting constituents (KNN, random forest, balanced logistic regression, and balanced SVC). OSEC retains its stacking architecture; its final estimator was tuned over $C \in \{0.1, 1, 10\}$.

C. Metrics and uncertainty

At a fixed probability threshold of 0.5, we report accuracy, precision, sensitivity, specificity, F1, ROC-AUC, and average precision (PR-AUC). Brier score and ten-bin equal-width expected calibration error (ECE), both macro-bin and sample-weighted, summarize calibration. Fold means and standard deviations describe nested evaluation. For pooled held-out predictions, 1,000 class-stratified percentile bootstrap replicates condition on observed class counts; these intervals do not account for patient clustering, cohort selection, or between-site variation. No hypothesis tests or superiority claims were prespecified.

Split-conformal sets use nonconformity $1 - p(y)$ and nominal error level $\alpha = 0.10$. Feature support is the mean distance to five nearest development observations after the fitted feature transformation; the alert cutoff is the 95th percentile of the source calibration scores. Cross-dataset domain AUROC distinguishes source calibration scores from target scores and is a feature-shift diagnostic, not an outcome-prediction metric. Confidence-ranked risk–coverage curves and the joint singleton-conformal / in-support accepted subset are descriptive. Conformal coverage guarantees require exchangeability and are not assumed to hold after transport [11].

III. Results

A. Within-task model comparison

All 42 dataset–model evaluations completed. Table II shows outer-fold ROC-AUC means and standard deviations; SPECTF is one official held-out test estimate and has no fold SD. The highest observed model by fold-mean ROC-AUC varied across tasks: random forest on Cleveland (0.911), OVEC on Hungary (0.890), logistic regression on Switzerland (0.772), random forest on VA Long Beach (0.679), OVEC on heart failure (0.752), random forest on MI complications (0.869), and OVEC on SPECTF (0.805). These maxima are descriptive comparisons, not unbiased estimates of a separately selected final model.

The complete thresholded and ranking results for each model appear in Tables III and IV; all metrics are also available as CSV. OVEC exceeded OSEC ROC-AUC on Cleveland, Hungary, heart failure, SPECTF, and VA Long Beach, while OSEC was higher on Switzerland and MI. Neither ensemble consistently exceeded all four individual baselines. Model selection results therefore do not support a general ensemble-superiority claim.

B. Overfitting, calibration, and selective diagnostics

Resubstitution scores were often higher than outer-fold performance, particularly for KNN and random forest. For example, MI-complication training ROC-AUC was 1.000 for KNN and 1.000 (approximately) for random forest, while outer-fold means were 0.736 and 0.869. This is evidence of a training–evaluation gap in these fits, not proof that every high-score model is overfit. The full report includes model-specific training, selected inner-validation, outer-test values, and per-fold gaps.

Empirical coverage for nominal 90% split-conformal sets varied from 0.791 to 1.000 across dataset–model groups; therefore the target was not uniformly met on these held-out samples. Average set sizes and singleton rates varied as well. Confidence-ranked selective coverage did not identify a universally reliable error-free region. The combined singleton-plus-support rule accepted between 0% and 88.5% of observations across task–model groups. One SPECTF model accepted none, while low-coverage zero-error subsets in other groups remain unstable and cannot certify safety.

The nearest-neighbor alert rate varied from 0.5% to 37.4% within held-out task predictions. Across cross-cohort transfers, source-versus-target feature-score AUROC ranged from 0.357 to 0.934, showing that feature-support separation depended strongly on source, target, and model. This distance score may detect some shifts in

TABLE I
Dataset characteristics. CAD cohorts share the UCI angiographic endpoint; heart failure, MI, and SPECTF are different tasks. Class counts are negative/positive.

Dataset	n	Predictors	Negative/positive	Missing cells	Task
UCI Cleveland	303	13	164/139	6	Legacy angiographic CAD cohort; retrospective s...
UCI Hungary	294	13	188/106	782	Legacy angiographic CAD cohort; retrospective s...
UCI Switzerland	123	13	8/115	396	Legacy angiographic CAD cohort; retrospective s...
UCI VA Long Beach	200	13	51/149	748	Legacy angiographic CAD cohort; retrospective s...
UCI Heart Failure	299	11	203/96	0	Separate death-event task; varying follow-up ho...
UCI MI Complications	1700	100	1429/271	15200	Separate MI lethal-outcome task; admission vari...
UCI SPECTF Heart	267	44	55/212	0	Separate SPECT normal/abnormal task; official 8...

TABLE II
ROC-AUC across all six candidates. Values are fold mean ± fold SD for nested evaluation. SPECTF reports the single official held-out test estimate.

Dataset	LR	KNN	SVM	RF	OVEC	OSEC
UCI Cleveland	0.911 ± 0.031	0.889 ± 0.031	0.909 ± 0.039	0.911 ± 0.032	0.909 ± 0.033	0.903 ± 0.029
UCI Hungary	0.879 ± 0.040	0.876 ± 0.020	0.877 ± 0.043	0.880 ± 0.042	0.890 ± 0.033	0.876 ± 0.031
UCI Switzerland	0.772 ± 0.097	0.530 ± 0.120	0.569 ± 0.312	0.736 ± 0.096	0.732 ± 0.125	0.751 ± 0.183
UCI VA Long Beach	0.623 ± 0.026	0.641 ± 0.046	0.633 ± 0.036	0.679 ± 0.037	0.655 ± 0.041	0.612 ± 0.023
UCI Heart Failure	0.735 ± 0.039	0.661 ± 0.020	0.565 ± 0.324	0.739 ± 0.037	0.752 ± 0.027	0.736 ± 0.041
UCI MI Complications	0.854 ± 0.006	0.736 ± 0.023	0.846 ± 0.017	0.869 ± 0.008	0.866 ± 0.006	0.868 ± 0.009
UCI SPECTF Heart	0.758	0.729	0.788	0.796	0.805	0.797

TABLE III
Complete classification metrics for the four UCI CAD cohorts. Each value is outer-fold mean ± SD; PR-AUC should be read alongside cohort prevalence because precision depends on case mix.

Dataset	Model	Acc.	Prec.	Sens.	Spec.	F1	ROC-AUC	PR-AUC
UCI Cleveland	K-Nearest Neighbors	0.809 ± 0.029	0.819 ± 0.033	0.748 ± 0.033	0.860 ± 0.029	0.782 ± 0.032	0.889 ± 0.031	0.873 ± 0.027
UCI Cleveland	Logistic Regression	0.812 ± 0.026	0.797 ± 0.054	0.798 ± 0.034	0.823 ± 0.066	0.796 ± 0.018	0.911 ± 0.031	0.909 ± 0.030
UCI Cleveland	OSEC	0.825 ± 0.055	0.829 ± 0.091	0.791 ± 0.033	0.853 ± 0.099	0.808 ± 0.047	0.903 ± 0.029	0.901 ± 0.024
UCI Cleveland	OVEC	0.818 ± 0.035	0.808 ± 0.063	0.799 ± 0.024	0.835 ± 0.067	0.802 ± 0.029	0.909 ± 0.033	0.902 ± 0.031
UCI Cleveland	Random Forest	0.802 ± 0.030	0.795 ± 0.071	0.777 ± 0.035	0.823 ± 0.084	0.783 ± 0.016	0.911 ± 0.032	0.902 ± 0.031
UCI Cleveland	Support Vector Machine	0.828 ± 0.049	0.822 ± 0.073	0.806 ± 0.019	0.847 ± 0.078	0.813 ± 0.043	0.909 ± 0.039	0.908 ± 0.031
UCI Hungary	K-Nearest Neighbors	0.813 ± 0.021	0.790 ± 0.113	0.688 ± 0.106	0.882 ± 0.089	0.725 ± 0.018	0.876 ± 0.020	0.816 ± 0.039
UCI Hungary	Logistic Regression	0.793 ± 0.041	0.715 ± 0.086	0.735 ± 0.068	0.824 ± 0.091	0.720 ± 0.029	0.879 ± 0.040	0.822 ± 0.077
UCI Hungary	OSEC	0.806 ± 0.018	0.705 ± 0.034	0.802 ± 0.031	0.808 ± 0.044	0.749 ± 0.008	0.876 ± 0.031	0.821 ± 0.049
UCI Hungary	OVEC	0.799 ± 0.039	0.749 ± 0.087	0.688 ± 0.052	0.861 ± 0.077	0.713 ± 0.031	0.890 ± 0.033	0.837 ± 0.055
UCI Hungary	Random Forest	0.803 ± 0.046	0.748 ± 0.092	0.707 ± 0.093	0.856 ± 0.086	0.721 ± 0.053	0.880 ± 0.042	0.824 ± 0.062
UCI Hungary	Support Vector Machine	0.796 ± 0.047	0.759 ± 0.102	0.660 ± 0.052	0.872 ± 0.085	0.702 ± 0.043	0.877 ± 0.043	0.839 ± 0.057
UCI Switzerland	K-Nearest Neighbors	0.935 ± 0.014	0.935 ± 0.014	1 ± 0	0 ± 0	0.966 ± 0.007	0.530 ± 0.120	0.937 ± 0.026
UCI Switzerland	Logistic Regression	0.846 ± 0.061	0.938 ± 0.026	0.897 ± 0.092	0.167 ± 0.289	0.914 ± 0.038	0.772 ± 0.097	0.981 ± 0.011
UCI Switzerland	OSEC	0.902 ± 0.049	0.940 ± 0.030	0.956 ± 0.031	0.167 ± 0.289	0.948 ± 0.027	0.751 ± 0.183	0.976 ± 0.022
UCI Switzerland	OVEC	0.935 ± 0.014	0.935 ± 0.014	1 ± 0	0 ± 0	0.966 ± 0.007	0.732 ± 0.125	0.974 ± 0.017
UCI Switzerland	Random Forest	0.894 ± 0.037	0.941 ± 0.026	0.948 ± 0.068	0.167 ± 0.289	0.943 ± 0.022	0.736 ± 0.096	0.975 ± 0.015
UCI Switzerland	Support Vector Machine	0.935 ± 0.014	0.935 ± 0.014	1 ± 0	0 ± 0	0.966 ± 0.007	0.569 ± 0.312	0.945 ± 0.061
UCI VA Long Beach	K-Nearest Neighbors	0.715 ± 0.013	0.759 ± 0.008	0.906 ± 0.041	0.157 ± 0.068	0.826 ± 0.013	0.641 ± 0.046	0.815 ± 0.027
UCI VA Long Beach	Logistic Regression	0.610 ± 0.030	0.788 ± 0.008	0.651 ± 0.050	0.490 ± 0.034	0.713 ± 0.031	0.623 ± 0.026	0.812 ± 0.032
UCI VA Long Beach	OSEC	0.650 ± 0.061	0.845 ± 0.042	0.658 ± 0.143	0.627 ± 0.180	0.731 ± 0.080	0.612 ± 0.023	0.805 ± 0.027
UCI VA Long Beach	OVEC	0.720 ± 0.021	0.760 ± 0.012	0.913 ± 0.046	0.157 ± 0.090	0.829 ± 0.017	0.655 ± 0.041	0.818 ± 0.039
UCI VA Long Beach	Random Forest	0.730 ± 0.068	0.818 ± 0.017	0.819 ± 0.100	0.471 ± 0.059	0.817 ± 0.055	0.679 ± 0.037	0.838 ± 0.027
UCI VA Long Beach	Support Vector Machine	0.755 ± 0.008	0.753 ± 0.006	1 ± 0	0.039 ± 0.034	0.859 ± 0.004	0.633 ± 0.036	0.816 ± 0.026

these samples, but does not provide a consistent rejection rule. Figure 2 shows calibration behavior; the complete risk–coverage curves are included in the accompanying experimental report.

C. Source-only cross-cohort transfers

All 72 directed model transfers across the four UCI CAD cohorts completed. Source-only tuning and calibration were used; target labels were used only for evaluation. Across model–direction pairs, target ROC-AUC ranged from 0.238 to 0.903. The average ROC-AUC across models varied by direction from 0.538 (Switzerland to VA Long Beach) to 0.886 (Cleveland to Hungary). The direction-level averages summarize model variability and should not obscure the full per-model matrix. These are retrospective stress tests because the component cohorts were part of the previously examined Heart5 lineage, and no patient identifier permits an independentness guarantee.

D. Failed analyses

Six learning-curve configurations failed, all for Switzerland. At the smallest requested training size, one or more stratified training subsets contained only one class; ROC-AUC learning curves were therefore not calculated for any model on that cohort. The remaining 36 task–model learning curves completed. All nested model evaluations

TABLE IV
Complete classification metrics for the task-distinct heart-failure, MI-complication, and SPECTF datasets. SPECTF has one official test split and no fold SD.

Dataset	Model	Acc.	Prec.	Sens.	Spec.	F1	ROC-AUC	PR-AUC
UCI Heart Failure	K-Nearest Neighbors	0.685 ± 0.034	0.508 ± 0.125	0.229 ± 0.095	0.901 ± 0.009	0.314 ± 0.115	0.661 ± 0.020	0.457 ± 0.033
UCI Heart Failure	Logistic Regression	0.682 ± 0.046	0.508 ± 0.055	0.656 ± 0	0.695 ± 0.068	0.572 ± 0.036	0.735 ± 0.039	0.580 ± 0.073
UCI Heart Failure	OSEC	0.719 ± 0.033	0.554 ± 0.049	0.656 ± 0.054	0.749 ± 0.043	0.600 ± 0.044	0.736 ± 0.041	0.556 ± 0.018
UCI Heart Failure	OVEC	0.706 ± 0.015	0.559 ± 0.033	0.396 ± 0.036	0.852 ± 0.014	0.463 ± 0.033	0.752 ± 0.027	0.575 ± 0.024
UCI Heart Failure	Random Forest	0.696 ± 0.013	0.534 ± 0.019	0.396 ± 0.100	0.838 ± 0.038	0.450 ± 0.074	0.739 ± 0.037	0.544 ± 0.043
UCI Heart Failure	Support Vector Machine	0.679 ± 0.002	0 ± 0	0 ± 0	1 ± 0	0 ± 0	0.565 ± 0.324	0.455 ± 0.212
UCI MI Complications	K-Nearest Neighbors	0.851 ± 0.004	0.798 ± 0.189	0.096 ± 0.025	0.994 ± 0.006	0.169 ± 0.038	0.736 ± 0.023	0.415 ± 0.039
UCI MI Complications	Logistic Regression	0.807 ± 0.010	0.438 ± 0.017	0.727 ± 0.026	0.822 ± 0.017	0.546 ± 0.008	0.854 ± 0.006	0.565 ± 0.030
UCI MI Complications	OSEC	0.799 ± 0.007	0.428 ± 0.009	0.768 ± 0.021	0.805 ± 0.012	0.550 ± 0.003	0.868 ± 0.009	0.615 ± 0.014
UCI MI Complications	OVEC	0.874 ± 0.006	0.686 ± 0.016	0.380 ± 0.039	0.967 ± 0.001	0.489 ± 0.036	0.866 ± 0.006	0.597 ± 0.011
UCI MI Complications	Random Forest	0.869 ± 0.012	0.829 ± 0.064	0.236 ± 0.114	0.990 ± 0.009	0.356 ± 0.124	0.869 ± 0.008	0.625 ± 0.015
UCI MI Complications	Support Vector Machine	0.862 ± 0.006	0.638 ± 0.054	0.321 ± 0.040	0.965 ± 0.010	0.426 ± 0.036	0.846 ± 0.017	0.537 ± 0.023
UCI SPECTF Heart	K-Nearest Neighbors	0.529	0.988	0.494	0.933	0.659	0.729	0.964
UCI SPECTF Heart	Logistic Regression	0.738	0.949	0.756	0.533	0.841	0.758	0.975
UCI SPECTF Heart	OSEC	0.733	0.969	0.733	0.733	0.834	0.797	0.980
UCI SPECTF Heart	OVEC	0.749	0.970	0.750	0.733	0.846	0.805	0.980
UCI SPECTF Heart	Random Forest	0.743	0.956	0.756	0.600	0.844	0.796	0.979
UCI SPECTF Heart	Support Vector Machine	0.727	0.948	0.744	0.533	0.834	0.788	0.978

TABLE V
Ranges across models for pooled Brier/ECE, mean empirical conformal coverage and set size, and combined singleton/support accepted coverage. These ranges summarize heterogeneous models; they are not model-selection recommendations.

Dataset	Brier range	Weighted ECE range	Conformal coverage range	Set size range	Joint coverage range
UCI Cleveland	0.120–0.131	0.050–0.083	0.894–0.931	1.201–1.290	0.680–0.762
UCI Heart Failure	0.184–0.219	0.051–0.173	0.890–0.910	1.418–1.616	0.381–0.555
UCI Hungary	0.131–0.142	0.037–0.091	0.881–0.918	1.156–1.296	0.680–0.820
UCI MI Complications	0.095–0.147	0.017–0.212	0.905–0.922	1.086–1.347	0.624–0.885
UCI SPECTF Heart	0.168–0.329	0.233–0.455	0.791–1.000	1.128–2.000	0.000–0.866
UCI Switzerland	0.062–0.145	0.017–0.286	0.951–0.976	1.154–1.846	0.138–0.707
UCI VA Long Beach	0.182–0.236	0.047–0.216	0.900–0.970	1.595–1.843	0.155–0.380

TABLE VI
Source-only transfer summary over the six models per direction. Mean and range summarize model scores on the same target cohort; they are descriptive. Domain OOD AUROC compares source calibration and target feature scores.

Source	Target	Mean AUC	Model range	Domain OOD AUROC	Target alert
UCI Cleveland	UCI Hungary	0.886	0.859–0.903	0.486	0.074
UCI Cleveland	UCI Switzerland	0.713	0.639–0.741	0.560	0.045
UCI Cleveland	UCI VA Long Beach	0.719	0.692–0.733	0.703	0.129
UCI Hungary	UCI Cleveland	0.843	0.820–0.873	0.896	0.421
UCI Hungary	UCI Switzerland	0.756	0.688–0.800	0.723	0.123
UCI Hungary	UCI VA Long Beach	0.713	0.689–0.739	0.727	0.160
UCI Switzerland	UCI Cleveland	0.635	0.534–0.773	0.874	0.612
UCI Switzerland	UCI Hungary	0.761	0.654–0.860	0.571	0.122
UCI Switzerland	UCI VA Long Beach	0.538	0.484–0.578	0.454	0.107
UCI VA Long Beach	UCI Cleveland	0.706	0.238–0.851	0.890	0.448
UCI VA Long Beach	UCI Hungary	0.813	0.702–0.868	0.772	0.235
UCI VA Long Beach	UCI Switzerland	0.662	0.610–0.711	0.586	0.046

and all cross-cohort transfers completed. Switzerland contains only eight negative records, so even the completed three-fold performance and OSEC stacking estimates have high uncertainty; the fixed five-fold internal stacking procedure generated minority-count warnings. The failures and warnings are preserved in the logs and failure CSV.

IV. Discussion

This study broadens the CARDIODIAG audit while preserving the model definitions and keeping outcomes separate. In the compatible CAD cohorts, internal scores were strongest on Cleveland and Hungary and substantially weaker on VA Long Beach; Switzerland had only eight negative cases and yielded unstable fold estimates. The cross-cohort results varied sharply by direction and model. Even where the transfer AUC remained high, historical dataset reuse and unavailable participant linkage prevent treating these experiments as independent external validation.

OVEC and OSEC did not dominate the individual baselines consistently. Calibration and selective diagnostics were also mixed: nominal conformal coverage fell below 90% in some model–task combinations, feature-support alerts varied widely, and low-coverage accepted subsets are not evidence of safe abstention. High apparent PR-AUC on imbalanced tasks must be interpreted against




Fig. 1. Outer-fold mean ROC-AUC and PR-AUC with fold standard deviations. The SPECTF point is from its official test set and has no fold-based interval.

prevalence and the target definition; it does not by itself establish clinical value.

The additional public datasets illustrate why dataset count alone does not establish evidence strength. Heart-failure death and MI lethal complications are distinct outcomes with different prediction times. SPECTF is an imaging classification task and retains a severe class imbalance in its official test set. These tasks are useful for checking implementation behavior but cannot validate angiographic CAD prediction.

A. Limitations

The datasets are small, historical, and in some cases severely imbalanced or incomplete. Exact-row matching cannot detect near duplicates or transformations, while persistent participant identifiers are unavailable. The source tables do not establish a prospective index prediction time. Patient clustering, site sampling, and between-site variation are not represented by the class-stratified bootstrap intervals. Switzerland learning curves were not estimable under the requested training fractions, and OSEC's fixed internal stacking folds are poorly matched to such a small minority class. Conformal exchangeability is not established under dataset shift. No prospective, real-world, fairness, clinical-utility, or deployment study was performed.

V. Conclusion

The experiments provide a traceable comparison of individual classifiers and the existing CARDIODIAG ensembles across seven separately defined public tasks. No ensemble was consistently superior, cross-cohort performance varied substantially, and conformal/OOD/selective diagnostics did not produce a uniform reliability guarantee. These results support reproducible software and retrospective benchmark analysis. They do not establish independent clinical validation or readiness for use. A confirmatory study requires an authorized, untouched cohort with a prespecified CAD reference standard, prediction time, patient-level overlap controls, and a frozen operating protocol.

Declarations

Ethics and data governance: Only public UCI repository datasets were used in this run. No institutional approval or




Fig. 2. Calibration curves from pooled held-out predictions. Bin counts and ECE variants are available in the result tables.




Fig. 3. Mean target ROC-AUC over six source-only models for each CAD cohort direction. Source and target are historically examined UCI cohorts; diagonal cells are omitted.

exemption is claimed; the project team should obtain the appropriate institutional determination before submission.
Funding and competing interests: Confirmation pending.
Author contributions: Contributions and author order require confirmation from all listed authors.
Data and code availability: Raw records and row-level predictions are not included in the public release. Aggregate metrics, source checksums, preprocessing scripts, and figures are included in the project deliverables. Dataset attribution is required under the listed UCI licenses.
Manuscript status: Working draft requiring author, affiliation, declaration, and venue review.

References

[1] G. S. Collins, K. G. M. Moons, P. Dhiman et al., "Tripod+ai statement: updated guidance for reporting clinical prediction models that use regression or machine learning methods," BMJ, vol. 385, p. e078378, 2024.

[2] K. G. M. Moons, J. A. A. Damen, T. Kaul et al., "Probast+ai: an updated quality, risk of bias, and applicability assessment tool for prediction models using regression or artificial intelligence methods," BMJ, vol. 388, p. e082505, 2025.

[3] S. Varma and R. Simon, "Bias in error estimation when using cross-validation for model selection," BMC Bioinformatics, vol. 7, p. 91, 2006.

[4] G. C. Cawley and N. L. C. Talbot, "On over-fitting in model selection and subsequent selection bias in performance evaluation," Neural Computation, vol. 22, no. 11, pp. 2818–2837, 2010.

[5] M. J. Khan, B. Raza, and A. Nadeem, "Cardiodiag: a machine learning approach for cardiovascular disease diagnosis using xai-guided hybrid dimensionality reduction and optimized-based ensemble classifiers," PeerJ Computer Science, vol. 12, p. e4059, 2026.

[6] A. Janosi, W. Steinbrunn, M. Pfisterer, and R. Detrano, "Heart Disease," UCI Machine Learning Repository, dataset 45, 1989. [Online]. Available: https://archive.ics.uci.edu/dataset/45/heart%2Bdisease

[7] UCI Machine Learning Repository, "Heart Failure Clinical Records," Dataset 519, cC BY 4.0; accessed 2026-11-08. [Online]. Available: https://archive.ics.uci.edu/dataset/519/heart%2Bfailure%2Bclinical%2Brecords

[8] ——, "Myocardial Infarction Complications," Dataset 579, cC BY 4.0; accessed 2026-11-08. [Online]. Available: https://archive.ics.uci.edu/dataset/579/myocardial%2Binfarction%2Bcomplications

[9] ——, "SPECTF Heart," Dataset 96, cC BY 4.0; accessed 2026-11-08. [Online]. Available: https://archive.ics.uci.edu/dataset/96/spectf%2Bheart

[10] ——, "Statlog (Heart)," Dataset 145, cC BY 4.0; all 270 rows match Cleveland after binary-label harmonization. [Online]. Available: https://archive.ics.uci.edu/dataset/145/statlog+heart

[11] A. N. Angelopoulos and S. Bates, "Conformal prediction: A gentle introduction," Foundations and Trends in Machine Learning, vol. 16, no. 4, pp. 494–591, 2023.