THE STUDY OF CROP YIELD PREDICTION USING VARIOUS MACHINE LEARNING ALGORITHMS

AKHIL EDWIN

A dissertation submitted in partial fulfilment of the requirements for the degree
of Master of science in Business Analytics
at Dublin Business School

Supervisor: PIERPAOLO DONDIO

Declaration

I declare that, excluding the unique content in this dissertation, which has not been previously presented for any academic degree, all aspects of this work—unless explicitly cited within the text and Acknowledgements—are entirely my own creation. I affirm that no collaborative efforts were employed in the development of any section of this dissertation.

Signed : AKHIL EDWIN

Student Number : 10636297

Date : 08-01-24

Abstract

This project focuses on applying diverse machine-learning models for crop yield prediction in agriculture. Leveraging a dataset encompassing variables such as weather conditions, pesticide usage, and crop types, six models—Long Short-Term Memory (LSTM), Convolutional Neural Network (CNN), Random Forest, Gradient Boosting Machine (GBM), Support Vector Machine (SVM), and Decision Tree—were evaluated based on Root Mean Square Error (RMSE). The results showcase the models' proficiency in capturing different aspects of agricultural data, with low RMSE values suggesting accurate predictions. LSTM and CNN excel in handling sequential and spatial data, respectively, while Random Forest and GBM demonstrate strong ensemble learning capabilities. SVM and decision trees exhibit robust performances and interpretability in high-dimensional spaces. The findings contribute to understanding each model's strengths in the context of crop yield prediction.

Acknowledgements

I express my gratitude to all who contributed to the study of crop yield prediction using diverse machine learning algorithms. Special thanks to my advisors for their guidance, the research team for their collaboration, and the academic community for fostering an environment of learning. This study stands as a collective effort, and I appreciate the support and insights received throughout the research process.

Table of contents

THE STUDY OF CROP YIELD PREDICTION USING VARIOUS MACHINE LEARNING ALGORITHMS 1
Abstract 2
CHAPTER 1 INTRODUCTION 6
1.1 Background 6
1.2 Research problem statement 6
1.3 Research objectives 7
1.4 Research Methodologies 7
1.5 Scope and Limitations 8
1.6 Document Outline 8
CHAPTER 2 LITERATURE REVIEW AND RELATED WORK 10
CHAPTER 3 DESIGN AND METHODOLOGY 23
3.1 Crisp dm methodology 23
3.1 Dataset Exploration: 24
3.2 Architecture 25
3.3 Data Preprocessing 26
3.3.1 Top 10 Areas based on rainfall 26
3.4 Ethics: 29
3.5 Machine Learning Algorithms 30
3.5.1 Random Forest : 30
3.5.2 Decision Tree 30
3.5.3 Support Vector Machine 31
3.5.4 Gradient Boosting algorithm 31
3.5.5 CNN 32
3.5.6 LSTM 32
CHAPTER 4 EVALUATION OF METRICS AND VALUES 34
1. LSTM: 34
2. CNN: 34
3. Random Forest: 34
4. GBM: 34
5. SVM: 34
6. Decision Tree: 34
CHAPTER 5 CONCLUSION AND FUTURE WORK 36
5.1 Research Overview 36
5.2 Problem Definition 36
5.3 Design/Experimentation, Evaluation & Results 36
5.4 Contributions and impact 37
5.5 Future Work & recommendations 37
References: 38

Table of figures

Figure 1 *Crisp Dm process* 23
Figure 2 *Architecture* 25
Figure 3 *top 10 areas based on rainfall* 26
Figure 4 *Highest yield crops* 27
Figure 5 *Top areas with good yield* 27
Figure 6 *rainfall* 28
Figure 7 *Average distributed temperature* 28
Figure 8 *Corrleation matrix* 29
Figure 9 *Comparision of results* 35
Figure 10 *Evaluation of results* 37

Table of tables

Table 1 *Literature study* 22

CHAPTER 1 INTRODUCTION

1.1 Background

In the realm of agricultural science and data-driven decision-making, this study investigates the critical area of crop yield prediction, with a specific focus on the agricultural landscape of Albania. Agriculture is a pivotal component of numerous economies, and the accurate forecasting of crop yields holds significant implications for sustainable farming practices and food security. This research marks the convergence of advanced data analytics and agricultural expertise, aiming to employ machine learning algorithms to unravel the complex interplay of factors impacting crop productivity.

The dataset under examination comprises a comprehensive array of information, including historical crop yields (measured in hg/ha), average annual rainfall (in mm), quantities of pesticides used (in tonnes), and average temperatures (in °C). These variables collectively serve as the bedrock for constructing predictive models, offering insights into future crop yields. The technical essence of this study lies in the adept application of diverse machine learning algorithms such as Linear Regression, Decision Trees, and Random Forests. These algorithms are trained on historical data to discern intricate patterns and relationships that contribute to the predictive accuracy of the models.

Beyond the technical complexities, the background of this research underscores the urgent need for precise crop yield predictions amid evolving climatic patterns and environmental dynamics. With burgeoning global populations and climate change introducing heightened uncertainty into agricultural systems, the imperative to optimize resource allocation, mitigate risks, and promote sustainable farming practices becomes increasingly pronounced. This study aspires to contribute meaningfully to this imperative by providing farmers and stakeholders with predictive tools capable of informing strategic decisions

1.2 Research problem statement

The research problem addressed in this study centers on the accurate prediction of crop yields in agriculture, employing a diverse set of machine learning algorithms. The challenge lies in developing models that can effectively leverage historical data, encompassing factors such as average rainfall, pesticide usage, and temperature, to forecast future crop yields reliably. Agricultural stakeholders face the critical need for advanced predictive tools that navigate the intricate relationships among these variables, offering insights into crop production's dynamic and often unpredictable nature.

Existing methodologies for crop yield prediction may need more precision to adapt to the evolving conditions of climate change and shifting environmental dynamics. Consequently, the research problem is defined by the imperative to overcome these limitations and provide more robust, accurate, and adaptable models. The study addresses the complexities inherent in predicting crop yields, acknowledging the multifaceted influences of environmental factors, agricultural practices, and technological interventions.

The research problem statement also encompasses the challenge of selecting and optimizing machine learning algorithms suitable for the specific characteristics of the dataset. Identifying the most effective algorithm or combination of algorithms is crucial for achieving reliable predictions. Furthermore, the study acknowledges the need for models that can account for temporal variations and trends, considering the time-dependent nature of agricultural processes.

1.3 Research objectives

- Evaluate and compare performance of diverse machine learning algorithms
- Conduct thorough data preprocessing, addressing missing values, outliers, and ensuring data consistency.
- Explore feature engineering techniques to enhance relevance and informativeness of input variables.
- Train machine learning models on historical data, optimizing hyperparameters for improved accuracy.
- Investigate ensemble methods or advanced algorithmic configurations to enhance model robustness.
- Incorporate temporal considerations into models to capture time-dependent trends and variations in crop yield patterns.
- Use robust performance metrics (e.g., MAE, MSE, R-squared) to comprehensively assess model accuracy.
- Develop visualizations to communicate model predictions and showcase significant features influencing crop yield.
- Enhance model interpretability to facilitate understanding among stakeholders.
- Implement cross-validation techniques to validate model performance and ensure generalization.
- Assess models' practical feasibility and applicability in real-world agricultural settings.
- Consider factors such as data availability, computational resources, and ease of implementation.
- Evaluate potential impact of accurate crop yield predictions on sustainable agricultural practices, resource optimization, and resilience to environmental changes.

1.4 Research Methodologies

- **Which machine learning algorithms demonstrate superior performance in predicting crop yields, and under what conditions?**
- *Explanation:* This question aims to identify the most effective algorithm or combination of algorithms for accurate crop yield prediction. Understanding the strengths and limitations of different models will contribute to the selection of an optimal approach based on the dataset's characteristics and the complexity of the problem.
- **How do various preprocessing techniques impact the accuracy of crop yield prediction models?**
- *Explanation:* This question delves into the influence of data preprocessing on model performance. By addressing issues such as missing values and outliers and exploring feature engineering, the study aims to determine the extent to which data quality and preprocessing contribute to the accuracy of crop yield predictions.
- **What is the temporal pattern in crop yield fluctuations, and how can it be effectively integrated into predictive models?**
- *Explanation:* Recognizing the temporal dynamics of crop yields is crucial for accurate predictions. This question seeks to understand the time-dependent trends and variations in crop yield patterns and how they can be incorporated into machine learning models to enhance their predictive capabilities.
- **To what extent do ensemble methods or advanced algorithmic configurations improve the robustness and generalization of crop yield prediction models?**

- *Explanation:* This question explores the potential benefits of employing ensemble methods or advanced algorithmic configurations. Understanding how these techniques contribute to model robustness and generalization will guide the development of more reliable and adaptable crop yield prediction models.
- **How do the developed crop yield prediction models contribute to sustainable agricultural practices, and what practical implications do they have for farmers and decision-makers?**
- *Explanation:* This question addresses the practical applicability and impact of the research. By assessing the real-world implications of the developed models, the study aims to provide insights into how accurate crop yield predictions can inform sustainable agricultural practices, optimize resource allocation, and support decision-making among farmers and stakeholders.

1.5 Scope and Limitations

Scope:

- **Geographical Scope:** Limited to the agricultural context of Albania.
- **Crop Types:** Encompasses maize, potatoes, rice, sorghum, soybeans, and wheat.
- **Time Frame:** Focuses on historical data with an emphasis on future crop yield predictions.
- **Algorithmic Focus:** Primarily explores machine learning algorithms like Linear Regression, Decision Trees, and Random Forests.

Limitations:

- **Data Limitations:** The accuracy depends on the quality and completeness of available data.
- **External Factors:** External variables not included in the dataset may impact model accuracy.
- **Generalizability:** Findings may not be universally applicable beyond the specified geographical and crop scope.
- **Algorithmic Constraints:** Limited to the chosen machine learning algorithms, excluding potential benefits from other advanced methods.
- **Temporal Constraints:** Predictions are based on historical trends, and unforeseen changes may not be adequately captured.

1.6 Document Outline

Chapter	Content
1. Introduction	- 1.1 Background: Overview of crop yield prediction significance - 1.2 Research Problem: Definition of challenges - 1.3 Research Objectives: Enumerated study goals - 1.4 Research Questions: Key inquiries guiding the study - 1.5 Scope & Limitations: Clearly defined study boundaries

2. Literature	- 2.1 Overview: Introduction to existing research in the field. - 2.2 Key Concepts: Exploration of relevant concepts and theories - 2.3 Methodologies: Review of methodologies used in similar studies| - 2.4 Gaps Identified: Identification of gaps in existing literature - 2.5 Relevance: How existing literature aligns with the current study|
3. Methodology	- 3.1 Data Collection: Detailed explanation of data sources - 3.2 Data Preprocessing: Techniques for cleaning and preparing data - 3.3 Model Selection: Choice and rationale behind machine learning algorithms - 3.4 Feature Engineering: Methods for enhancing input variables - 3.5 Temporal Analysis: Incorporation of temporal patterns - 3.6 Performance Metrics: Criteria for evaluating model performance - 3.7 Validation Techniques: Methods for ensuring model reliability
4. Results & Performance	Presentation of how different algorithms perform in crop yield prediction
5. Conclusion	Summary of Findings: Recapitulation of key research outcomes

CHAPTER 2 LITERATURE REVIEW AND RELATED WORK

[1] The study by Whitmire et al. employs various ML algorithms, including linear regression, regression trees, support vector machines, neural networks, Bayesian regression, and k-nearest neighbours, to predict alfalfa yield. The research compares the performance of these algorithms with different feature selection methods, such as correlation-based, Relief, and wrapper methods. The results reveal intriguing insights into the effectiveness of these techniques. Linear regression, often used as a baseline, shows its utility in predicting alfalfa yield. Neural networks, capable of handling complex, nonlinear functions, offer flexibility in capturing intricate relationships in the data. Support vector machines, employing a 'kernel trick' for nonlinear data, demonstrate their suitability for this agricultural prediction task-nearest neighbours, a spatially based method, showcase its ability to remember past data and make predictions based on proximity. Regression trees, forming decision trees to learn patterns, visually represent the key features influencing alfalfa yield Bayesian ridge regression, a probabilistic method, leverages probability distributions based on training data for predictions.

The study's emphasis on feature selection methods is noteworthy. The correlation-based method, outperforming others, identifies critical features like Julian's day of harvest, days between sown and harvest, cumulative solar radiation, and cumulative rainfall since the previous harvest. These features prove instrumental in achieving high prediction accuracy, particularly with the k-nearest neighbour and random forest methods.

The overall findings highlight the promising role of ML in predicting crop yields, even with relatively simple datasets. The research contributes valuable insights for agricultural planning, crop variety testing, and achieving global sustainability goals. It demonstrates that ML, coupled with effective feature selection, can play a pivotal role in addressing challenges related to poverty, health, and environmental conditions outlined by the United Nations.

[2] Their study by Juan Cao et al. aimed to address the pressing need for accurate and timely crop yield estimates, given the challenges posed by climate change, a growing global population, and increasing food demand. Focusing on China, the world's largest producer and consumer of wheat, particularly in the major winter wheat production regions, the researchers compared the performance of a traditional machine learning method, RF, with three deep learning models: DNN, 1D-CNN), and LSTM. They were integrating diverse datasets, including climate, satellite, and soil properties, and within the Google Earth Engine (GEE) platform allowed for a comprehensive analysis.

The study covered thirteen provinces, one municipality (Beijing), and one autonomous region (Ningxia) in China, comprising approximately 4.15 million square kilometres. The results showcased the efficacy of all four models in capturing winter wheat yield variations at the county level, with impressive performance metrics—mean R2 values exceeding 0.85 and RMSE below 768 kg/ha for the years 2011 to 2015. Notably, the DNN and RF models exhibited superior performance at the field level, emphasizing their ability to discern spatial heterogeneity and yield differences between individual fields across a county. With mean R2 values of 0.71 and 0.66 and RMSE values of 1127 kg/ha and 956 kg/ha, respectively, these models demonstrated a nuanced understanding of the complex factors influencing crop yields.

The study's findings extended beyond the immediate implications for crop management, emphasizing the proposed framework's scalability, simplicity, and cost-effectiveness. The ability to estimate crop yields reliably at various scales, including county and field levels, holds significance for agricultural disaster monitoring, formulation of food trade policies, and ensuring food security. One noteworthy aspect of the research was comparing traditional machine learning (RF) and deep learning models, challenging the prevailing assumption that deep learning consistently outperforms conventional methods. The conclusion highlighted the importance of a pragmatic and versatile approach, advocating for Random Forest as a competitive alternative in crop yield prediction.

This research contributes crucial insights into a scalable and efficient framework for timely and accurate crop yield estimation, addressing critical global challenges in agriculture and food security. The comprehensive analysis of machine learning and deep learning models provides valuable guidance for researchers, policymakers, and practitioners involved in sustainable and resilient agricultural practices.

[3] The literature review on crop yield prediction, with a particular emphasis on palm oil, offers a comprehensive exploration of recent studies in precision agriculture. Authored by Mamunur Rashid et al., the review underscores the importance of early and reliable crop yield estimation for strategic planning in the agricultural sector. The authors highlight the challenges and complexities associated with palm oil production, pivotal in global vegetable oil consumption.

The fundamental phases of the crop yield prediction process, encompassing data collection, preprocessing, partition, and analysis, are meticulously elucidated. Noteworthy features influencing crop yield, ranging from soil properties and climatic information to socio-economic factors, are identified as critical components in the predictive modelling framework. The study critically reviews machine learning algorithms employed in crop yield prediction, offering insights into RF, SVM, KNN, and ANN. A comprehensive summary of these algorithms and their benefits and drawbacks provides a valuable resource for researchers and practitioners.

Integrating wireless sensor WSNs and the IoT emerges as a transformative element in precision agriculture. The potential of WSNs to monitor climate and nutrient data, forecast crop health, and facilitate irrigation planning is underscored. Moreover, the revolutionary impact of IoT on smart agriculture is highlighted, offering solutions to challenges such as soil sampling, fertilization, disease monitoring, and yield measurement.

Concerning corn yield prediction, the review references studies employing machine learning techniques, including spectral vegetation indices, distance-based VI techniques, predictive models, and ensemble methods. The study by Shah Hosseini et al. introduces semiparametric neural networks (SNN) as a promising approach, showcasing superior out-of-sample prediction performance.

In conclusion, this literature review serves as a valuable synthesis of recent advancements in crop yield prediction, providing a nuanced understanding of the challenges and opportunities in the agricultural landscape. Integrating machine learning algorithms, wireless sensor networks, and IoT technologies emerges as a promising avenue for enhancing precision agriculture and addressing the complexities of global food production and security.

[4] The paper by Christopher D. Whitmire et al. delves into precision agriculture by employing ML and feature selection techniques to predict alfalfa yield—a pivotal factor in livestock feed production. The study, published in AI, draws on extensive yield data from diverse alfalfa varieties across several years in

Kentucky and Georgia. The authors systematically compare various ML models, including linear regression, regression trees, support vector machines, neural networks, Bayesian regression, and nearest neighbours. The impact of different feature selection methods is thoroughly examined, encompassing correlation-based, ReliefF, and wrapper methods.

The standout feature selection method is the correlation-based approach, revealing a feature set comprising the Julian day of harvest, days between sowing and harvest, cumulative solar radiation, and cumulative rainfall since the previous harvest. Integrating these features into the k-nearest neighbour and random forest methods yields remarkable results, with an average R-value surpassing 0.95 and an average mean absolute error below 200 lbs./acre. Notably, the achieved top R2 of 0.90 outperforms previous benchmarks, showcasing the potential of ML, coupled with feature selection, in predicting crop yields even with modest datasets and a limited feature set.

The significance of reporting accuracies in R and R2 for intuitive cross-crop comparisons is underscored throughout the paper. Within the Anaconda environment, Python serves as the primary programming language for data cleaning, visualization, and ML model development. Various packages, such as Pandas, matplotlib, seaborn, sci-kit learn, and Numpy, are leveraged for these tasks. Beyond conventional yield prediction, the researchers construct a decision tree classifying data into three distinct yield classes, offering valuable insights into the features influencing yield classification.

In conclusion, this work contributes substantially to the evolving landscape of precision agriculture, demonstrating the potential of ML and feature selection in advancing crop yield prediction. The findings provide a robust foundation for alfalfa yield forecasting and underscore the adaptability of these techniques for diverse crops, offering a valuable tool for informed decision-making in the agricultural sector. The authors' commitment to transparency and reproducibility is evident through the provision of open-access code on Github, enhancing the accessibility and applicability of their research.

[5] ML techniques have been emerged as powerful tools in the realm of precision agriculture, particularly in the prediction of crop yield and estimation of nitrogen status. In a comprehensive review by Anna Chlingaryana et al. from the Australian Centre for Field Robotics and the Centre for Carbon, Water and Food at the University of Sydney, the integration of ML with sensing technologies is explored to enhance decision support tools for modern farming systems.

Accurate crop yield estimation and optimized nitrogen management are crucial for agricultural success. While widely used, remote sensing (RS) systems require processing large volumes of data, prompting a shift towards ML methods due to their ability to handle vast datasets and nonlinear tasks. The paper discusses developments in ML techniques over the last 15 years for precise crop yield prediction and nitrogen status estimation.

ML techniques, such as Gaussian Processes, Dirichlet Processes, and Indian Buffet Process, offer probabilistic approaches that consider sensor noise, enabling the fusion of information from different sensors. The paper emphasizes the autonomy of ML techniques in solving significant nonlinear problems, providing a flexible framework for data-driven decision-making and incorporating expert knowledge into the system.

In precision agriculture, the primary goal is to provide information for improved management decisions across space and time. ML techniques contribute to the identification of management zones within fields, aiding in the application of specific treatments. The review explores how ML has been applied to detect biotic stress in crops, including identifying weeds, plant diseases, and insect pests.

Moreover, ML techniques applied to hyperspectral imaging data reveal physiological and structural characteristics in plants, enabling the tracking of environmental effects on crops. The authors discuss the advantages of ML techniques, such as Neural Networks, in automating the extraction of relevant features from data-enhancing crop yield prediction.

The paper also delves into studies using ML for precision nitrogen management. ML techniques help identify management zones for optimal nitrogen application, considering soil properties' complex correlations and spatial variability. The authors highlight studies that employ different ML methods for estimating nitrogen status and making informed decisions about fertilizer application.

Despite the successes of ML techniques, the review acknowledges their limitations, emphasizing the importance of data quality, model representativeness, and careful consideration of dependencies between input and target variables. The authors conclude that ongoing advancements in sensing technologies and ML techniques will provide cost-effective solutions for precise crop and environment state estimation, paving the way for transformative changes in crop yield management.

[6] The research paper titled "Agricultural Crop Yield Prediction Using Artificial Neural Network Approach" addresses the critical objective in agrihikar and Dr. Sandeep V. Rode from Sipna College of Engineering, Maharashtra, India, the study delves into the intricate connections between large-scale meteorological phenomena and crop yield. Recognizing the profound impact of climatological conditions on agriculture, the authors argue that early detection and management of issues affecting crop yield indicators are crucial for optimizing production and subsequent profits.

The research situates itself within the broader context of predicting crop yield for strategic plants like wheat, corn, and rice, considering the economic implications at both national and international levels. The authors contend that dry farming crop production is intricately linked to climatic events, necessitating accurate prediction models. Traditionally, two main categories of models—Statistical and Crop Simulation Models like CERES—have been employed. However, the paper asserts that recent Artificial Intelligence (AI) advancements, particularly Artificial Neural Networks (ANNs), have demonstrated increased efficiency in addressing complex issues within natural systems.

The study focuses on the application of feed-forward backpropagation ANNs, a widely used neural network architecture. ANNs are computational models , capable of machine learning and pattern recognition. As illustrated in Figure 1, the feed-forward backpropagation neural network processes information through interconnected "neurons" organized in layers. The "feed-forward" aspect describes the forward flow of information. At the same time, "backpropagation" refers to the supervised training method where anticipated outputs are compared against actual outputs.

The paper demonstrates the application of ANNs in predicting suitable crops based on various soil and atmospheric parameters. These parameters include soil type, pH, nitrogen, phosphate, potassium, organic carbon, temperature, rainfall, and humidity. The authors argue that ANNs offer a versatile approach to handling complex situations, mainly when certain factors are challenging or costly to measure. Using MATLAB for implementation, the research highlights the efficacy of the feed-forward backpropagation ANN in accurately predicting crop yields, particularly when considering regional soil characteristics.

In conclusion, the study affirms that ANNs, specifically the feed-forward backpropagation model, are beneficial for crop prediction. The integration of regional soil parameters enhances the accuracy of

predictions, showcasing the potential of AI in optimizing agricultural practices and contributing to the broader field of agro-meteorology.

[7] In this research paper, authors Khadijeh explore the application of DL models in predicting end-of-season yields for tomato and potato crops. The study recognizes the challenges in agriculture, such as climate change, environmental impacts, and resource limitations, emphasizing the need for innovative technologies to ensure higher productivity and sustainability.

The authors focus on two deep learning models, LSTM and GRU, and their extensions, Bidirectional Long Short-Term Memory (BLSTM) and Bidirectional Gated Recurrent Units (BGRU). These models analyze time-series data, including climate data. The research involves testing these models on datasets from a site in Portugal, specifically the Fadagosa region, where an agricultural weather station collected big climate data.

The results indicate that the BLSTM model outperformed the other models, including GRU, LSTM, and BGRU, on the validation dataset. The model demonstrated the nonlinear relationship between irrigation amount, climate data, and soil water content, achieving a mean squared error (MSE) of 0.017 to 0.039. The performance of the BLSTM model was also compared with other commonly used deep learning and CNN, MLP, and Random Forest Regression, with the BLSTM model showing superior accuracy.

The study underscores the potential of deep learning, particularly the BLSTM model, in predicting crop yields at the end of the season. The models leverage historical data to make accurate estimates, contributing to developing decision support systems for optimized irrigation scheduling and resource management in agriculture. The research emphasizes the importance of data-driven approaches and technological innovations, such as deep learning, in addressing the evolving challenges of modern agriculture.

[8] Crop yield prediction is a critical facet of contemporary agriculture, vital for addressing food security challenges and facilitating informed decision-making at regional and national levels. Despite the proliferation of models in the literature, the current performance of crop yield prediction models still needs to improve, necessitating continuous improvements. In response to this imperative, Oikonomidis et al. undertook a study leveraging deep learning-based models to assess their efficacy in predicting crop yield. The models evaluated encompassed a spectrum of algorithms, including the XGBoost machine learning algorithm, Convolutional Neural Networks (CNNs), Deep Neural Networks (DNNs), CNN-XGBoost, CNN-Recurrent Neural Networks (RNNs), and CNN-Long Short-Term Memory (LSTM). The study employed a comprehensive dataset derived from soybean crops in nine states in the United States, encompassing diverse features such as weather, soil parameters, and management data, totalling 395 features and 25,345 samples spanning the period from 1980 to 2018. Before model training, meticulous data processing techniques were applied, including handling missing values, removing outliers, and standardization through scaling. Feature selection was conducted using the SelectFromModel method based on importance weights. The results of the experiments demonstrated that the hybrid CNN-DNN model outperformed other models, achieving a Root Mean Squared Error (RMSE) of 0.266, Mean Squared Error (MSE) of 0.071, and Mean Absolute Error (MAE) of 0.199, with predictions aligning closely with an R2 value of 0.87. Notably, the CNN-DNN model, integrating convolutional and dense layers, exhibited superior predictive performance, showcasing the efficacy of deep learning techniques in extracting and leveraging essential features for crop yield prediction. The study emphasized the significance of computation speed and efficiency, especially in the context of time-series data, where hybrid models of CNN layers combined with either RNN or LSTM layers proved instrumental. These findings contribute valuable insights for stakeholders in agriculture, policy-making, and seed companies, offering a nuanced understanding of model performance and feature importance in crop yield prediction.

[9] In the pursuit of enhancing agricultural breeding and bolstering resilience against climatic challenges, Shook et al. conducted a pioneering study on crop yield prediction by integrating genotype and weather variables using deep learning methodologies. The research, emanating from the Department of Agronomy and Mechanical Engineering at Iowa State University, delves into the intricate dynamics of predicting crop performance in varying environments. The authors harnessed data from the Uniform Soybean Tests (UST) in North America, specifically focusing on soybean crops, a staple in the region's agricultural landscape. The study employed a Long Short-Term Memory (LSTM)—Recurrent Neural Network (RNN) based model, incorporating pedigree-relatedness measures and weekly weather parameters to dissect and predict genotype responses across diverse climatic conditions.

The research showcased their proposed models' superiority over existing machine learning counterparts, including Support Vector Regression with Radial Basis Function kernel (SVR-RBF), LASSO regression, and the data-driven USDA model for yield prediction. Notably, the models outperformed by leveraging a temporal attention mechanism for LSTM models, providing interpretability of crucial time windows during the growing season. The interpretability aspect is crucial for informing plant breeders offering valuable insights into the temporal nuances of genotype responses. The authors aptly addressed predicting crop performance in unseen environments, a pivotal concern in plant breeding and crop production.

The dataset, spanning over a decade and encompassing 5839 unique genotypes, was meticulously prepared, considering performance records, weather variables, and genotype clustering. Incorporating a wholly connected pedigree and K-means algorithm, the clustering approach enhanced model performance by accounting for relatedness among genotypes. The study emphasized the necessity of considering genetic and environmental variables in crop yield prediction, shedding light on the pivotal nexus between the two in plant breeding strategies.

The research advocated adopting deep learning models, specifically Long Short-Term Memory networks, citing their efficacy in capturing long-term temporal dependencies in complex multivariate sequences. The proposed models demonstrated efficiency in learning non-linear dependencies between various inputs, including weather variables, maturity groups, and genotype information, thereby presenting a robust solution to the intricate task of yield prediction. The findings of this study contribute significantly to the intersection of agriculture, machine learning, and climate science, offering a comprehensive and advanced approach to crop yield prediction that holds implications for future research, agricultural practices, and policy-making.

[10] Guoyong Leng and Jim W Hall's (2020) paper contributes to the literature on crop yield prediction by conducting an inter-comparison of machine learning (ML), regression, and process-based models. Focusing on maize yields in the United States, the study utilizes data from the US Department of Agriculture (USDA) and simulations from the Agricultural Modelling Inter-comparison and Improvement Project (AgMIP) and the Inter-Sectoral Impact Model Inter-comparison Project (ISI-MIP). Notably, the paper addresses the limitations of process-based models, emphasizing biases in yield simulations related to agricultural management representations.

In addition to process-based models, the study employs ML, specifically the Random Forest algorithm, and a traditional multiple linear regression model. The ML models, trained on historical data, exhibit advantages over traditional statistical approaches, offering the capability to handle complex interactions and nonlinear relationships among variables without assuming specific response function shapes. The regression model,

on the other hand, incorporates temperature, precipitation, and the year term to predict maize yield variations.

The comparison of these models reveals nuanced insights. While process-based models struggle with biases, ML and regression models show better skills in reproducing observed spatial distribution patterns of maize yields across the US. However, caution is urged in the interpretation, given the calibration differences between observation-driven ML and regression models and less-calibrated process-based models.

The study extends its analysis to assess the contribution of climate variations to maize yield variability. Climate variables and detrended yields are used to construct reduced models. Results indicate that ML estimates of climate variations explain 93% of historical yield variability, contrasting with process-based models' smaller estimates.

Furthermore, the paper extends its findings to project future maize yields under 1.5°C and two °C warming scenarios. Using the ML model, the study predicts a decrease in maize yield, particularly pronounced under the two °C warming scenario, aligning with previous estimations. The spatial distribution of yield loss indicates hot-spot regions in the central and southeastern US, emphasizing the importance of targeted adaptation measures.

In conclusion, this paper comprehensively compares modelling approaches to understanding crop yield variability and climate impacts.

[11] This comprehensive study endeavours to enhance the accuracy of tea crop yield prediction, recognizing its pivotal role in global consumption. The research was conducted in the Pakhal Valley from 2016 to 2019, utilizing data from the National Tea and High-Value Crop Research Institute. The novel approach combines the AquaCrop simulation model and machine learning algorithms for regression-based yield predictions. AquaCrop, calibrated through the Food and Agriculture Organization (FAO), harnesses weather, soil, crop, and agro-management data to estimate crucial parameters such as reference evapotranspiration, crop transpiration, biomass, and eventual yield. Simultaneously, an ensemble of linear regression-based techniques, including Linear SVR, AdaBoost Regressor, ARD Regression, Decision Tree Regressor, MLP Regressor, MLR, RANSAC Regressor, SLR, XGBoost, and SVM Regressor, are employed to predict yield, presenting a hybridized strategy that integrates both simulation and machine learning methodologies. *Rigorous data pre-processing* is a fundamental step involving cleaning and conversion to the plot level to ensure the robustness of subsequent analyses. Performance evaluation is anchored in well-established metrics— MAE, MSE, and RMSE —enabling a rigorous comparative analysis of AquaCrop and machine learning models. The study's noteworthy findings emphasize the superior predictive capabilities of machine learning algorithms, achieving comparable accuracy with reduced data requirements when juxtaposed with the AquaCrop model. However, the narrative is somewhat constrained by the absence of a detailed exposition of potential limitations or challenges faced during the study. Despite this, the research significantly contributes to the evolving landscape of crop yield prediction, shedding light on the potential benefits of amalgamating diverse methodologies. Future research trajectories could explore expanding the geographical scope, incorporating additional relevant variables, or exploring more advanced machine learning models, offering the prospect of revolutionizing agricultural forecasting practices on a broader scale.

[12] The research from the University of Waterloo, Canada, delves into the intricate task of predicting strawberry yield and farm prices, presenting a significant departure from conventional approaches. Addressing the limitations of existing models, which often rely on either traditional machine learning or simplistic deep learning architectures, the study introduces more sophisticated and comprehensive deep learning models. Notably, the Convolutional Long Short-Term Memory Recurrent Neural Network (CNN-LSTM) stands out significantly when augmented with attention mechanisms, showcasing a groundbreaking application in predicting strawberry yield and prices directly from weather input parameters. By leveraging a diverse dataset encompassing weather, soil, and yield data from 2006 to 2019, the research sheds light on the broader set of influencing factors crucial for accurate predictions—performance evaluations, employing metrics position the Attention-CNN-LSTM model as the most effective among various models tested. The emphasis on incorporating attention mechanisms into compound models demonstrates innovation and significantly improves prediction accuracy. The research's focus on the California strawberry dataset, a significant producer, provides valuable insights into the complexities of predicting agricultural outcomes, especially in the face of diverse environmental factors. The study underscores the potential of advanced artificial intelligence techniques and intense learning in revolutionizing fresh produce price predictions. By introducing attention mechanisms, the research contributes not only to understanding strawberry yield and pricing dynamics but also to the broader field of supply chain management in agriculture. In summary, this research represents a substantial advancement in predicting strawberry yield and farm prices, offering a novel perspective on integrating deep learning and attention mechanisms for enhanced accuracy in agricultural predictions, with potential implications for decision-making in the agricultural supply chain.

[13] Zhao Zhang's article presents a comprehensive study to predict maize yield in China. The research is crucial considering the global demand for maize and China's significant contribution to its production amid decreasing arable land. The authors emphasize the importance of incorporating various satellite data, including optical, fluorescence, and thermal, along with environmental factors, to enhance the accuracy of maize yield predictions.

The study focuses on four agroecological zones in China, employing LASSO, RF and XGBoost, and deep learning networks (LSTM). The results reveal that integrating multi-source data, including satellite observations and environmental variables, explains over 75% of the yield variation satellite data during the silking stage, and solar-induced chlorophyll fluorescence (SIF) demonstrates significant contributions to yield predictions. The study also identifies high temperature and vapour pressure deficit during the reproductive period as critical climate variables affecting maize production in China.

A key finding is the superiority of machine learning (ML) and deep learning (DL) approaches over regression-based methods. ML exhibits enhanced computational efficiency and generalization capabilities compared to DL.

The research contributes to the evolving field of crop yield prediction by offering insights into the optimal combinations of input variables. By systematically analyzing the influence of different variables at various stages of the growing season, the authors provide a nuanced understanding of the factors impacting maize yield. The proposed methodology, utilizing satellite and environmental data fusion, sets a paradigm for large-scale crop yield prediction with implications for global food security.

In conclusion, the article by Zhang et al. serves as a valuable contribution to the field of agricultural science, demonstrating the efficacy of integrating diverse datasets and advanced machine learning techniques for accurate maize yield predictions in China. The findings hold significance for agricultural practices, policy-making, and research endeavours worldwide.

[14] The research article addresses a critical aspect of precision agriculture by exploring integrating cultivar information and uncrewed aerial vehicle (UAV) remote sensing data to enhance potato yield prediction. Authored by Dan Li, Yuxin Miao, Sanjay K. Gupta, Carl J. Rosen, Fei Yuan, Chongyang Wang, Li Wang, and Yanbo Huang, the study delves into the significance of accurate high-resolution yield maps for identifying spatial yield variability patterns and understanding key factors influencing yield variability in potato crops.

The investigation, conducted through small plot experiments in 2018 and 2019, involves different cultivars and nitrogen rates. UAV-based multispectral images collected throughout the growing season serve as the primary dataset. The authors employ machine learning models, specifically for SVR, to amalgamate various vegetation indices with cultivar information.

A noteworthy discovery is that spectral data obtained from the early growing season, specifically at the tuber initiation stage in late June, exhibits a superior correlation with potato marketable yield compared to data collected during the tuber maturation stage in August. The study concludes that integrating cultivar information into machine learning algorithms significantly improves yield prediction, as evidenced by enhanced performance metrics compared to models solely utilizing remote sensing data.

The research underscores the global importance of potatoes as a high-yield food crop and the need for advanced methodologies in predicting and managing yield variability. Additionally, the article highlights the potential for further refinement of predictions by incorporating more detailed cultivar information, soil and landscape variables, and employing advanced machine learning models. Overall, the study contributes valuable insights to the ongoing discourse in precision agriculture, emphasizing the necessity for continued research to optimize potato yield predictions and support global food security.

[15] The authors compare the efficacy of a traditional machine learning method (Random Forest, RF) with three deep learning (DL) models—Deep Neural Networks (DNN), 1D Convolutional Neural Networks (1D-CNN), and LSTM. The goal is to predict winter wheat yields at county and field scales. The study covers the years 2011 to 2015, and the models are assessed based on recorded and simulated yields, focusing on the R^2 and RMSE.

Results demonstrate that all four models effectively capture winter wheat yield variations at the county level, exhibiting R^2 values ranging from 0.83 to 0.90 and RMSE from 561.18 to 959.62 kg/ha. Importantly, the DL models, particularly DNN and RF, excel at the field level, providing R^2 values of 0.71 and 0.66 and RMSE values of 1127 kg/ha and 956 kg/ha, respectively.

The study highlights the significance of scalable, simple, and cost-effective frameworks for estimating crop yields, emphasizing the application of publicly available datasets and the potential of DL and ML algorithms. The Google Earth Engine platform emerges as a valuable resource, leveraging its vast data repository and computational capabilities. The findings offer crucial insights for crop yield forecasting, agricultural disaster monitoring, food trade policy, and national food security initiatives

Title	**year**	**Research Question**	**Methods used**	**Results achieved**
Predicting Alfalfa Yield	2023	What is the effectiveness of various ML	Linear regression, regression trees, support vector	- Linear regression proves useful as a

		algorithms in predicting alfalfa yield,?	machines, neural networks, Bayesian regression	baseline for predicting alfalfa yield.
Enhancing Crop Yield Estimates in China's Wheat Production Regions: A Comparative Analysis of Traditional and Deep Learning Models	2022	How do traditional machine learning methods, specifically Random Forest (RF)	RF, DNN, 1D-CNN, LSTM;	- All models demonstrate efficacy in capturing winter wheat yield variations at the county level.
Advances in Crop Yield Prediction: A Literature Review with Emphasis on Palm Oil	2021	What are the recent advancements in precision agriculture, specifically in crop yield prediction.	(RF, SVM, KNN, ANN), WSNs, IoT	Comprehensive exploration of recent studies in precision agriculture, with a specific focus on palm oil production.
Precision Agriculture: ML and Feature Selection in Alfalfa Yield Prediction	2021	How do various machine learning models, perform in predicting alfalfa yield, ?	ML models: linear regression, regression trees, support vector machines, neural networks, Bayesian regression, nearest neighbours; Feature selection methods	-Correlation-based feature selection identifies critical features
ML in Precision Agriculture: Advances, Challenges, and Opportunities	2022	How can machine learning techniques enhance decision support tools for modern farming systems,	ML techniques: Gaussian Processes, Dirichlet Processes, Indian Buffet Process, Neural Networks;	ML techniques offer probabilistic approaches, considering sensor noise, enabling fusion of information from different sensors.
Agricultural Crop Yield Prediction Using Artificial Neural Network Approach	2023	How effective is the application of ANNs, specifically the feed-forward backpropagation model, in predicting crop	ANN architecture: feed-forward backpropagation.	ANNs, especially feed-forward backpropagation, efficiently predict suitable crops based on soil .

		yields based on various soil ?		
Application of DL Models in Predicting End-of-Season Yields for Tomato and Potato Crops	2021	How effective are deep learning models, specifically LSTM and GRU, and their extensions (BLSTM and BGRU), in predicting end-of-season yields for tomato and potato crops using time-series data, including climate data?	Deep learning models: LSTM, GRU, Bidirectional LSTM (BLSTM), Bidirectional Gated Recurrent Units (BGRU); Dataset: Fadagosa region, Portugal; Comparison with CNN, MLP, Random Forest Regression	BLSTM outperforms other models, showcasing a nonlinear relationship between irrigation amount, climate data, and soil water content. - Achieves a mean squared error (MSE) of 0.017 to 0.039
Deep Learning-Based Models for Improved Crop Yield Prediction	2022	How do various deep learning-based models, including XGBoost, CNNs, DNNs, CNN-XGBoost, CNN-RNNs, and CNN-LSTM, perform in predicting soybean crop yields ?	Deep learning models: XGBoost, CNNs, DNNs, CNN-XGBoost, CNN-RNNs.	Hybrid CNN-DNN model outperforms other models with RMSE of 0.266, MSE of 0.071, MAE of 0.199, and R2 of 0.87.
Integrating Genotype and Weather Variables for Crop Yield Prediction	2023	How effective is a Long Short-Term Memory (LSTM)—Recurrent Neural Network (RNN)?	Deep learning model: LSTM-RNN; Comparison with SVR-RBF, LASSO regression, and USDA model; Dataset:	LSTM-RNN model outperforms existing machine learning counterparts. time windows
Inter-comparison of ML, Regression, and Process-Based Models for Maize Yield Prediction	2020	How do machine learning (ML), regression, and process-based models compare in predicting maize yields ?	Models: Random Forest (ML), Multiple Linear Regression (Regression), AgMIP and ISI-MIP (Process-Based)	ML and regression models show better skills in reproducing observed spatial distribution patterns of maize yields compared to process-based models.
Hybrid Approach: AquaCrop	2021	How effective is the combination	Models: AquaCrop	Machine learning algorithms

Simulation Model and ML for Tea Crop Yield Prediction		of the AquaCrop simulation model and various machine learning algorithms?	simulation model, Linear SVR, AdaBoost Regressor, ARD Regression, Decision Tree Regressor, MLP Regressor, MLR, RANSAC Regressor, SLR, XGBoost, SVM Regressor;	achieve comparable accuracy with reduced data requirements compared to the AquaCrop model.
Advanced Deep Learning Models for Strawberry Yield and Price Prediction	2022	How effective are advanced deep learning models, particularly the Attention-CNN-LSTM, in predicting strawberry yield? ?		- Attention-CNN-LSTM model, with attention mechanisms, outperforms other models in predicting strawberry yield and farm prices.
Maize Yield Prediction in China: Integrating Satellite Data	2022	How effective is the integration of various satellite data ?	Models: LASSO, Random Forest (RF), XGBoost, Long Short-Term Memory (LSTM);	- Over 75% of yield variation explained by integrating multi-source data during the silking stage. - Solar-induced chlorophyll fluorescence (SIF) significantly contributes to yield predictions.
Potato Yield Prediction: Integrating Cultivar and UAV Data	2021	How does integrating cultivar information and uncrewed aerial vehicle (UAV) remote sensing data enhance potato yield prediction, and what is the significance of spectral data obtained during different stages of the growing season?	Model: Support Vector Regression (SVR);	Spectral data from the early growing season (late June) correlates better with potato marketable yield than data from the tuber maturation stage (August). - Integrating cultivar information improves yield prediction compared to models using only remote sensing data

Winter Wheat Yield Prediction: RF vs. DL Models	2021	How do traditional machine learning methods (Random Forest, RF) compare with three deep learning models (DNN, 1D-CNN, LSTM) in predicting winter wheat yields at county and field scales, and what are the implications for scalable and cost-effective frameworks?	Models: Random Forest (RF), Deep Neural Networks (DNN), 1D Convolutional Neural Networks (1D-CNN), Long Short-Term Memory (LSTM); Evaluation metrics: R^2, RMSE; Data: Years 2011-2015	-All four models effectively capture winter wheat yield variations at the county level (R^2: 0.83 to 0.90, RMSE: 561.18 to 959.62

Table 1 Literature study

CHAPTER 3 DESIGN AND METHODOLOGY

3.1 Crisp dm methodology




Figure 1 Crisp Dm process

CRISP-DM :

CRISP-DM is a widely used and well-established methodology for guiding data mining and machine learning projects. It provides a structured and systematic approach to data mining, from understanding business objectives to deploying the final model. The CRISP-DM methodology consists of the following six phases:

1. **Business Understanding:**
- Define the business problem and objectives.
- Establish project goals and criteria for success.
- Develop an initial plan to achieve the objectives.

2. **Data Understanding:**
- Collect and explore the available data.
- Describe the data, identify data quality issues, and assess its suitability for analysis.
- Verify data quality and address issues as needed.

3. **Data Preparation:**
- Select data relevant to the analysis.
- Clean and preprocess the data, addressing missing values and outliers.

- Transform variables and create new features as necessary.

4. **Modelling:**

- Select appropriate modelling techniques based on the business problem.
- Build and train machine learning models using the prepared data.
- Evaluate and refine models to achieve optimal performance.

5. **Evaluation:**

- Assess the model's performance against business objectives.
- Validate the model on independent datasets to ensure generalization.
- Review and interpret results to determine the model's effectiveness.

6. **Deployment:**

- Implement the model into the production environment.
- Develop a deployment plan and monitor the model's performance in real-world scenarios.
- Document and communicate the results and insights.

CRISP-DM is iterative, allowing for revisiting earlier phases based on the insights gained during subsequent stages. It emphasizes collaboration between business stakeholders and data professionals throughout the project lifecycle. It has become a widely accepted framework in data science and machine learning due to its practical and flexible nature.

3.1 Dataset Exploration:

The provided dataset offers a comprehensive glimpse into agricultural dynamics, focusing on crop yield prediction in Albania over several years. Encompassing diverse crops such as maize, potatoes, rice, sorghum, soybeans, and wheat, the dataset provides a multifaceted view of agricultural productivity. Each entry includes crucial parameters such as the type of crop, the corresponding year, yield per hectare , average annual rainfall, the number of pesticides used, and the average temperature .

With its varied climate and geographical features, Albania poses a unique challenge for crop yield prediction, making this dataset particularly valuable. Including different crops allows for a nuanced analysis of how varying environmental conditions impact distinct agricultural outputs. The temporal dimension is critical in understanding the dataset, as it spans multiple years, enabling the exploration of trends, patterns, and potential cyclical variations in crop yields.

The dataset also sheds light on the influence of external factors such as pesticide usage, rainfall, and temperature on crop productivity. The correlation between these variables and crop yield can be examined to uncover dependencies and refine predictive models.

However, as with any dataset, there are potential challenges. Missing data, outliers, or inconsistencies may need careful handling during preprocessing. Additionally, the dataset's temporal nature suggests the need for specialized time-series analysis techniques to capture the evolving patterns over the years.

In machine learning and agricultural science, this dataset serves as a valuable resource for developing and fine-tuning predictive models. The interplay of variables provides an opportunity to employ algorithms such as Linear Regression, Decision Trees, or Random Forests to discern complex relationships and enhance the accuracy of crop yield predictions. By delving into this dataset, researchers can contribute to advancements in sustainable farming practices, resource optimization, and the broader goal of ensuring food security in the face of changing environmental conditions.

3.2 Architecture




Figure 2 Architecture

3.3 Data Preprocessing

3.3.1 Top 10 Areas based on rainfall:

This graph gives the information on rainfall recored in different countries from our dataset, India recieves the highest, republic recieves the lowest.




Figure 3 top 10 areas based on rainfall

This graph gives the information on highest yieled crops in the history, potatoes seems to have more yield when compared with all the other crops, where as soyabeans is less in recent years.




Figure 4 Highest yield crops

Since hg/ha is our target column, this graph shows the overall percentahes by area and by the crops occupancy in catageory.




Figure 5 Top areas with good yield

This graph gives the information on overall rainfall received over the years given in the dataset.




Figure 6 rainfall

This graph gives the information on overall temperature distributions




Figure 7 Average distributed temperature

The below graph shows the coorelation of overall features present in the dataset, some have positive correlation, some have negative correlation.




Figure 8 Corrleation matrix

3.4 Ethics:

In the realm of data-driven agriculture, ethical considerations are paramount. Using machine learning to predict crop yields demands transparency, fairness, and privacy. Ethical practices involve informed consent for data collection, ensuring unbiased representation of diverse agricultural practices, and safeguarding against unintended consequences. Researchers must prioritize responsible data handling, protecting farmers' privacy, and avoiding discriminatory outcomes. Upholding ethical standards is integral to fostering trust among stakeholders and ensuring the equitable and responsible application of machine learning in agriculture.

3.5 Machine Learning Algorithms

3.5.1 Random Forest :

Random Forest, an ensemble learning algorithm, emerges as a robust contender in the landscape of crop yield prediction. Its strength lies in constructing many decision trees during training and amalgamating their outputs to yield a final prediction. The algorithm's notable advantage is its capacity to achieve high accuracy, making it particularly valuable in scenarios where precision in crop yield forecasting is paramount. Its resilience against overfitting, a pervasive concern in predictive modelling, sets Random Forest apart. The infusion of randomness in feature selection and data sampling contributes to its ability to generalize well to unseen datasets.

One of the distinguishing features of Random Forest is its capability to provide insights into the importance of features. This attribute proves crucial in agricultural contexts, where understanding the variables significantly influencing crop yields is instrumental for informed decision-making. However, the algorithm has its challenges. The computational expense can be a limiting factor, especially when dealing with a substantial ensemble of decision trees. Additionally, interpreting the individual decision trees within the ensemble can be intricate, impacting the algorithm's overall interpretability.

In the specific realm of crop yield prediction, Random Forest finds applicability due to its adeptness at handling complex datasets with numerous variables and potential nonlinear relationships. Its resilience to overfitting is particularly advantageous in scenarios where the dataset may be subject to noise or exhibit intricate patterns. Nevertheless, implementers must exercise caution regarding computational resources, significantly as the ensemble size increases, and carefully balance the trade-off between accuracy and model interpretability. In conclusion, Random Forest is a potent tool in the agricultural data scientist's arsenal, offering a versatile solution for accurate and reliable crop yield predictions with due consideration to its computational complexity and interpretability challenges.

3.5.2 Decision Tree

In the realm of crop yield prediction, Decision Trees emerge as versatile and interpretable models. Operating on a hierarchical structure of decision nodes, they navigate through features to make predictions or classifications. One of their distinct advantages lies in interpretability, offering a transparent representation of decision-making processes. This feature is precious in agricultural contexts where stakeholders may need more deep machine learning expertise. Decision Trees excel in capturing nonlinear relationships within datasets, providing a robust approach for scenarios characterized by intricate variable interactions. Moreover, the algorithm inherently reveals the importance of different features, enabling a nuanced understanding of the factors influencing crop yield predictions.

Despite these strengths, Decision Trees are not immune to challenges. Their susceptibility to overfitting, especially in cases where the tree grows excessively deep, poses a concern. Deep trees may capture noise rather than meaningful patterns in the data, emphasizing the need for careful pruning and parameter tuning to balance complexity and generalization. Regularization techniques, such as limiting tree depth, can mitigate overfitting.

Decision trees are valuable tools in data scientists' arsenals in crop yield prediction. Their transparency facilitates communication with stakeholders, and their ability to discern complex relationships is advantageous in the multifaceted domain of agriculture. Furthermore, Decision Trees provide insights into the relative importance of different variables, aiding in identifying critical drivers for crop productivity.

While mindful consideration of overfitting is essential, Decision Trees contribute significantly to the interpretability and accuracy of crop yield prediction models, making them a compelling choice for agricultural data analysis and decision support.

3.5.3 Support Vector Machine

In machine learning, the Support Vector Machine (SVM) emerges as a robust algorithm, and its applications extend seamlessly to the domain of crop yield prediction. SVM is designed to excel in scenarios where data points need to be classified or predicted, and its effectiveness lies in identifying the optimal hyperplane in a high-dimensional space. This hyperplane acts as a boundary that maximally separates data points belonging to different classes. When applied to crop yield prediction, SVM showcases notable strengths. Its versatility becomes apparent in handling datasets with numerous dimensions, an attribute particularly valuable in agriculture where diverse factors influence crop productivity. SVM is adept at capturing intricate relationships between variables, making it suitable for agricultural datasets' complex and multifaceted nature.

One of SVM's distinguishing features is its robustness against overfitting. This ensures the model's predictive capabilities extend beyond the training dataset, offering reliable predictions for new and unseen data. In the agricultural context, where predicting crop yields accurately is essential for informed decision-making, SVM's ability to generalize well is paramount. Furthermore, SVM's effectiveness in delineating complex decision boundaries contributes to its utility in scenarios where the relationships between variables are not linear.

As agriculture embraces data-driven approaches for optimization and sustainability, SVM emerges as a valuable tool for crop yield prediction. Its application goes beyond mere classification; it aids in understanding the intricate interplay of variables influencing crop outcomes. The algorithm's capacity to navigate the complexities of agricultural datasets positions SVM as a critical contributor to advancements in precision farming and the overarching goal of optimizing agricultural practices for increased efficiency and sustainability.

3.5.4 Gradient Boosting algorithm

Gradient Boosting, an ensemble learning algorithm, is a formidable force in predictive modelling, finding applications in diverse domains, including crop yield prediction. This iterative method amalgamates weak learners, often decision trees, refining their predictions with each step. Renowned for its high predictive accuracy, Gradient Boosting excels in capturing intricate nonlinear relationships inherent in agricultural datasets. It sequentially fits models to correct the errors of the combined ensemble, progressively enhancing the model's predictive capabilities.

A distinguishing feature of Gradient Boosting is its ability to unveil the importance of features, providing valuable insights into the variables influencing crop yields. This interpretability is crucial for stakeholders seeking to understand the driving factors behind predictions. Additionally, Gradient Boosting exhibits resilience against overfitting when appropriately tuned, contributing to the model's reliability.

In the realm of crop yield prediction, Gradient Boosting serves as a sophisticated and powerful tool. Its iterative refinement process, emphasis on correcting errors, and capacity to handle complex relationships make it well-suited for agricultural data's dynamic and multifaceted nature. By harnessing the collective strength of weak learners, Gradient Boosting is a key player in improving the accuracy and interpretability of crop yield prediction models, offering valuable insights for informed agricultural decision-making.

3.5.5 CNN

The Convolutional Neural Network (CNN) stands as a formidable deep learning architecture, predominantly recognized for its image processing and pattern recognition prowess. Its application extends into diverse domains, including agriculture, where CNNs exhibit a spatial analysis capacity crucial for crop yield prediction tasks. CNNs excel in capturing intricate spatial hierarchies within images, making them particularly well-suited for tasks where the arrangement of features holds significance, such as identifying crop patterns within fields. One of the critical advantages of CNNs is their ability to autonomously learn hierarchical features from raw data, reducing the manual effort required for feature engineering. Their inherent translation invariance further enhances their utility, enabling the recognition of patterns irrespective of their position in the image.

Despite their strengths, CNNs come with computational demands, especially during training, requiring substantial computational resources. Additionally, the optimal performance of CNNs often hinges on large, diverse datasets, posing a challenge in agriculture, where comprehensive datasets may be limited. Another consideration is the interpretability of CNNs, as their complex architectures can make understanding the rationale behind their decisions challenging compared to simpler models.

In crop yield prediction, CNNs find application in analyzing aerial or satellite imagery, capturing crucial visual cues such as field boundaries, crop types, and growth stages. While challenges like computational intensity and data availability persist, the promise of CNNs in revolutionizing precision agriculture and enhancing crop monitoring remains substantial. As technology advances and datasets become more comprehensive, CNNs stand poised to play a pivotal role in reshaping how we approach and optimize crop yield prediction in the agricultural landscape.

3.5.6 LSTM

Long Short-Term Memory, a subtype of recurrent neural networks, has established itself as a potent solution for tasks involving sequential data, and its relevance extends notably to crop yield prediction. Uniquely designed to mitigate the vanishing gradient problem, LSTMs excel in capturing long-range dependencies within sequential data, a crucial attribute in the context of agricultural variables influenced by historical trends. Their proficiency in sequential learning makes LSTMs well-suited for unravelling the temporal intricacies inherent in agricultural processes, such as the stages of crop growth and the patterns of weather conditions. A key strength lies in their capacity to retain information across extended sequences, enabling

the incorporation of lasting effects from past conditions on crop outcomes. The adaptability of LSTMs to varying input sequences and different time intervals enhances their suitability for the dynamic nature of agricultural datasets. However, the advantages of LSTMs come with computational challenges, as their training can be resource-intensive, and their intricate architecture introduces complexity in parameter tuning. Despite these hurdles, the potential of LSTMs to elevate the accuracy of crop yield predictions is substantial, especially with ongoing advancements in technology, increased accessibility to computational resources, and the expansion of comprehensive agricultural datasets. The unique ability of LSTMs to capture temporal dependencies positions them as key contributors to refining the precision and reliability of models in the ever-evolving landscape of crop yield prediction.

CHAPTER 4 EVALUATION OF METRICS AND VALUES

Crop yield prediction is a critical task in agriculture, aiming to provide accurate forecasts of agricultural outcomes. Various machine learning models, including Long Short-Term Memory (LSTM), Convolutional Neural Network (CNN), Random Forest, Gradient Boosting Machine (GBM), Support Vector Machine (SVM), and Decision Tree, have been employed for this purpose. Evaluating the performance of these models is essential to gauge their accuracy and reliability. One common approach to assess model performance is using evaluation metrics, with Root Mean Square Error (RMSE) being a prominent measure. Let us delve into the evaluation of each model based on their respective RMSE values.

1. LSTM:

The LSTM model, designed to capture temporal dependencies in sequential data, yielded an RMSE of 1.0598. This metric quantifies the average magnitude of the errors between predicted and observed crop yield values. The relatively low RMSE suggests that the LSTM model performed well in predicting crop yields, capturing the dataset's nuanced temporal patterns and dependencies. The model's proficiency in handling sequential data and adaptability to varying time intervals contribute to its effectiveness in crop yield prediction.

2. CNN:

The CNN, known for its spatial analysis capabilities in image data, demonstrated an RMSE of 1.1032. While slightly higher than the LSTM, this metric still indicates a reasonably accurate prediction performance. CNNs excel in capturing spatial hierarchies and patterns within images, making them suitable for tasks where the arrangement of features is crucial. In crop yield prediction, the CNN's ability to analyze aerial or satellite imagery contributes to its effectiveness. However, potential challenges related to computational intensity and data availability should be considered.

3. Random Forest:

The Random Forest model exhibited a notably low RMSE of 0.1550, signifying predictive solid accuracy. Random Forests are ensemble models that aggregate multiple decision trees, leveraging their collective strength to enhance predictive capabilities. The low RMSE suggests that the Random Forest effectively captured the complex relationships within the agricultural data, showcasing its versatility in handling diverse variables and potential nonlinear patterns.

4. GBM:

The Gradient Boosting Machine (GBM) produced an RMSE of 0.4399, indicating a robust performance in predicting crop yields. Like Random Forest, GBM is an ensemble learning method that sequentially builds decision trees. The lower RMSE suggests that GBM excelled in correcting errors from previous models, capturing intricate relationships and patterns within the agricultural data. Its ability to handle complex nonlinearities contributes to its effectiveness in crop yield prediction.

5. SVM:

Support Vector Machines (SVMs), recognized for their effectiveness in high-dimensional spaces, yielded an RMSE of 0.9474. SVMs are adept at handling intricate relationships within datasets, making them suitable for scenarios where numerous dimensions influence agricultural variables. While the RMSE is higher than Random Forest and GBM, it still suggests a reasonable level of accuracy in predicting crop yields. The SVM's effectiveness in capturing complex relationships and its resilience to overfitting contribute to its utility in agriculture.

6. Decision Tree:

The Decision Tree model, known for its interpretability, demonstrated a low RMSE of 0.2114. Decision Trees operate by recursively partitioning data based on features, making them suitable for understanding

the importance of different variables. The low RMSE suggests that Decision Trees effectively captured patterns within the agricultural data, providing valuable insights into the factors influencing crop yields.
In conclusion, evaluating crop yield prediction models through metrics like RMSE provides a comprehensive understanding of their accuracy and suitability for agricultural applications. The choice of a specific model depends on the specific characteristics of the dataset, the nature of the variables involved, and the computational resources available. The ensemble models, Random Forest and GBM, demonstrated predictive solid capabilities, while the LSTM and CNN showcased their effectiveness in handling sequential and spatial data, respectively. SVM and Decision Tree models also performed well, leveraging their unique strengths in high-dimensional spaces and interpretability. As agriculture increasingly adopts data-driven approaches, these evaluation metrics play a crucial role in guiding the selection and optimization of models to enhance the precision and reliability of crop yield predictions.




Figure 9 Comparision of results

CHAPTER 5 CONCLUSION AND FUTURE WORK

5.1 Research Overview

This research embarks on a crucial exploration of the intersection of agriculture and data science, focusing on crop yield prediction in Albania. With global agricultural sustainability and food security in mind, the study leverages diverse machine-learning algorithms to analyze a comprehensive dataset encompassing historical crop yields, rainfall, pesticide usage, and temperatures. The research addresses the challenge of accurately predicting crop yields amidst climate change and evolving environmental dynamics. Objectives include algorithm performance evaluation, robust data preprocessing, feature engineering, and the incorporation of temporal considerations for more precise predictions. The study aspires to contribute impactful tools for stakeholders, fostering sustainable farming practices, resource optimization, and resilience in the face of environmental uncertainties.

5.2 Problem Definition

The identified problem centres on accurately predicting crop yields amid the complexities of climate change and evolving environmental dynamics. Aligned with the global imperative for agricultural sustainability and food security, the research employs diverse machine-learning algorithms to analyze a rich dataset, including historical crop yields, rainfall, pesticide usage, and temperatures.

5.3 Design/Experimentation, Evaluation & Results

Model	RMSE
LSTM	1.0598
CNN	1.1032
Random Forest	0.1550
GBM	0.4399
SVM	0.9474
Decision Tree	0.2114

Figure 10 Evaluation of results

5.4 Contributions and impact

The section outlines how the project contributes to the enhancement of existing methodologies for crop yield prediction, showcasing its potential to revolutionize agricultural practices. Furthermore, it highlights the practical applications of the research findings, demonstrating their relevance to farmers, agricultural stakeholders, and policymakers. By elucidating the project's distinctive contributions and underscoring its broader impact, this section aims to emphasize the transformative potential of the research in optimizing crop production, fostering sustainable farming practices, and addressing challenges in food security.

5.5 Future Work & recommendations

Future work in crop yield prediction could focus on refining model architectures and exploring hybrid models that combine the strengths of different approaches. Additionally, incorporating more diverse and comprehensive datasets, including factors like socio-economic variables and advanced sensing technologies, could enhance model accuracy. Integration of real-time data and continuous model updates could further improve predictions. Exploring explainability techniques for complex models like LSTM and CNN would enhance interpretability. Collaboration between data scientists, agronomists, and stakeholders is crucial for tailoring models to specific agricultural contexts. Future research should aim to develop more robust, interpretable, and adaptive models that contribute to sustainable and data-driven agricultural practices.

References:

1. Whitmire, C. D., Vance, J. M., Rasheed, H. K., Missaoui, A., Rasheed, K. M., & Maier, F. W. (2021). Using Machine Learning and Feature Selection for Alfalfa Yield Prediction. *AI*, 2(1), 71-88. https://doi.org/10.3390/ai2010006.
2. Zhang, J., Tian, H., Wang, P., Tansey, K., Zhang, S., & Li, H. (2021). Improving wheat yield estimates using data augmentation models and remotely sensed biophysical indices within deep neural networks in the Guanzhong Plain, PR China. *Computers and Electronics in Agriculture*, 106616. https://doi.org/10.1016/j.compag.2021.106616.
3. Rashid, M., Bari, B. S., Yusup, Y., Kamaruddin, M. A., & others. (2021). A Comprehensive Review of Crop Yield Prediction Using Machine Learning Approaches With Special Emphasis on Palm Oil Yield Prediction. *IEEE Access*, 9. https://doi.org/10.1109/ACCESS.2021.3075159
4. Vance, J. M., Whitmire,.Comparing Machine Learning Techniques for Alfalfa Biomass Yield Prediction.
5. Sharma, A., Jain, A., Gupta, P., & Chowdary, V. (2020). "Machine Learning Applications for Precision Agriculture: A Comprehensive Review." *IEEE Access*. DOI: 10.1109/ACCESS.2020.3048415.
6. Dahikar, S. S., & Rode, S. V. (2014). "Agricultural Crop Yield Prediction Using Artificial Neural Network Approach." *International Journal of Innovative Research in Electrical, Electronics, Instrumentation and Control Engineering*, 2(1), 683. Retrieved from www.ijireeice.com
7. Alibabaei, K., Gaspar, P. D., & Lima, T. M. (2021). "Crop Yield Estimation Using Deep Learning Based on Climate Big Data and Irrigation Scheduling." *Energies*, 14(11), 3004. https://doi.org/10.3390/en14113004.
8. Oikonomidis, A., Catal, C., & Kassahun, A. (2022). "Hybrid Deep Learning-based Models for Crop Yield Prediction." Article ID: 2031822. Received: October 16, 2021. Accepted: January 18, 2022. Published online: January 22, 2022.
9. Leng, G., & Hall, J. W. (2020). "Predicting spatial and temporal variability in crop yields: an inter-comparison of machine learning, regression and process-based models." Environmental Research Letters, 15(4), 044027. Published 20 April 2020. https://doi.org/10.1088/1748-9326/ab7b24
10. Reddy, D. J., & Kumar, M. R. (2021). Crop Yield Prediction using Machine Learning Algorithm. In *5th International Conference on Intelligent Computing and Control Systems (ICICCS)*. IEEE.
11. D. Batool, M. Shahbaz, H.S. Asif, et al., "A Hybrid Approach to Tea Crop Yield Prediction Using Simulation Models and Machine Learning," *Plants*, vol. 11, no. 15, p. 1925, 2022. [DOI: 10.3390/plants11151925].
12. X. Chen, L. Feng, R. Yao, X. Wu, J. Sun, W. Gong, "Prediction of Maize Yield at the City Level in China Using Multi-Source Data," *Remote Sens.*, vol. 13, no. 1, p. 146, 2021. [DOI: 10.3390/rs13010146]
13. D. Li, Y. Miao, S.K. Gupta, C.J. Rosen, F. Yuan, C. Wang, L. Wang, Y. Huang, "Improving Potato Yield Prediction by Combining Cultivar Information and UAV Remote Sensing Data Using Machine Learning," *Remote Sens.*, vol. 13, no. 16, p. 3322, 2021. [DOI: 10.3390/rs13163322]
14. Champaneri, M., Chachpara, D., Chandvidkar, C., & Rathod, M. (2020). Crop Yield Prediction Using Machine Learning. *International Journal of Science and Research (IJSR)*, *9*(4). Retrieved from http://www.ijsr.net. (ISSN: 2319-7064)

15. Srivastava, N. Safaei, S. Khaki, G. Lopez, W. Zeng, F. Ewert, T. Gaiser, J. Rahimi, "Winter wheat yield prediction using convolutional neural networks from environmental and phenological data," *Sci. Rep.*, vol. 12, article number: 3215, 2022. [DOI: 10.1038/s41598-022-06124-z]