

# Orbital-use fees could more than quadruple the value of the space industry

Akhil Rao<sup>a</sup>, Matthew G. Burgess<sup>b,c,d</sup>, and Daniel Kaffine<sup>d</sup>

<sup>a</sup>Department of Economics, Middlebury College, Middlebury, VT 05753; <sup>b</sup>Cooperative Institute for Research in Environmental Sciences, University of Colorado Boulder, CO 80309; <sup>c</sup>Environmental Studies Program, University of Colorado Boulder, CO 80303; <sup>d</sup>Department of Economics, University of Colorado Boulder, 80302

This manuscript was compiled on December 3, 2019

**The space industry's rapid recent growth represents the latest Tragedy of the Commons. Satellites launched into orbit contribute to—and risk damage from—a growing buildup of space debris and other satellites. Collision risk from this orbital congestion is costly to satellite operators. Technological and managerial solutions—such as active debris removal or end-of-life satellite deorbit guidelines—are currently being explored by regulatory authorities. However, none of these approaches address the underlying incentive problem: satellite operators do not account for costs they impose on each other via collision risk. Here, we show an internationally harmonized orbital-use fee can correct these incentives and substantially increase the value of the space industry. We construct and analyze a coupled physical-economic model of commercial launches and debris accumulation in low-Earth orbit. Similar to carbon taxes, our model projects an optimal fee that rises at a rate of 14.8% per year, equal to roughly \$250,000 per satellite-year in 2040. The long-run value of the satellite industry would more than quadruple by 2040—increasing from around \$600 billion under business-as-usual to around \$3 trillion. In contrast, we project that purely technological solutions are unlikely to fully address the problem of orbital congestion. Indeed, we find debris removal sometimes worsens economic damages from congestion by increasing launch incentives. In other sectors, addressing the Tragedy of the Commons has often been a game of catch-up, with substantial social costs. The infant space industry has a rare opportunity to get out ahead, if the right incentives can be put in place.**

satellite tax | common-pool resources | externalities

In 2017, 466 new satellites were launched—more than double the previous year's launches, and more than 20% of all active satellites in orbit in 2017 (1, 2). Rapid space industry growth is projected to continue, driven largely by commercial satellites (Fig. 1). This growth is driving buildup of debris in low-Earth orbit, currently including over 15,000 objects (3). Collision risk from debris is costly; collisions damage or destroy expensive capital assets that are difficult or impossible to repair. Debris buildup could eventually make some low-Earth orbits economically unviable and other orbits difficult or impossible to access (4). In the worst case—though uncertain and occurring over long time horizons—debris growth could become self-sustaining due to collisions between debris objects, a tipping point called Kessler Syndrome (4, 5).

Proposed solutions have so far largely been technological and managerial, aimed at mapping, avoiding, and removing debris (6, 7). These include end-of-life deorbit guidelines and “keep-out” zones for active satellites, and nets, harpoons and lasers to deorbit debris (6). However, with open access to orbits, reducing debris and collision risk incentivizes additional satellite launches, which eventually restores the debris and risk. For instance, if firms were willing to tolerate a 0.1%

annual risk of satellite loss before a technological improvement in debris removal, they will be willing to launch more satellites until the 0.1% annual risk of satellite loss is restored.

Thus, the core of the space debris problem is incentives, not technology. Since satellite operators are unable to secure exclusive property rights to their orbital paths or recover collision-related costs imposed by others, prospective operators face a choice between launching profitable satellites, thereby imposing current and future collision risk on others, or not launching and leaving those profits to competitors. This is a classic Tragedy of the Commons problem (3, 8–10). It can be economically efficiently addressed via incentive-based solutions, such as fees or tradable permits per year in orbit, analogous to carbon taxes or cap-and-trade (8). Incentives should target objects in orbit—rather than launches—because orbiting objects are what directly impose collision risk on other satellites (11). We provide the first analysis, to our knowledge, quantifying the economic benefits of implementing such incentives.

We use a coupled physical-economic model combining rich physical dynamics with satellite economics to quantify the benefits of an internationally harmonized ‘orbital-use fee’ (OUF) relative to a business-as-usual (BAU) open-access scenario, and relative to a scenario with active debris removal. Our model accounts for the effects of each scenario on satellite launch decisions (see Materials and Methods, and Supporting Information). While we focus on an orbital-use fee for analytical convenience, it is conceptually equivalent to other mechanisms for pricing orbits, such as tradable permits.

Our physical model of satellite and debris evolution in orbit

## Significance Statement

The commercial satellite industry is rapidly expanding. A side effect of this expansion is a growing buildup of space debris that imposes costly collision risk on satellite operators. Proposed solutions to this debris have been primarily technological, but the core of the problem is incentives—satellites are being launched without consideration of the collision risks they impose on other operators. We show this incentive problem can be solved with an internationally harmonized ‘orbital-use fee’ (OUF)—a tax on orbiting satellites. Using a coupled physical-economic model, we project an optimally-designed OUF could more than quadruple the long-run value of the satellite industry by 2040.

A.R. and D.K. conceived the project; A.R. performed the analysis, with input from M.G.B. and D.K.; A.R., M.G.B., and D.K. wrote the paper.

The authors have no competing interests.

<sup>2</sup>To whom correspondence should be addressed. E-mail: akhilar@middlebury.edu

obeys relevant accounting identities and utilizes reduced-form approximations of physical processes validated in other works (12, 13). We fit and calibrate the model using data on collision risk and orbital debris produced by the European Space Agency (ESA) (14), and data on active satellites from the Union of Concerned Scientists (UCS) (2) (see Materials and Methods, and Supporting Information). The ESA dataset covers 1958–2017, and the UCS dataset covers 1957–2017. Our physical model assumes that runaway debris growth (Kessler Syndrome) cannot occur, which likely leads our model to understate the benefits of OUFs (see Methods). Our economic model assumes satellites are launched and operated to maximize per-satellite private profits, net of any fees, subject to collision risk. We calibrate the model by fitting the BAU scenario (no fees or debris removal) to historical industry data and launch trends (2, 9) (see Materials and Methods, and Supporting Information).

We project future launch rates to 2040 under the BAU scenario using our fitted model and published projections of future growth of the space economy (15). We then calculate launch rates that would maximize the long-run value of the industry, and we calculate the time series of OUFs that would incentivize these optimal launch rates. The industry value is measured as net present value (NPV)—the long-run value of the entire fleet of satellites in orbit, accounting for both the financial costs of replacing satellites due to natural retirement and collisions, as well as the opportunity cost of investing funds in satellites rather than capital markets. For instance, an NPV of \$1 trillion in 2020 means that the sum total of the stream of net benefits, looking from 2020 into the future and accounting for the timing of the net benefits, is \$1 trillion.

Though our models are deliberately simplified for tractability, they are based on previously-validated approaches to orbital object modeling (12, 13), and our calibrations allow us to reproduce observed trends and magnitudes in the growth of orbital debris and satellite stocks, as well as the calculated collision risk (Figure 3). Nonetheless, our projections should be interpreted as order-of-magnitude approximations that can be refined as needed by more detailed models. In these respects, our approach mirrors integrated assessment modeling approaches that have been useful in developing solutions to other natural resource management problems (e.g., ref (16)).

## Results

We project that shifting from open access to the optimal series of OUFs in 2020 would increase the NPV of the satellite industry from around \$600 billion under business-as-usual to around \$3 trillion—a more than four-fold increase (Fig. 2) (3.61 to 8.4-fold increases in 90% of parameter sets randomly drawn from their calibrated distributions; see Figure 2d). Assuming a 5% market rate of return, an increase of \$2.5 trillion in NPV would be equivalent to annual benefits of approximately \$120 billion in perpetuity. The large immediate increase in NPV that we project in each OUF scenario, relative to BAU (Fig. 2a), comes primarily from the immediate effect of reducing launch activity while the satellite and debris stocks are sub-optimally high (see Supporting Information).

Based on our calculations (see Methods), the optimal OUF starts at roughly \$13,500 per satellite-year in 2020 and escalates at roughly 14.8% per year to around \$244,000 per satellite-year in 2040. Rising optimal price paths are common

in environmental pricing such as carbon taxes (17). The rising price path in this case partly reflects the rising value of safer orbits (resulting in rising industry NPV; see Fig. 2a) created by the OUF. For comparison, the average annual profits of operating a satellite in 2015 were roughly \$2.1 million. The 2020 and 2040 OUF values we describe amount to roughly 0.6% and 11% of average annual profits generated by a satellite in 2015.

Forgone NPV from the satellite industry in 2040—which is the cost of inaction under BAU—escalates from around \$300 billion if optimal management begins in 2025 to around \$750 billion if optimal management begins in 2035. Without OUFs, losses remain substantial even when active debris removal is available. In a best-case analysis where we assume debris removal is costless (i.e., it is provided for free, and requires no additional satellites to implement), debris removal can only recover up to 8.6% of the value lost under open access. At worst, debris-removal can exacerbate orbital congestion via a rebound-type effect, causing additional losses on the order of 1.8% of the value already lost from open access (see Figure 4 and Supporting Information). The inability of debris removal to induce efficient orbit use is driven by open-access launching behavior and underscores the importance of policies to correct economic incentives to launch satellites.

## Discussion

The costly congestion in low-Earth orbit, caused by a buildup of debris and satellites, is fundamentally a problem of incentives—satellite operators currently lack the incentives to factor into their launch decisions the collision risks their satellites impose on other operators. Our analysis suggests that correcting these incentives, via an orbital-use fee, could have substantial economic benefits to the satellite industry, and failing to do so could have substantial and escalating economic costs.

Escalating costs of inaction is a common feature of the Tragedy of the Commons, evident in several other sectors in which it went unaddressed for long periods of time (18). For example, tens of billions of dollars in net benefits are lost annually from open-access or poorly-managed fisheries around the globe (19). Similarly, open access to oil fields in U.S. at the turn of the century drove recovery rates down to 20–25% at competitively drilled sites, compared to 85–90% potential recovery under optimal management (20). Open access to roadways—somewhat analogous to orbits—is estimated to create traffic congestion costs in excess of \$120 billion a year in the U.S. alone (21). In contrast, there is a rare opportunity to get out ahead of the Tragedy of the Commons in the young space industry.

The international and geopolitically complex nature of the space sector poses challenges to implementing orbital-use pricing systems, but these challenges need not be insurmountable. Theory suggests countries could each collect and spend orbital-use fee revenues domestically, without losing economic efficiency, as long as the fee's magnitude was internationally harmonized (22). An example of such a system is the Vessel Day Scheme (VDS) used by the Parties to the Nauru Agreement (PNA) to manage tuna fisheries. Under the VDS, PNA countries each lease fishing rights within their waters, using a common price floor (23). The European Union's (E.U.) Emissions Trading System (ETS) provides an example of an

173 internationally coordinated tradable permit system (24). Notably, each of these pricing programs is built on a pre-existing international governance institution (the Nauru Agreement and the E.U.).

177 An OUF could also be built within existing space governance institutions, such as the Outer Space Treaty (25). For example, Article VI states that countries supervise their space industries, which provides a framework for orbital-use fees to be administered nationally. Article II prohibits national appropriation of outer space, but it does not prohibit private property rights, thus potentially allowing for tradable orbital permitting.

185 Military and other non-commercial uses of satellites are also important to consider in designing any incentive-based orbital management policy. In December of 2018, 430 of 1957 satellites in orbit had acknowledged military users (2). Some ostensibly-commercial satellites may also fill national security needs. Further, governments provide satellite-based public goods to their citizens, such as remote sensing and GIS. In addition to collision risk, satellite and debris buildup may impose costly interference on ground-based space observation systems. OUFs targeting the commercial sector should account for these interests and monitor, as best possible, any "leakage" (see ref (26)) from the commercial sector to the government sector. Antarctica and the Arctic Circle offer two models of internationally managed regions in which commercial interests, environmental interests, and geopolitical interests interact (e.g., ref (27)).

201 Despite the simplifying assumptions of our model, its key qualitative results may be robust to geopolitical and other complexities. Experience in other resource contexts supports our projection that technological or managerial solutions to the space debris problem are unlikely to be as effective as incentive-based solutions. For instance, in fisheries, attempts to restrict entry have led to "capital stuffing," where existing vessels heavily invest in harvest capacity (28). Similarly, catch limits led to the "race to fish" phenomena (29), whereby fishermen exert extreme effort to harvest as much as possible before the catch limit is reached. Attempts to build new highway capacity to reduce traffic congestion have been frustrated because new capacity induces new demand for travel (30). By contrast, policies such as individual tradable quotas (31) and congestion charges (32) directly target the underlying incentive issue and have proven more successful, though are not without some controversy due to distributional effects (33).


218 The Tragedy of the Space Commons is an international incentives problem and therefore needs an internationally coordinated incentive-based solution. Our analysis suggests such solutions could add trillions in value to the industry. Programs such as the EU Emissions Trading System and the PNA's Vessel Day Scheme offer potential models for success.

## 224 SI Datasets.

## 225 Materials and Methods

226 Here we describe the data sources, calibration procedures, and dynamic optimization model used to quantify the economic benefits of orbits under BAU and under optimal management via an OUF.

229 **Data.** We use data on collision risk, orbital debris counts, and satellite counts provided by the European Space Agency (ESA) (14) and the Union of Concerned Scientists (UCS) (2) to calibrate our physical models of aggregate active satellite and debris evolution.



**Fig. 1.** State of the industry (data from refs. 6, 7, 17). a: Projected growth of satellite industry costs and revenues. Revenues and costs are yearly flows. Costs are not annuitized over the satellite lifetime. b: Observed growth in commercial space sector revenues and government space budgets.


We calibrate our economic model of open access launching using historical aggregate returns and costs for the space industry obtained from ref. (9) and launch data calculated from ref. (2). Our physical data cover 1957–2017, while our historical economic data cover 2005–2015.

For each year, the ESA data provides counts of the number of debris objects in orbit within a specified 50 km. altitude band (average orbital altitude) beginning at 100 km. above mean sea level and extending to 2000 km. above sea level, as well as the estimated probability that a collision occurs in the same bands. Each observation in the ESA data corresponds to a year, and each column corresponds to an altitude band. The UCS data describe all known active satellites currently in orbit, including variables for stated purpose, date of launch, and date of deorbit (if applicable). Each observation in this dataset corresponds to an active satellite, with corresponding variables describing that satellite. This dataset is spread over a series of files, released at roughly quarterly to semi-annually frequencies between 2005–2018, from which we reconstruct yearly counts of the number of active satellites in the 100–2000 km. range from 1957–2015.


The economic dataset we use comes from refs. (9) and (15), with data from ref. (9) representing historical data. An observation in the original dataset represents a year, with variables for dollars spent on different space sector activities, e.g. satellite construction and launch services, ground observation services, government spending by different national governments, etc. We classify these entries into yearly revenues flowing to satellite owners, and yearly costs of building, launching, and operating satellites (which we refer to as "launch costs" for brevity). We merge these data with the estimated collision probability data from the ESA dataset (14).

An observation in the final, merged dataset we use for physical model calibration represents a year in 1957–2015, which includes variables for active satellite count, debris count, estimated collision probability, and estimated returns to and costs of launching a satellite in the 100–2000 km. range. An observation in the final, merged dataset we use for economic model calibration represents a year in 2005–2015, with variables for aggregate satellite sector revenues and costs, the yearly gross rate of return on a satellite, the year-over-year change in launch costs, and the estimated collision probability. We describe our data and variable construction procedures in greater detail in the Supporting Information.

**Calibrating the physical model.** We construct the laws of motion for aggregate active satellite and debris stocks from accounting relationships. We assume that a) a constant fraction (3.3%) of undestroyed active satellites naturally deorbit each period (without creating additional debris), b) a constant fraction (49%) of debris deorbits each



**Fig. 2.** Projected gains from optimal management via orbital-use fees. a: NPV of orbit recovery, with optimal management beginning in different years. BAU is shown in black. The NPV gain is the difference between optimal management and BAU. b: Loss in permanent orbit use value from delaying optimal management, relative to 2020 optimal management start. c: Time path of the optimal OUF. d: Distribution of the ratio of optimal NPV in 2040 (assuming optimal management begins in 2020) to BAU NPV in 2040 using 250 draws of alternate parameter sets. The distribution is calculated via a residual-bootstrap resampling procedure which illustrates the effect of calibration uncertainty. We describe the bootstrap in more detail in section 1.4 of the Supporting Information.



**Fig. 3.** Historical and future model simulations. a-d: Time paths of satellites launched, the satellite stock, collision risk, and the debris stock over time in both open access and optimal management simulations. The solid black lines show the observed data from 2006–2015. The vertical dashed black line shows the end of the historical sample in 2015. The red dashed lines show the fit of the open-access model simulation, and the blue dashed lines show the fit of the optimal model simulation. Simulation paths begin from the initial condition of observed satellite and debris levels in 2006 to emphasize that the open-access model is consistent with the observed data from 2006–2015. Simulation procedures and discussion of the discrepancies between observed and modeled outcomes are found in section 1.3 and figure S1 of the Supporting Information.

**Fig. 4.** Model simulations with debris removal. a-d: Time paths of launches, satellites, debris, and collision risk with 50% debris removal beginning in 2029. The red lines show open access paths, while the blue lines show optimal management paths. The dashed lines show time paths with debris removal. The vertical dashed black line marks 2029, when removal begins in the simulation. e: Change in fleet NPV in 2040 with debris removal and open access, relative to a baseline of switching to optimal management in 2020 with debris removal beginning in 2029. f: Summary statistics for changes in open-access welfare loss 50% debris removal for free beginning in each year from 2021–2034. Positive values in panels e and f indicate increased losses due to open access (debris removal makes open access worse), and negative values indicate reduced losses due to open access (debris removal makes open access better). We discuss the intuition for the results in section 1.8 of the Supporting Information.



period, and c) new fragment creation occurs due to launch debris, anti-satellite missile tests, and collisions between orbital objects. We parameterize the probability that an active satellite is destroyed in a collision using an approximate form validated in other works (12, 13), and from that functional form derive an expression for the number of new fragments created in collisions. The parameters in these functions are calibrated separately. Specific formulas are presented in the Supporting Information.

We first calibrate the collision probability function by fitting it to data on the estimated probability of a collision between 100-2000 km provided by ESA, using constrained nonlinear least squares. We constrain the parameter estimates to be positive, which is consistent with the physical interpretation of the collision probability parameters as products of positive values: the magnitude of velocity differences between colliding objects, the total cross-sectional area of the collision, and scaling constants that relate object counts to object densities.

We then calibrate the debris law of motion, using the estimated collision probability parameters, by constrained ridge regression. We use ridge regression to improve the debris model fit, given that we have relatively few observations for the number of parameters. We constrain the parameters to comply with their physical interpretations: parameters representing numbers of fragments created in collisions must be positive, while the debris decay rate must be between 0 and 1. Given the uncertainties involved in modeling and calibrating the potential for Kessler Syndrome, we disallow the possibility of debris objects colliding with each other. This likely makes our conclusions conservative, as it likely understates the growth in collision risk over time due to debris-debris collisions. Consequently, the estimated optimal OUF is likely less than what a model with debris-debris collisions would predict, as are our estimated benefits of implementing the OUF.

Finally, we calibrate the fraction of active satellites that naturally deorbit by fitting the active satellite law of motion using ordinary least squares. This yields an estimated average lifespan of approximately 30 years (the 3.3% deorbit rate mentioned above), which is consistent with an average mission length of 5 years followed by compliance with the IADC's 25-year deorbit guideline (34). We assume full compliance with the guideline to be conservative in our estimates of debris production. We describe our physical calibration procedures in greater detail in Supporting Information. We show sensitivity analyses of our calibrations in SI Figure S3, and of our model projections in SI Figures S4-6. We describe our sensitivity analysis procedure in the Supporting Information.

**Calibrating the economic model.** Economic theory predicts that the collision probability will be determined by the excess rate of return on an active satellite (11). However, the excess rate of return includes the internal rate of return on a satellite asset, which we do not observe, as well as factors relating to the economic structure of the various satellite-using industries involved, which we do not model. To account for these issues, we model the excess rate of return using observed aggregate revenues and costs for the satellite industry, and then include the various unknown and unmodeled factors as parameters to estimate. We then fit this model to the estimated collision probability data using ordinary least squares. Since the signs of the parameters representing the unknown and unmodeled factors are ambiguous, we do not constrain the estimates. The estimated parameters are incorporated into the economic model and used to adjust for unmodeled economic factors. The adjustment produces estimated "implied aggregate costs", shown in Table S5 of the SI. We describe this adjustment procedure and discuss the relevant unmodeled factors in the Supporting Information. We assume a market discount rate of 5% (35). The economic gains from an OUF would be even larger in magnitude at lower social discount rates.

**Projecting open access and optimal launch rates.** To project the launch rates under BAU and optimal management via an OUF, we solve the dynamic optimization problems associated with launching new satellites. Under open access BAU, each firm wishes to maximize their own lifetime satellite profits, and the optimization problem determines each firm's decision of whether to launch or not, which in turn determines the total launch rate across all firms. This involves calculating whether the expected lifetime revenues from

another satellite exceed the cost of launching it and aggregating up from the individual decisions to the total launch rate. Under optimal management, the optimization problem directly determines the total launch rate which maximizes the NPV of the satellite fleet. This involves calculating whether the expected lifetime revenues from another satellite exceed the marginal industry-wide costs of launching it, including the expected costs of replacing other satellites lost due to collision risk created by the new satellite and its associated debris.

Solving these dynamic optimization problems yields "launch policy functions", which map the possible orbital states – satellite and debris counts, given current revenues and costs – into an annual launch rate. As the launch policy functions encode behavioral and institutional assumptions, the launch rates they generate are more realistic approximations of orbit management behavior than simple sensitivity analyses over all possible launch rates. That is, the launch decisions are consistent with, and respond to changes in, economic and orbital conditions. By using the launch policy functions along with the laws of motion for the satellite and debris stocks and an initial condition, we can project the number of satellites and debris in orbit each year under a specified type of management institution. We describe the computation of the launch policy functions in more detail in Algorithms 1-3 in the Supporting Information. Figure 3 compares optimal and open-access projections from the calibrated model assuming no debris removal against the observed data, while figure 4 compares projections with and without debris removal.

**Calculating the optimal orbital-use fee and its benefits.** To calculate the optimal OUF for each year  $t$ , we compare the expected cost of satellite-destroying collisions in year  $t + 1$  under BAU and under optimal management (see Equation 18 in Supporting Information). The OUF is positive whenever the optimal management solution maintains a lower collision probability than what would occur under open access BAU. The OUF is simply the marginal external cost of another satellite launch, which is the additional industry-wide cost of another satellite in orbit (additional collision risk and debris production now and in the future) that is not internalized by individual firms under BAU. By charging firms the marginal external cost of their satellite launch through an OUF, each firm's incentives and thus their launch rates are aligned with those under industry-wide, NPV-maximizing optimal management.

To calculate the benefits of imposing the OUF, we use the launch policy functions described above to compute the NPV of the entire satellite fleet under BAU and optimal management. These NPVs reflect the value of the entire satellite fleet, in perpetuity, assuming society stays on the BAU or optimal management path. The difference between the NPVs yields the gains from implementing the optimal OUF and moving from BAU to the optimal management path.

Although we abstract from many of the economic and physical complications in modeling orbit use, we consider how those factors would affect our analysis in the Supplementary Methods and Supplementary Discussion. In particular, Supplementary Methods details our sensitivity analyses with respect to physical parameter uncertainty, and Supplementary Discussion considers how these concerns may impact our conclusions. Our conclusions are likely robust to the complications we abstract from, with our calculated optimal OUF and the NPV benefits of implementing it providing the correct order of magnitude with the correct qualitative features. Future research will improve our estimates and provide more detailed guidance to policymakers.

**ACKNOWLEDGMENTS.** We acknowledge funding from the University of Colorado Boulder. We thank Francesca Letizia of the European Space Agency for data on debris fragments and collision risk in low-Earth orbit, and Teri Grimwood of the Union of Concerned Scientists for the full catalog of UCS active satellite data. We thank Waleed Abdalati for helpful comments on a previous draft, and attendees of a European Space Agency Space Debris Office seminar for providing helpful feedback on our results.

1. M Wienzierl, Space, the final economic frontier. *J. Econ. Perspectives* 32, 173–192 (2018).

2. Union of Concerned Scientists, Ucs satellite database (2018) From UCS at <https://www.ucsusa.org/nuclear-weapons/space-weapons/satellite-database>. Accessed April 2019.

3. A Witze, The quest to conquer earth's space junk problem. *Nature* **561**, 24–26 (2018).

4. DJ Kessler, BG Cour-Palais, Collision frequency of artificial satellites: The creation of a debris belt. *J. Geophys. Res.* **83**, 2637–2646 (1978).

5. J Liou, N Johnson, Instability of the present leo satellite populations. *Adv. Space Res.* **41**, 1046–1053 (2008).

6. J Pearson, J Carroll, E Levin, Active debris removal: Edde, the electrodynamic debris eliminator. *Space Manuf.* **14** (2010).

7. J Morin, Four steps to global management of space traffic. *Nature* **567**, 25–27 (2019).

8. N Adilov, PJ Alexander, BM Cunningham, Earth orbit debris: An economic model. *Environ. Resour. Econ.* **60**, 81–98 (2015).

9. M Weinzierl, Space, the final economic frontier. *J. Econ. Perspectives* **32**, 173–192 (2018).

10. C Munoz-Patchen, Regulating the space commons: Treating space debris as abandoned property in violation of the outer space treaty. *Chi. J. Int. Law* **18**, 27–44 (2018).

11. A Rao, Ph.D. thesis (University of Colorado at Boulder) (2019).

12. AM Bradley, LM Wein, Space debris: Assessing risk and responsibility. *Adv. Space Res.* **43**, 1372–1390 (2009).

13. F Letizia, C Colombo, H Lewis, H Krag, Extending the ecob space debris index with fragmentation risk estimation. (2017).

14. European Space Agency, Discos database (2018) From ESA at <https://discosweb.esoc.esa.int/web/guest/home>.

15. A Jonas, et al., Space: Investment implications of the final frontier, (Morgan Stanley), Technical report (2017).

16. JR et al, Scenarios towards limiting global mean temperature increase below 1.5 c. *Nat. Clim. Chang.* **8**, 325–332 (2018).

17. WD Nordhaus, To tax or not to tax: Alternative approaches to slowing global warming. *Rev. Environ. Econ. Policy* **1**, 26–44 (2007).

18. GD Libecap, *Contracting for property rights*. (2003).

19. K Kelleher, R Willmann, R Arnason, The sunken billions: the economic justification for fisheries reform, (The World Bank), Technical report (2009).

20. GD Libecap, SN Wiggins, Contractual responses to the common pool: Prorating of crude oil production. *Am. Econ. Rev.* **74**, 87–98 (1984).

21. D Schrank, T Lomax, B Eisele, Urban mobility report, (Texas Transportation Institute, College Station, TX), Technical report (2012).

22. ML Weitzman, Can negotiating a uniform carbon price help to internalize the global warming externality? *J. Assoc. Environ. Resour. Econ.* **1**, 29–49 (2014).

23. E Havice, Rights-based management in the western and central pacific ocean tuna fishery: economic and environmental change under the vessel day scheme. *Mar. Policy* **42**, 259–267 (2013).

24. E Commission, Eu ets handbook, (DG Climate Action), Technical report (2015).

25. UNO for Disarmament Affairs, Treaty on principles governing the activities of states in the exploration and use of outer space, including the moon and other celestial bodies, (UNODA), Technical report (1967).

26. S Felder, TF Rutherford, Unilateral co2 reductions and carbon leakage: the consequences of international trade in oil and basic materials. *J. Environ. Econ. Manag.* **25**, 162–176 (1993).

27. CM Brooks, Competing values on the antarctic high seas: Ccamlr and the challenge of marine-protected areas. *The Polar J.* **3**, 277–300 (2013).

28. RE Townsend, On capital-stuffing in regulated fisheries. *Land Econ.* **61**, 195–197 (1985).

29. AM Birkenbach, DJ Kaczan, MD Smith, Catch shares slow the race to fish. *Nature* **544**, 223–226 (2017).

30. G Duranton, MA Turner, The fundamental law of road congestion: Evidence from us cities. *Am. Econ. Rev.* **101**, 2616–2652 (2011).

31. C Costello, SD Gaines, J Lynham, Can catch shares prevent fisheries collapse? *Science* **321**, 1678–1681 (2008).

32. J Leape, The london congestion charge. *J. Econ. Perspectives* **20**, 157–176 (2006).

33. NC Ban, et al., Diverse fisheries require diverse solutions. *Science* **323**, 338–339 (2009).

34. IADC, Iadc space debris mitigation guidelines, Technical Report IADC-02-01 (2007).

35. LH Goulder, RN Stavins, Discounting: an eye on the future. *Nature* **419**, 673–674 (2002).

# Supporting Information for “Orbital-use fees could more than quadruple the value of the space industry”

Akhil Rao, Matthew G. Burgess, Daniel Kaffine

December 3, 2019

## 1 Extended description of methods and discussion of results

We generate the path of an optimal orbital-use fee (OUF) in three steps. First, we calibrate functions describing the physics and economics of orbit use to match observed data on satellite and debris stock levels and aggregate satellite industry costs and revenues prior to 2015. Then, using the calibrated values, we generate open access and optimal launch paths from 2006 to 2040. Finally, by comparing the open access path of collision risk to the optimal path of collision risk, we calculate the path of the optimal OUf which induces open access satellite owners to internalize the externality they impose on other orbit users.

As we state in the main text, our goal in this article is to provide order-of-magnitude estimates of the optimal OUf and the net present value (NPV) gains from implementing it, and to show the qualitative features of both the OUf path and NPV gains. As we discuss below, our conclusion that a globally-harmonized OUf is necessary to improve the value of the satellite industry is robust to the limitations in our estimation methodology.

We measure the value of the satellite industry as the NPV of cash flows generated by the satellite fleet. NPV accounts for the time value of money — the fact that cash flows received sooner are preferable to cash flows received later — by discounting future cash flows according to the interest that could have been earned on those cash flows had they been received in the present and immediately reinvested in capital markets. In this article, we use the infinite-horizon NPV of the satellite fleet. To calculate this value, we assume that satellite costs and revenues evolve according to the projection in Jonas et al. (2017) until 2050, after which they stabilize at 2050 levels and are constant for the infinite future.<sup>1</sup> The resulting NPVs reflect the sum of short- and long-run costs and returns from the satellite fleet. To express these in terms of equivalent constant cash flows received each year, we convert the NPV into an annuitized present value in the main text.<sup>2</sup> For example, given a discount rate of 5%, a net present value of \$1 trillion in 2020 is equivalent to an annuitized present value of \$47 billion received in perpetuity in each year from 2020 onwards.<sup>3</sup>

We obtain physical functions relating launches, satellites, and debris stocks to collisions, new fragments, and satellite and debris growth from the engineering literature, and economic functions relating the decision to launch to collision risk, costs, and returns from the economics literature. To calibrate the physical functions, we estimate the unknown parameters from satellite stocks, debris stocks, and launches observed over 1957–2014. We constrain the parameters to comply with theoretical restrictions imposed by the engineering model. To calibrate the economic functions, we estimate the unknown parameters from satellite stocks, debris stocks, launches, aggregate satellite industry costs, and aggregate satellite industry returns over 2006–2014.<sup>4</sup> To allow the estimation process to adjust for unobserved launch market frictions, we do not constrain these parameter estimates.

<sup>1</sup>The projection in Jonas et al. (2017) only goes till 2040 — we extend it to 2050 by calculating the compound annual growth rate of the costs and revenues, and assuming that they continue to grow at those rates from 2040 until 2050.

<sup>2</sup>The annuitized present value (PV) of an NPV level is the constant number of dollars received each year such that the discounted sum of annuitized PVs is equal to the target NPV level. Formally, defining the NPV of a stream of uneven cashflows  $x_t$  as  $NPV = \sum_{t=1}^{\infty} \beta^{t-1} x_t = \sum_{t=1}^{\infty} \beta^{t-1} A = A(1 - \beta)^{-1}$  for some constants  $A > 0$  and  $\beta \in (0, 1)$ , the annuitized PV is  $A = NPV(1 - \beta)$ . These conversions facilitate comparisons between uneven streams of cash flows, as they can be expressed in a common unit at a single point in time (NPV terms) or in a common unit at every point in time (annuitized PV terms).

<sup>3</sup>We discuss the interpretation of the discount rate as the opportunity cost of funds invested below.

<sup>4</sup>While we have physical data series extending till 2017, and economic data series covering 2005–2015, we omit observations after 2014 in all calibration data and the 2005 observations in the economic data series. Equations 7 and 15 are the bottlenecks: the former requires omission of the

## 1.1 Data

### 1.1.1 Satellite and debris counts, 1957–2014

We use data on satellites in orbit from the Union of Concerned Scientists’ (UCS) lists of active satellites to construct the satellite stock and launch rate series (Union of Concerned Scientists, 2018). The UCS data provide details on payloads in low-Earth orbit (LEO) and their projected lifetimes. The data are described in Table S1.

Variable	Sample period	Mean	St. Dev.	Min	Max
<i>Active satellite count</i>	1957–2014	1057.9	720.3	4	2271
<i>Launch successes</i>	1957–2014	89.9	47.1	5	334
<i>Launch failures</i>	1957–2014	13.8	9.8	1	42
<i>Payloads decayed</i>	1957–2014	50.8	24.1	3	90
<i>Anti-satellite missile tests</i>	1957–2014	0.2	0.5	0	3
<i>Active debris count</i>	1957–2014	3375.6	2681.3	0	9662
<i>Expected collision rate (active satellites hit)</i>	1957–2014	2.6	1.8	0	6
<i>Historical satellite industry revenues (billion nominal USD)</i>	2006–2014	101.7	20.4	70.4	126.6
<i>Historical satellite industry costs (billion nominal USD)</i>	2006–2014	185.4	40.1	136.2	254.4
<i>Projected satellite industry revenues (billion nominal USD)</i>	2015–2040	217.9	76.5	120.8	366.6
<i>Projected satellite industry costs (billion nominal USD)</i>	2015–2040	362.2	87	210.7	525.1

Table S1: Summary statistics for the data we use to calibrate our physical and economic models and to project open access and optimal launch rates.

We construct the number of active satellites in each year by calculating the number of objects launched in a particular year, adding the number of satellites previously calculated in orbit, and then subtracting the number of satellites listed as having decayed in that year.<sup>5</sup>

Letting  $\ell_t$  be the number of collisions observed in year  $t$  and  $Z_t$  be the number of payloads listed as decayed in  $t$ , we construct the launch rate in  $t$ ,  $X_t$ , from the law of motion for the satellite stock series as

$$\begin{aligned} S_{t+1} &= S_t - Z_t - \ell_t + X_t \\ \implies X_t &= S_{t+1} - S_t + Z_t + \ell_t, \end{aligned} \quad (1)$$

where  $S_t$  is the number of active payloads in  $t$  and  $Z_t$  is the number of payloads listed as decayed in  $t$ .

The debris and collision risk series’ we use were provided by the European Space Agency. We use debris data from the DISCOS database (European Space Agency, 2018) and collision probability data used in Letizia, Lemmens, and Krag (2018) (the variable  $p_c$  in that paper). We use only objects with a semi-major axis of 2000km or less in all our data series. We prefer to use the DISCOS fragment data rather than the Space-Track fragment data (Combined Space Operations Center, 2018) as DISCOS attributes fragments to the time they were created or detected, whereas the Space-Track data attributes fragments to the time their parent body was launched. The DISCOS attribution method is closer to how economic agents in our model receive information and make decisions. Given the difficulties in determining operational status, the collision probability estimates account for the probability of collisions with all

first year’s data to construct the cost growth variable, while the latter cannot be used to construct the necessary implied cost growth for the final year. As a result, our economic data calibration only uses data covering 2006–2014. Since our economic model implies that the number of objects in orbit and the collision risk convey economic information about returns and costs, we omit physical variables after 2014 from our physical model calibration to avoid unintended data snooping.

<sup>5</sup>This procedure is likely to produce an upward-biased estimate of the returns-generating satellite stock in any given year, since satellites which are no longer operational will not be removed from the estimated stock until they have deorbited. Thus, the satellite stock in this procedure includes some objects which are, economically speaking, “socially-useless debris”. We use this procedure despite the attribution issue for two reasons. First, we do not have data on when specific satellites were declared nonoperational by their owners. Such a determination can be particularly tricky when a mission has ended, but the satellite still has fuel and could be repurposed for another mission. Second, to the extent that our estimates of the satellite stock are biased upward (toward positive infinity), our physical and economic parameters estimates will be biased downwards. The downward bias in economic parameters will deflate both the open access and socially optimal launch rates, while the downward bias in physical parameters will inflate both the open access and socially optimal launch rates, with the net effect being difficult to determine. However, the downward bias in our estimated collision risk coefficients and the upward bias in our estimated satellite stock will bias our estimated OUF downward, so that it is a lower bound.



intact bodies. This produces an upward-biased estimate of the probability of collisions with only operational satellites. This upward bias likely deflates the number of open access and optimal launches we project. However, since the open access and optimal launch rates are chosen to equate collision risk with measures of economic returns (described in equations 6 and 9), the resulting estimated OUF paths will not be biased upward by the same degree as the collision probability estimates.

### 1.1.2 Aggregate satellite industry returns and costs, 2006–2040

We use data on historical satellite industry revenues from Wienzierl (2018), and UCS data on satellites in LEO (semi-major axis less than 2000km) (Union of Concerned Scientists, 2018). The economic data provide a breakdown of revenues across satellite manufacture, launch, insurance, and products and services. The satellite industry revenues data cover 2006–2015, while the active satellites data cover 1957–2014. To generate launch rate and OUF projections out to 2040, we use revenue and cost projections from Jonas et al. (2017). These projections are shown in Figure 1a of the main text. The historical and projected data we use are described in Table S1.

We calculate the per-period returns on owning a satellite ( $\pi_t$ ) as the revenues generated from commercial space products and services, and the per-period costs of launching a satellite ( $F_t$ ) as the sum of revenues from commercial infrastructure and support industries, ground stations and equipment, commercial satellite manufacturing, and commercial satellite launching. The ratio  $\pi_t/F_t$  then gives a time series of the rate of return on a single satellite, as the number of satellites cancels out of the numerator and denominator.<sup>6</sup> Since the numbers provided in Wienzierl (2018) are for the satellite industry as a whole, the ratio still needs to be adjusted to represent satellites in LEO. We do not explicitly conduct this adjustment, but let the adjustment be calculated during the estimation of equation 7.<sup>7</sup>

Note that the data we use for  $\pi_t$  and  $F_t$  are industry-level aggregates, rather than satellite-level figures. To convert the data from industry-level figures to per-satellite figures, we must apply a scaling factor which “disaggregates” the data. This unknown factor is common to both cost and revenue aggregates, and so cancels out of  $\pi_t/F_t$  such that the ratio correctly represents the rate of return per satellite. Since we use  $\pi_t/F_t$  to compute the open access and optimal policy and value functions, the unknown scaling factor does not affect our solutions (launch rates). However, it does affect our calculated time paths of NPV under business-as-usual (BAU, i.e. open access) and optimal management as well as the OUF, since those values do not involve ratios which would cancel out the unknown scaling factor. Since the unknown scaling factor is on the order of the reciprocal of the number of satellites in orbit (or projected to be in orbit) in each period, we proxy for it in our projected time paths by dividing by the BAU satellite stock path. This choice of proxy does not affect our qualitative results or the order of magnitude of the OUF or NPV under BAU and optimal management — using alternate proxies such as the optimal management path of satellites or the observed number of satellites in 2015 produces similar results, although our proxy results in more conservative projections than those alternative choices.

## 1.2 Models

### 1.2.1 Orbital mechanics with limited lifespans, missile tests, and certainty

Our physical model uses physical accounting relationships in the aggregate stocks of satellites and debris for the laws of motion, and draws on (Letizia et al., 2017) for the functional forms of the new fragment creation and collision probability functions  $G(S, D)$  and  $L(S, D)$ . The time scale is set as one calendar year to match our data.  $S_t$  denotes the number of active satellites in an orbital shell in period  $t$ ,  $D_t$  the number of debris objects in the shell in  $t$ ,  $X_t$  the number of satellites launched in  $t$ ,  $L(S_t, D_t)$  the probability that an active satellite in the shell will be destroyed in a collision in  $t$ ,  $\mu$  is the fraction of satellites which do not deorbit in  $t$ , and  $m$  is the average amount of debris generated by launching satellites (such as rocket bodies).  $\delta$  is the average proportion of debris objects which deorbit in  $t$ , and  $G(S_t, D_t)$  is the number of new debris fragments generated due to all collisions between satellites and debris.<sup>8</sup>  $A_t$  is

<sup>6</sup>This is true whether the satellites were launched individually on separate rockets or in groups on the same rockets.

<sup>7</sup>Another way to perform this adjustment is by calculating the yearly share of satellites in LEO and multiplying the ratio  $\pi_t/F_t$  by the share in LEO. This approach is difficult to generalize to future years since it requires projections of satellites in other orbits. It is also not clear that the returns of satellites in LEO are truly proportional to the LEO share of the total number of active satellites in all orbits; it seems more likely that LEO satellites earn less revenue per satellite than geostationary satellites.

<sup>8</sup>For most of our sample, the number of observed collisions is zero. We use the probability of collisions in our models rather than the observed number for two reasons. First, it proxies for unobserved collisions, including non-catastrophic ones. Second, a model with stochastic collisions

the number of anti-satellite missile tests conducted in  $t$ , and  $\gamma$  is the average number of fragments created by one test. We assume that satellites which deorbit do so without creating any additional debris.

The number of active satellites in orbit is modeled as the number of launches in the previous period plus the number of satellites which survived the previous period (also shown in equation 1). The amount of debris in orbit is the amount from the previous period which did not decay, plus the number of new fragments created in collisions, plus the amount of debris in the shell created by new launches.<sup>9</sup> Formally,

$$S_{t+1} = S_t(1 - L(S_t, D_t))\mu + X_t \quad (2)$$

$$D_{t+1} = D_t(1 - \delta) + G(S_t, D_t) + \gamma A_t + mX_t. \quad (3)$$

For simplicity, we assume all non-operational satellites are immediately deorbited, making our OUF estimates conservative. Letizia et al. (2017) use an analogy to kinetic gas theory to parameterize the probability of a collision as a negative exponential function, with the density of colliding objects one of the arguments of the exponential function. We therefore parameterize  $L(S_t, D_t)$  as

$$L(S_t, D_t) = 1 - \exp(-\alpha_{SS}S_t - \alpha_{SD}D_t), \quad (4)$$

where  $\alpha_{SS}$  and  $\alpha_{SD}$  include the difference in velocities between the objects colliding, the total cross-sectional area of the collision, and scaling parameters which relate the number of objects to their density in the volume. We use these probability functional forms to parameterize  $G(S_t, D_t)$  as

$$G(S_t, D_t) = \beta_{SS}(1 - \exp(-\alpha_{SS}S_t))S_t + \beta_{SD}(1 - \exp(-\alpha_{SD}D_t))S_t, \quad (5)$$

where the  $\beta_{jk}$  parameters are interpreted as “effective” numbers of fragments from collisions between objects of type  $j$  and  $k$ .<sup>10</sup> We refer to the  $\alpha_{jk}$  and  $\beta_{jk}$  as “structural physics parameters”, as they represent physical entities which are exogenous to our model.

We ignore the possibility of collisions between debris objects for two reasons. First, the data we have do not allow us to identify the effective number of fragments from such collisions, or the probability of such collisions, using our calibration approach. Second, our focus here is not on the probability of Kessler Syndrome, but on launch patterns and their response to the extant stock of orbiting satellites and debris. Our estimates of the optimal OUF path and the benefits of implementing it are likely understated due to this omission. Incorporating the possibility of Kessler Syndrome is an important piece of optimal orbit use analysis and policy design, and will likely require higher-fidelity physical modeling than the “aggregate calibration” approach we take here. This is an important area for future research.

Equations 3, 4, and 5 can be viewed as reduced-form statistical models which recreate the results of higher-fidelity physics models of debris growth and the collision probability. While higher-fidelity physics models may use similar functional forms, the key difference between our approach and the approach in such models is how we calibrate the models: rather than derive the appropriate parameter values from physical first principles given the data, we estimate the values of those parameters which maximize the fit between the data and model-predicted collision probabilities, satellite evolution, and debris stocks. Though our approach is computationally convenient, it likely sacrifices some predictive power.

### 1.2.2 Open access orbit use with time-varying aggregate returns and costs

The economic model of open access here is based on the model of open access in Rao (2019) to determine the satellite launch rate under open access,  $X_t$ , as a function of the collision probability,  $L(S_{t+1}, D_{t+1})$ , and the excess return on a

complicates the process of solving for the optimal time path by adding another state variable to the dynamic programming algorithm. As the number of objects in a single period increases, the fraction of satellites destroyed in collisions in that period converges to the probability of destruction, so this assumption provides a “mean field”-type approximation.

<sup>9</sup>Empirically, we only consider the population of trackable fragments, i.e. those with size greater than 5-10 cm in LEO.

<sup>10</sup>“Effective” numbers of fragments measure the number of new fragments weighted by the time they spend inside the volume of interest. This approach is used in the debris modeling literature, for example Bradley and Wein (2009).

satellite,  $r_s - r$ .<sup>11</sup> In the simplest case, where all of the economic parameters are constant over time, the open access launch rate equates the collision probability with the excess return:

$$L(S_{t+1}, D_{t+1}) = \underbrace{r_s - r}_{\text{excess return on a satellite}}, \quad (6)$$

where  $r_s$  is the per-period rate of return on a single satellite ( $\pi/F$ , where  $\pi$  is the per-period return generated by a satellite and  $F$  is the cost of launching a satellite, inclusive of non-launch expenditures such as satellite manufacturing and ground stations) and  $r$  is the risk-free interest rate.<sup>12</sup>

Equation 6 can therefore also be used to calculate the implied internal rate of return (IRR) for satellite investments from observed data on collision risk and satellite returns.  $r$  is not observed in our data. When costs and returns are time-varying, equation 6 becomes

$$\begin{aligned} L(S_{t+1}, D_{t+1}) &= 1 + r_{s,t+1} - (1+r) \frac{F_t}{F_{t+1}} \\ \Rightarrow L(S_{t+1}, D_{t+1}) &= \underbrace{\left( r_{s,t+1} - r \frac{F_t}{F_{t+1}} \right)}_{\text{excess return on a satellite}} + \underbrace{\left( 1 - \frac{F_t}{F_{t+1}} \right)}_{\text{capital gains from open access and satellite launch cost variation}} \end{aligned} \quad (7)$$

where  $r_{s,t+1} = \pi_{t+1}/F_{t+1}$ . With time-varying economic parameters, two sources of returns drive the collision risk. One is the excess return realized in  $t+1$  from launching a satellite in  $t$ . The other is the capital gain (or loss) due to open access and the change in satellite costs. Since open access drives the value of a satellite down to the total cost of launching and operating it,  $F_t$  becomes the cost of receiving  $F_{t+1}$  in present value the following period, and the returns are given as percentages of  $F_{t+1}$ . Since the discount rate is unobserved, we fix it to be constant over time to facilitate estimation.<sup>13</sup> While we abstract from the fact that satellite lifetimes are finite, this extension was considered in Rao (2019) and shorter planned operational lifetimes were shown to reduce the expected collision risk. We discuss this issue further when describing our calibration methodology in section 1.3.2, including why it is unlikely to affect our estimates of the optimal OUF and the benefits of implementing it.

### 1.2.3 Optimal orbit use with time-varying aggregate returns and costs

Determining the launch plan to ensure optimal orbit use is more complicated. Economists commonly refer to this type of problem as “the (fleet) planner’s problem”, imagining a planner tasked with maximizing fleet-wide NPV. The fleet planner launches satellites to maximize the value of the entire fleet into the (discounted) infinite future, subject to the laws of motion of satellite and debris stocks. Formally, letting  $\beta = (1+r)^{-1}$  be the discount factor, the planner solves

$$\begin{aligned} W(S, D) &= \max_{X \geq 0} \{ \pi S - FX + \beta W(S', D') \} \\ S' &= S(1 - L(S, D))\mu + X \\ D' &= D(1 - \delta) + G(S, D) + \gamma A + mX. \end{aligned} \quad (8)$$

We drop time subscripts and use primes on a variable’s right to indicate future values, in keeping with the convention for infinite-horizon dynamic programming problems. The economic parameters  $\pi$  and  $F$  are allowed to be time-varying in our solution approach, though all other physical and economic parameters are constant over time.

<sup>11</sup>While we consider LEO as a whole, this approach could be generalized to individual spherical shells within LEO. Such generalization could incorporate the substantial heterogeneity in orbital-use values. For example, some orbits are more valuable because they offer ideal conditions for Earth observation, and will likely need a different fee schedule than orbits which do not offer such conditions.

<sup>12</sup>More precisely,  $r$  is the opportunity cost of funds invested in launching a satellite, and may diverge from the risk-free rate if the satellite launcher’s most-preferred alternate investment is not a risk-free security. This rate is sometimes referred to as the internal rate of return.

<sup>13</sup>This equation was derived in the Appendix of Rao (2019). In that setting the discount rate was not constant over time.

Solving the planner's problem by taking the first-order condition and applying the envelope condition to recover the unknown functional derivatives, we obtain an expression for the collision risk in period  $t + 1$  which characterizes the optimal launch rate in period  $t$ :

$$L(S_{t+1}, D_{t+1}) = 1 + r_{s,t+1} - (1 + r) \frac{F_t}{F_{t+1}} - \xi(S_{t+1}, D_{t+1}), \quad (9)$$

where

$$\begin{aligned} \xi(S_{t+1}, D_{t+1}) = & S_{t+1} L_S(S_{t+1}, D_{t+1}) F + \frac{\pi - rF - L(S_t, D_t) F - S_t L_S(S_t, D_t) F}{\beta(1 - \delta + G_D(S_{t+1}, D_{t+1}) + mL_D(S_{t+1}, D_{t+1}) S_{t+1})} \\ & - \frac{\beta G_S(S_{t+1}, D_{t+1}) + m(1 - L(S_{t+1}, D_{t+1}) - S_{t+1} L_S(S_{t+1}, D_{t+1}))}{\beta(1 - \delta + G_D(S_{t+1}, D_{t+1}) + mL_D(S_{t+1}, D_{t+1}) S_{t+1})} L_D(S_{t+1}, D_{t+1}) S_{t+1} F \end{aligned} \quad (10)$$

is defined as the marginal external cost of another satellite. These equations are derived in Section 1.7.

### 1.3 Calibration and simulation

#### 1.3.1 Physical parameters: deorbit, decay, collisions, and fragments

We calibrate the rate at which satellites deorbit,  $\mu$ , by estimating the following analog to equation 1 by ordinary least squares (OLS):

$$S_{t+1} = S_t(1 - L(S_t, D_t))\hat{\mu} + X_t + \varepsilon_{S_t}. \quad (11)$$

We use hats over variables to indicate a parameter being estimated, e.g.  $\mu$  is the true (unknown) value while  $\hat{\mu}$  is the estimate of  $\mu$ .  $\varepsilon_{S_t}$  is the error term.

Equation 11 yields an estimated average lifespan of about 30 years, i.e.  $\hat{\mu} = 0.967$ . This is consistent with an average mission length of 5 years, followed by compliance with the 25-year deorbit guideline issued by the IADC (IADC, 2007). Using this rate in our forward simulations is conservative in the sense that the estimated OUF becomes a lower bound relative to a model with imperfect compliance. Bradley and Wein (2009) show full compliance to be optimal.

We calibrate equations 3 and 4 by estimating the following equations:

$$L(S_t, D_t) = 1 - \exp(-\hat{\alpha}_{SS} S_t - \hat{\alpha}_{SD} D_t) + \varepsilon_{L_t} \quad (12)$$

$$D_{t+1} = (1 - \hat{\delta}) D_t + \hat{\beta}_{SS}(1 - \exp(-\hat{\alpha}_{SS} S_t)) S_t + \hat{\beta}_{SD}(1 - \exp(-\hat{\alpha}_{SD} D_t)) S_t + \quad (13)$$

$$\hat{\gamma} A_t + \hat{m} X_t + \varepsilon_{D_t}. \quad (14)$$

Theory predicts  $\alpha_{jk}$ ,  $\beta_{jk}$ , and  $m$  are nonnegative, and  $\delta$  is in  $(0, 1)$ . We constrain the parameter estimates to comply with the theoretical restrictions.

We estimate equations 12 and 13 in two stages. First, we estimate equation 12 by constrained nonlinear least squares (NLS). Then, using the estimated values of  $\hat{\alpha}_{SS}$  and  $\hat{\alpha}_{SD}$  to generate  $(1 - \exp(-\hat{\alpha}_{SS} S_t)) S_t$  and  $(1 - \exp(-\hat{\alpha}_{SD} D_t)) S_t$ , we estimate equation 13 by constrained ridge regression, estimating  $(1 - \delta)$  directly.<sup>14</sup> We estimate both equations on the sample from 1957-2014. The fitted values for all three equations are shown against the actual values in Figure S1.

Tables S2 and S3 show the calibrated parameters for equations 12 and 13. Since our goal is predicting the time series rather than inference on the physical coefficients, we show prediction uncertainty estimates for equations 12 and 13 rather than coefficient standard errors in Figure S3.

<sup>14</sup>We use ridge regression for the debris equation to improve the model fit, despite bias in the estimated parameters. Ridge estimates are biased toward zero relative to OLS estimates. For a given penalty parameter  $\lambda \geq 0$ , the relationship between a ridge coefficient estimate  $\hat{\beta}^{\text{ridge}}$  and the corresponding OLS estimate  $\hat{\beta}^{\text{OLS}}$  is  $\hat{\beta}^{\text{ridge}} = \hat{\beta}^{\text{OLS}} / (1 + \lambda)$ .

<b>Collision probability parameters</b>	$\alpha_{SS}$	$\alpha_{SD}$
<i>Parameter values</i>	1.29e-06	2.56e-08

Table S2: Parameter values from estimating equation 12.

<b>Debris law of motion parameters</b>	$\delta$	$m$	$\gamma$	$\beta_{SS}$	$\beta_{SD}$
<i>Parameter values</i>	0.49	4.84	144.13	292.72	5026.17

Table S3: Parameter values from estimating equation 13. All values are rounded to two decimal places. The penalty parameter  $\lambda$  was selected through cross-validation.

The estimated physical parameter values are physically plausible, with the values estimated for equation 13 being lower bounds due to ridge estimation bias. For example, the value of  $m$  suggests that every satellite launched creates (at least) 4.84 pieces of debris on average, while the value of  $\gamma$  suggests that anti-satellite missile tests create (at least) 144.13 pieces of debris on average. While higher-fidelity physical models which derive these quantities from first principles will yield more accurate results, the estimated values appear to be a reasonable first-order approximation to the true values based on the model fits (shown in Figure S1).

### 1.3.2 Economic parameters: returns, costs, and discounting

Since the discount rate  $r$  is unobserved, we calibrate equation 7 by estimating

$$L(S_t, D_t) = a_{L1} + a_{L2}r_{st} + a_{L3}\frac{F_{t-1}}{F_t} + \varepsilon_{rt}, \quad (15)$$

using OLS on the sample of returns data from 2005-2015, omitting the first observation (for 2005) to construct  $F_{t-1}/F_t$ .  $\varepsilon_{rt}$  is a mean-zero error term,  $a_{L2}$  is a scale parameter, and  $a_{L3}$  measures the gross IRR,  $1 + r$ . We do not explicitly incorporate satellite lifetimes net of exogenous failure (the parameter  $\mu$ ) as the coefficient is not separately identifiable — it enters each term in equation 15 as a scalar multiplying the parameters  $(a_{L1}, a_{L2}, a_{L3})$ , and does not affect the model's predictions. Table S4 shows the calibrated parameters.

<b>Economic calibration parameters</b>	$a_{L1}$	$a_{L2}$	$a_{L3}$
<i>Parameter values</i>	0.004	0.009	-0.0004
<i>Standard errors</i>	0.002	0.002	0.001

Table S4: Parameter values from estimating equation 15. All values are rounded to the first non-zero digit.

If our data perfectly measured the costs and returns of satellite ownership, and our theoretical model held exactly, we would expect  $a_{L1} = 1$ ,  $a_{L2} = 1$ , and  $a_{L3} < -1$ . Our estimates therefore suggest that our returns and cost series are measured with error or that our theoretical model is missing some important factors, such as constraints on the number of launches possible each period. In this situation, using the raw economic data with the theoretical model could yield overly-pessimistic predictions. To account for this, we adjust our satellite launch cost data to be consistent with the simple open access model by using the estimated parameter values from the economic calibration. Table S5 compares the adjusted data to the original series. The adjusted cost data are of the same magnitude as the unadjusted data, but typically smaller. This suggests that the unmodeled factors include cost efficiencies in satellite production, launch, and management, or constraints on launch services available each period, which are consistent with the analytical features we abstract from in our theoretical model.

The open access model described so far assumes that any firm which wants to launch a satellite can do so. If launches are limited, as they are in practice, this assumption will be violated. The limits will prevent open access launching from equating the excess return on a satellite with the risk of its destruction. In this way, firms which are able to launch earn rents from having a satellite while the collision risk is below the excess return. The wedge between the collision risk and excess return will reflect the value of those rents. To get a sense of how a binding



launch constraint would affect our estimates, we adapt the flow-controlled equilibrium condition from Rao (2019) to our situation with time-varying parameters.

$$L(S_{t+1}, D_{t+1}) = \left(1 - \frac{p_t}{F_{t+1} + p_{t+1}}\right) + \frac{\pi_{t+1}}{F_{t+1} + p_{t+1}} - (1 + r_t) \frac{F_t}{F_{t+1} + p_{t+1}}, \quad (16)$$

$p_t$  can be interpreted in two ways. It can be viewed as the implied rent received by a firm which already owns a satellite in  $t$  due to launches in  $t$  being restricted. It can also be viewed as the implied launch fee paid by a firm which is allotted a launch slot in  $t$ . In either view, a binding launch constraint results in positive values of  $p_t$  and  $p_{t+1}$ , biasing the coefficients from regression 7.  $a_{L1}$  is biased toward negative infinity, while  $a_{L2}$  and  $a_{L3}$  are biased toward zero. If the data were free of measurement error and an atomistic homogeneous open access model with a constant discount rate  $r$  was correct, we would have  $a_{L3} = -(1 + r)$ . We set the discount rate to be 5% ( $r = 0.05$ , implying a discount factor of  $\beta = (1 + r)^{-1} \approx 0.95$ ).

Regardless of the factors missing from the theoretical model, we use equation 15 to recursively calculate the sequence of launch costs implied by the combination of open access, observed launch rates, and observed satellite returns as

$$\begin{aligned} L(S_t, D_t) &= a_{L1} + a_{L2}r_{st} + a_{L3} \frac{F_{t-1}}{F_t} + \varepsilon_{rt} \\ \implies \hat{F}_t &= \frac{a_2\pi_t + a_3F_{t-1}}{L_t - a_1}. \end{aligned} \quad (17)$$

This “adjusted cost” accounts for these missing factors for the historical period. When it exceeds the observed costs, the missing factors tend to reflect unobserved costs to launching satellites, such as limited launch availability. When it is below the observed costs, the missing factors tend to reflect unobserved returns to launching satellites, such as returns to scale.

Table S5 shows the observed satellite returns ( $\pi_t$ ), observed launch costs ( $F_t$ ), and implied launch costs ( $\hat{F}_t$ ). The adjusted costs can be interpreted as the costs such that observed launch patterns result from open access under a non-binding launch constraint and with no other unmeasured factors impacting launch costs. Using the adjusted costs instead of the observed costs in the historical sample ensures that our parameter estimates are consistent with our model of open access.<sup>15</sup>

Year	Observed aggregate cost	Implied aggregate cost
2006	161.02	194.26
2007	185.5	178.88
2008	170	154.86
2009	137.81	131.47
2010	136.16	140.48
2011	166.99	149.09
2012	186.88	165.69
2013	215.9	170.79
2014	254.39	170.68

Table S5: Launch costs (inclusive of satellite construction, integration, and maintenance) implied by open access and observed revenues and costs. Since aggregate launch costs do not reflect the various unmodeled frictions (changing shadow price of the launch constraint, economies of scale in satellite launching, and the share of satellites which are in LEO), we calibrate the economic costs to the collision probability and use the calibration to back out implied costs that reflect these frictions. All values are given in billions of nominal US dollars per year. 2015 is omitted due to the recursive nature of equation 17.

### 1.3.3 Algorithms for open access and optimal policy functions

We generate two sequences of policy functions: one function for each period under consideration, and one sequence for each management regime type. We compute each sequence through backwards induction: beginning at the final

<sup>15</sup>Estimating the economic parameters using an open access model with a binding launch constraint is challenging as equation 6 becomes an inequality, giving sets of open access-consistent parameters rather than a unique combination.

period in our projection horizon, and iteratively working backwards to the initial period. This procedure implies “perfect foresight” planning under each management regime, i.e. that all agents under any management regime are able to perfectly forecast the sequence of returns, costs, interest rates, launch rates, and other model objects. The perfect foresight assumption is clearly unrealistic, but our purpose is not to show how uncertainty in economic parameters propagates over time. Rather, our purpose is to show how an optimal OUF would vary over time and the time paths of orbital aggregate stocks under different management regimes. Such assumptions are used in integrated assessment models of climate change with similar rationales, e.g. the models studied in Kelly and Kolstad (1999); Nordhaus (2013); Wilkerson et al. (2015). Our work here is conceptually similar to integrated assessment modeling.

To compute the open access time path, we first generate a grid of satellite and debris levels,  $(grid_S, grid_D)$ . We generate this grid as an expanded Chebyshev grid to reduce numerical errors from interpolation, provide higher fidelity near boundaries, and economize on overall computation time. In contrast to a standard Chebyshev grid, an expanded Chebyshev grid allows for computation (rather than extrapolation) at the boundary points. The formula for the  $k^{th}$  expanded Chebyshev node on an interval  $[a, b]$  with  $n$  points is

$$x_k = \frac{1}{2}(a+b) + \frac{1}{2}(b-a) \sec\left(\frac{\pi}{2n}\right) \cos\left(\frac{k}{n} - \frac{1}{2n}\right)$$

We set different values of  $b$  for  $S$  and  $D$ , creating a rectangular grid. The main issue in setting  $b$  is ensuring that the time paths we solve for (described in section 1.3.4) do not run into or beyond the boundary. To avoid this issue while minimizing the number of points in regions the time paths never reach, we set different  $b$  bounds for open access and the optimal plan, with the open access grid being strictly larger in both dimensions than the optimal plan grid.

In general, computing decentralized solutions under open access is simpler than computing the planner’s solutions. This is because open access simplifies the continuation value to the cost of launching a satellite. We use R for all simulations, parallelizing where possible. To facilitate convergence of policy and value functions, we normalize the returns and costs parameters so that  $\pi_1 = 1$  during computation, and rescale the parameters after the time paths have been generated.

We compute optimal value functions by value function iteration on the grid. We initialize the algorithm with a guess of the value and policy functions. Then, at each point on the grid, we solve the first-order condition for the planner’s problem (equation 9). Since there may be multiple solutions, only one of which leads to a global maximum, we then evaluate the value function at each solution (including zero) and select the launch rate attached to the largest level of the value function. Algorithm 1 describes how we compute the optimal policy and value functions for a given grid and given value function guess  $guess(S, D)$ , while algorithm 2 describes how we compute the open access policy and value functions.

We construct our initial guess of the planner’s value function as the terminal value of the fleet. In the penultimate period, we assume it is not optimal to launch any satellites ( $X_{T-1}^* = 0$ ), making the final fleet size

$$S_T = S_{T-1}(1 - L(S_T, D_T)).$$

In the final period ( $T$ ), the payoff of the fleet is  $\pi_T S_T$ . Our assumption that it is not optimal to launch any satellites in the penultimate period implies the one-period returns of a satellite do not cover the cost of building and launching ( $\beta\pi_T < F_{T-1}$ ), which we verify to hold in every period of our data. We use the implied series of  $F_t$  given the observed  $\pi_t$  and launch rate series in solving for open access and optimal policies.

### 1.3.4 Projected time paths

We use algorithms 1 and 2 to compute policy and value functions in each period, and run them sequentially from the final period to the first period to generate a series of policy and value functions for each period’s set of economic parameters. Algorithm 3 describes this process.

It is important to note that when obtaining the sequence of policy functions we do not do backwards induction within each economic time period prior to the final period. Instead, we hold the continuation value ( $W(S_{t+1}, D_{t+1})$ )

---

**Algorithm 1:** Value function iteration

---

1 Set

$$W_0(S, D) = \text{guess}(S, D), \\ X_0 \equiv 0$$

for all  $(S, D) \in (\text{grid}_S, \text{grid}_D)$

2 Set  $i = 1$  and  $\delta = 100$  (*some large initial value*).

3 **while**  $\delta > \epsilon$  **do**

4     At each grid point in  $(\text{grid}_S, \text{grid}_D)$ , use a numerical rootfinder to obtain

$$X_i : W_D(S, D) + SL_D(S, D)\hat{F} - \beta[1 - \delta + G_D(S, D) + mSL_D(S, D)]W_D(S', D') = 0,$$

where  $W_D(S, D)$  is given by equation 37.

5     Evaluate  $W_i(S, D) = \pi S - \hat{F}X_i + \beta W_{i-1}(S', D')$  at  $X_i \cup \{0\}$ , and select whichever value of  $X_i \cup \{0\}$  maximizes  $W_i(S, D)$ .  $W_{i-1}(S', D')$  is computed by linear interpolation.

6      $\delta \leftarrow \|W_i(S, D) - W_{i-1}(S, D)\|_\infty$ .

7      $i \leftarrow i+1$

8 **end**

---

---

**Algorithm 2:** Open access launch plan computation

---

1 Use a numerical rootfinder to find the  $X_{t-1}^o$  which solves

$$L(S_t, D_t) = a_{L1} + a_{L2}\hat{f}_{st} + a_{L3}\frac{\hat{F}_{t-1}}{\hat{F}_t},$$

using the estimated laws of motion for  $S_t$  and  $D_t$  as functions of  $X_{t-1}$ , and the estimated function for  $L(S_t, D_t)$ .

2 Approximate  $W_i^\infty(S, D) = \sum_{t=1}^\infty \beta^{t-1}(\pi S_t - \hat{F}X_t^o)$  as  $W_i^T(S, D) = \sum_{t=1}^{T-1} \beta^{t-1}(\pi S_t - \hat{F}X_t^*)$  by backwards induction, using the estimated laws of motion for  $S_{t+1}$  and  $D_{t+1}$  and the estimated function for  $L(S_t, D_t)$ . We use  $T = 500$ .

---

---

**Algorithm 3:** Generating a perfect-foresight sequence of policy functions

---

```

1 Set economic parameters to final period values.
2 Run algorithm 1 (for an optimal path) or 2 (for an open access path) to obtain  $W_T(S_T, D_T)$ .
3 for  $t$  in  $T-1:1$  do
4   Set  $i = 1$  and  $\delta = 100$  (some large initial value). Set  $X_{0t} = X_T$ . while  $\delta > \varepsilon$  do
5     Using the value function from the previous time step as  $W_{t+1}(S_{t+1}, D_{t+1})$ , calculate
        
$$X_{it}^* : W_{Dt}(S_t, D_t) + S_t L_D(S_t, D_t) \hat{F}_t -$$


$$\beta [1 - \delta + G_D(S_t, D_t) + m S_t L_D(S_t, D_t)] W_{Dt}(S_{t+1}, D_{t+1}) = 0,$$

        (for an optimal path, where  $W_{Dt}(S, D)$  is given by equation 37),
        or
        
$$X_{it}^o : L(S_{t+1}, D_{t+1}) = a_{\ell 1} + a_{\ell 2} \hat{F}_{st} + a_{\ell 3} \frac{\hat{F}_{t-1}}{\hat{F}_t} \quad (\text{for an open access path}),$$

        using the estimated laws of motion for  $S_{t+1}$  and  $D_{t+1}$ , the estimated function for  $L(S_t, D_t)$ , linearly
        interpolating to compute  $W_{t+1}(S_{t+1}, D_{t+1})$ .
6      $\delta \leftarrow \|X_{it} - X_{i-1t}\|_\infty$ 
7   end
8   If calculating an optimal path, set  $W_t(S_t, D_t) = \pi_t S - \hat{F}_t X^* + W_t(S_{t+1}^*, D_{t+1}^*)$ . If calculating an open access
        path, set  $W_t(S_t, D_t) = \pi_t S - \hat{F}_t X^o + W_t(S_{t+1}^o, D_{t+1}^o)$ .
9 end

```

---

fixed and iterate on the policy functions, using previous iterations' policies as starting points. This ensures that the continuation value incorporates each period's returns and costs only once until the final period, while allowing for any numerical errors in initial policy solves to be corrected. This type of "policy iteration" typically takes 1-2 iterations to converge to within  $1e-3$ . Backwards induction on the value function in the final period treats that period's costs and returns as steady state values. This is why we change the notation for the fleet value function for algorithm 3, indexing by time to indicate that the launch cost and satellite per-period return are changing in each period.

Once we have a sequence of policy functions for each period's economic parameters, we generate time paths by picking a starting condition  $(S_0, D_0)$ , computing the launch rate  $X_0$  by thin-plate spline interpolation of the policy function, using the launch rate to compute the next-period state variables, and repeating the process until the terminal period. Figure 3 of the main text shows the simulated open access and optimal paths of launches, satellites, debris, and collision risk over the in-sample period, 2005-2015, as well as projections from 2016-2040.

### 1.3.5 Accounting for launch availability constraints

The maximum number of satellites which can be launched in a year are limited by a variety of factors, including weather, availability of rockets, and availability of launch sites. To prevent the model from violating the limited availability of launches, we estimate the launch constraint from the observed historical data and then project it forward. In each historical period, we calculate the maximum number of satellites which can be launched as the cumulative maximum of launch attempts (successes+failures). From the historical calculation, we project the launch constraint forward with a linear time trend and an intercept. Table S6 shows the estimated coefficients, and Figure S2 shows the projected launch constraint time path.

Launch constraint model parameters	Intercept	Time trend
Parameter values	30.13	12.5
Standard errors	16.43	2.65

Table S6: Parameter values from linear model of launch constraint. All values are rounded to two decimal places. We estimate these coefficients using OLS on the historical launch constraint.

In the historical period, we use the adjusted launch costs (described in section 1.3.2) When the zero-profit or optimal number of launches exceeds the launch constraint in the projection period, we impose the constraint. If the constraint binds for both the planner and open access firms, then the estimated optimal OUF will be zero. If the constraint binds on open access firms but not on the planner, the optimal OUF will be lower than if the constraint had not bound open access firms. If we impose the constraint when in reality it would not bind, we will underestimate the optimal OUF.

## 1.4 Sensitivity analyses of physical equation calibration

To study the sensitivity of our conclusions to uncertainty in our physical parameter estimates, we conduct a sensitivity analysis of the model simulations given different physical parameter values. We use a residual bootstrap procedure to obtain sets of alternative parameter values.

First, we estimate equations 12 and 13 as described above. Then, we sample from the distribution of residuals to generate “bootstrap worlds”. We add these residuals to the estimated models to generate bootstrap world outcome variables. Finally, we re-estimate equations 12 and 13 using the bootstrap world outcomes to generate alternate sets of physical parameter estimates, and simulate the open-access and optimal models under a random sample of those estimates. Algorithm 4 describes our procedure precisely.

---

### Algorithm 4: Bootstrap procedure

---

- 1 Estimate equations 12 and 13 to obtain base parameter estimates,  $\Phi$ , and residuals,  $\{(\varepsilon_{Lt}, \varepsilon_{Dt})\}_t$ .
  - 2 Draw from  $\{(\varepsilon_{Lt}, \varepsilon_{Dt})\}_t$  with replacement to generate  $b = 1, \dots, B$  bootstrap resamples  $\{(\varepsilon_{Lt}^b, \varepsilon_{Dt}^b)\}_t$ .
  - 3 Generate  $B$  bootstrap time series,  $L^b(S_t, D_t)$  and  $D_{t+1}^b$ , using equations 12 and 13 with base estimates  $\Phi$  and resampled residuals  $\{(\varepsilon_{Lt}^b, \varepsilon_{Dt}^b)\}_t$ .
  - 4 Re-estimate equations 12 and 13 using  $L^b(S_t, D_t)$  and  $D_{t+1}^b$  as dependent variables to obtain bootstrap parameter estimates  $\{\Phi^b\}_1^B$ .
- 

One issue to note is that, because we estimate equation 12 with a constrained procedure and the coefficients are near one of the constraint boundaries, the asymptotic properties of this procedure are difficult to obtain (Ketz, 2018). Since our goal is not asymptotic analysis of standard errors but rather to generate alternate parameter sets in a principled way for counterfactual simulations, we select the main model estimates as the mean of the bootstrap world parameters. This ensures that our sensitivity analysis selects parameters around the main model estimates. Ultimately this is inconsequential for the outcomes of interest — collision risk under open access and optimal management, and the resulting OUF — since the outcomes are endogenous variables which satisfy economic conditions irrespective of the specific physical parameter values. The physical parameter values affect the specific paths of launches, satellites, and debris, but only such that the collision risk continues to satisfy the economic conditions.

## 1.5 Projecting the optimal OUF path

With the calibrated parameter values, we turn to projecting the optimal OUF path. We split this process into two stages. In the first stage, we compute the time paths of the satellite stock, debris stock, and launch rate, given the open access and fleet planner models of orbit use. These describe the projected evolution of the orbital aggregates. In the second stage, we use the computed time paths with the estimated collision probability function and launch cost path to calculate the optimal OUF. The OUF is derived from the same open access and fleet planner models. It describes the amount which a satellite owner would have to be charged every year, beginning from the projection horizon’s initial conditions, in order to align their incentives with the fleet planner’s.<sup>16</sup> We show the in-sample fit of our open access projections to establish that our approach can approximate the observed history, and then use predictions of space economy revenues and costs from Jonas et al. (2017) to project out the open access and optimal launch rates given those predictions.

We calculate the time path of an optimal OUF from equation 18:

---

<sup>16</sup>This can also be thought of as “How much of the profits from orbit use currently reflect resource rents which should not have been dissipated?”



$$\tau_t = (L(S_{t+1}^o, D_{t+1}^o) - L(S_{t+1}^*, D_{t+1}^*))F_{t+1}, \quad (18)$$

where  $S_{t+1}^o$  and  $D_{t+1}^o$  are satellite and debris stocks in  $t + 1$  under open access management, and  $S_{t+1}^*$  and  $D_{t+1}^*$  are satellite and debris stocks in  $t + 1$  under optimal management. The optimal OUF is positive whenever the planner would maintain a lower collision probability than firms under open access would. The planner, in turn, will maintain a lower collision probability if the lifetime returns from another satellite in orbit are less than that satellite's expected future damages to other satellites in the fleet. By charging open access firms the marginal external cost of their satellite as an OUF, their incentives are aligned with those of the planner despite the institutional differences. With their incentives aligned, their decisions to launch or not are shifted to optimize the total intertemporal economic value from orbit use rather than their own individual profit.

Formally, equation 18 can be derived by comparing the open access equilibrium condition (equation 7) to the fleet planner's optimality condition for launching (the first-order condition of system of equations 8). These conditions can be written to express the expected loss in satellite value (collision probability multiplied by replacement cost) in terms of economic and, in the case of the optimality condition, physical parameters. Those economic parameters include terms for the current excess return on a satellite in addition to the capital gain or loss from changes in the cost of a replacement satellite. By subtracting the optimal expected collision probability from the open-access expected collision probability, we recover the additional physical and economic term the social planner accounts for — the marginal external cost of a satellite. The marginal external cost is the optimal OUF value to levy on each satellite.

## 1.6 Projecting the effects of active debris removal under open access

Finally, we consider the effects of active debris removal technologies on the NPV losses due to open access, shown in Figure 4 of the main text. We assume that debris removal is available for zero cost, and that 50% of all debris is removed from orbit each period once the technology is available. For example, in a scenario where removal begins in 2030, we assume that 50% of all debris in orbit is removed every year beginning in 2030. We assume debris is removed before it collides with any other orbiting object, and that implementing the removal technology does not require any additional satellites.

These assumptions help us bound a “best case” removal scenario. In reality, removal will cost more than \$0 per unit removed, will require some additional satellites on orbit to implement, and will not be guaranteed to be successful in all cases. Since debris removal in LEO is not commercially available yet, we experiment with different removal start years between 2021–2034.

The laws of motion with debris removal are

$$S_{t+1} = S_t(1 - L(S_t, D_t(1 - R_t)))\mu + X_t \quad (19)$$

$$D_{t+1} = D_t(1 - R_t)(1 - \delta) + G(S_t, D_t(1 - R_t)) + \gamma A_t + mX_t, \quad (20)$$

where  $R_t = 0.5$  if removal technologies are available and 0 otherwise.

The open access equilibrium condition is unchanged from the condition without debris removal (equation 7). The planner's optimality condition with freely-provided debris removal is similar to the condition without debris removal (equation 9), but with  $(1 - R_t)$  terms scaling the debris variable and all derivatives with respect to debris:

$$\begin{aligned} W_{D,t}(S_t, D_t(1 - R_t)) = & -S_t L_D(S_t, D_t(1 - R_t))(1 - R_t)F_t + \\ & \beta[1 - \delta + G_D(S_t, D_t(1 - R_t))(1 - R_t) + \\ & mS_t L_D(S_t, D_t(1 - R_t))(1 - R_t)]W_{D,t+1}(S_{t+1}, D_{t+1}(1 - R_{t+1})), \end{aligned} \quad (21)$$

where

$$W_{D,t}(S_t, D_t(1-R_t)) = \left[ \frac{F_{t-1}}{\beta} - \pi_t - (1 - L(S_t, D_t(1-R_t)) - S_t L_S(S_t, D_t(1-R_t))) F_t - \left( \frac{G_S(S_t, D_t(1-R_t)) - m(1 - L(S_t, D_t(1-R_t)) - S_t L_S(S_t, D_t(1-R_t)))}{1 - \delta + G_D(S_t, D_t(1-R_t))(1-R_t) + mL_D(S_t, D_t(1-R_t))(1-R_t)} \right) L_D(S_t, D_t(1-R_t))(1-R_t) S_t F_t \right] \cdot \left[ \frac{G_S(S_t, D_t(1-R_t)) - m(1 - L(S_t, D_t(1-R_t)) - S_t L_S(S_t, D_t(1-R_t)))}{1 - \delta + G_D(S_t, D_t(1-R_t))(1-R_t) + mL_D(S_t, D_t(1-R_t))(1-R_t)} + m \right]^{-1}. \quad (22)$$

## 1.7 Derivation of the optimal launch rate

In this section we derive the equations characterizing the planner's launch rule, equations 9 and 10. Period  $t$  values are shown with no subscript, and period  $t+1$  values are marked with a  $'$ , e.g.  $S_t \equiv S, S_{t+1} \equiv S'$ . The fleet planner's problem is

$$W(S, D) = \max_{X \geq 0} \{ \pi S - FX + \beta W(S', D') \} \quad (23)$$

$$\text{s.t. } S' = S(1 - L(S, D)) + X \quad (24)$$

$$D' = D(1 - \delta) + G(S, D) + \gamma A + mX. \quad (25)$$

The fleet planner's launch plan will satisfy

$$X^* : \beta [W_S(S', D') + mW_D(S', D')] = F, \quad (26)$$

that is, the planner will launch until the marginal value to the fleet of a new satellite plus the marginal value to the fleet of its launch debris is equal to the launch cost.

Assuming an optimal policy function  $X^* = H(S, D)$  exists and applying the envelope condition, we have the following expressions for the fleet's marginal value of another satellite and another piece of debris:

$$W_S(S, D) = \pi + \beta [W_S(S', D')(1 - L(S, D) - SL_S(S, D)) + W_D(S', D')G_S(S, D)] \quad (27)$$

$$W_D(S, D) = \beta [W_D(S', D')(1 - \delta + G_D(S, D)) + W_S(S', D')(-SL_D(S, D))] \quad (28)$$

Rewriting equation 26, we have

$$W_S(S', D') = \left[ \frac{F}{\beta} - mW_D(S', D') \right] \quad (29)$$

Plugging equation 29 into equations 27 and 28,

$$W_S(S, D) = \pi + F(1 - L(S, D) - SL_S(S, D)) - \beta W_D(S', D')[m(1 - L(S, D) - SL_S(S, D)) - G_S(S, D)] \quad (30)$$

$$W_D(S, D) = (-SL_D(S, D))F + \beta W_D(S', D')[1 - \delta + G_D(S, D) - m(-SL_D(S, D))] \quad (31)$$

Define the following quantities:

$$\alpha_1(S, D) = \pi + (1 - L(S, D) - SL_S(S, D))F$$

$$\alpha_2(S, D) = -SL_D(S, D)F$$

$$\Gamma_1(S, D) = G_S(S, D) - m(1 - L(S, D) - SL_S(S, D))$$

$$\Gamma_2(S, D) = 1 - \delta + G_D(S, D) + mSL_D(S, D).$$

These allow us to rewrite equations 30 and 31 as

$$W_S(S, D) = \alpha_1(S, D) + \beta \Gamma_1(S, D)W_D(S', D') \quad (32)$$

$$W_D(S, D) = \alpha_2(S, D) + \beta \Gamma_2(S, D)W_D(S', D'). \quad (33)$$

As long as  $\delta < 1$ ,  $\Gamma_2(S, D) \neq 0 \forall (S, D)$ , allowing us to rewrite equation 33 as

$$W_D(S', D') = \frac{W_D(S, D) - \alpha_2(S, D)}{\beta \Gamma_2(S, D)}. \quad (34)$$

Plugging equation 34 into equation 32, we get

$$\begin{aligned} W_S(S, D) &= \alpha_1(S, D) + \beta \Gamma_1(S, D) \frac{W_D(S, D) - \alpha_2(S, D)}{\beta \Gamma_2(S, D)} \\ &= \alpha_1(S, D) + \frac{\Gamma_1(S, D)}{\Gamma_2(S, D)} (W_D(S, D) - \alpha_2(S, D)) \\ \implies W_S(S, D) &= \alpha_1(S, D) - \frac{\Gamma_1(S, D)}{\Gamma_2(S, D)} \alpha_2(S, D) + \frac{\Gamma_1(S, D)}{\Gamma_2(S, D)} W_D(S, D) \end{aligned} \quad (35)$$

Iterating equation 29 one period backwards and plugging it into equation 32, we get

$$\begin{aligned} \frac{F}{\beta} - m W_D(S, D) &= \alpha_1(S, D) - \frac{\Gamma_1(S, D)}{\Gamma_2(S, D)} \alpha_2(S, D) + \frac{\Gamma_1(S, D)}{\Gamma_2(S, D)} W_D(S, D) \\ \implies W_D(S, D) &= \left[ \frac{F}{\beta} - \alpha_1(S, D) - \frac{\Gamma_1(S, D)}{\Gamma_2(S, D)} \alpha_2(S, D) \right] \left[ \frac{\Gamma_1(S, D)}{\Gamma_2(S, D)} + m \right]^{-1}. \end{aligned} \quad (36)$$

Substituting in the forms for  $\alpha_1(S, D)$ ,  $\alpha_2(S, D)$ ,  $\Gamma_1(S, D)$ , and  $\Gamma_2(S, D)$ , equation 36 yields

$$\begin{aligned} W_D(S, D) &= \left[ \frac{F}{\beta} - \pi - (1 - L(S, D) - S L_S(S, D)) F - \frac{G_S(S, D) - m(1 - L(S, D) - S L_S(S, D))}{1 - \delta + G_D(S, D) + m L_D(S, D)} L_D(S, D) S F \right] \\ &\quad \left[ \frac{G_S(S, D) - m(1 - L(S, D) - S L_S(S, D))}{1 - \delta + G_D(S, D) + m L_D(S, D)} + m \right]^{-1}. \end{aligned} \quad (37)$$

Combining equation 37 with equation 36 iterated one period forwards, we obtain

$$L(S', D') = 1 + r'_s - (1 + r) \frac{F}{F'} - \xi(S', D'),$$

where

$$\begin{aligned} \xi(S', D') &= S' L_S(S', D') F' + \frac{\pi - r F - L(S, D) F - S L_S(S, D) F}{\beta(1 - \delta + G_D(S', D') + m L_D(S', D') S')} \\ &\quad - \frac{\beta G_S(S', D') + m(1 - L(S', D') - S' L_S(S', D'))}{\beta(1 - \delta + G_D(S', D') + m L_D(S', D') S')} L_D(S', D') S' F'. \end{aligned}$$

## 1.8 Discussion

### 1.8.1 Interpreting the optimal OUF and long-run industry value paths

Figure 2 in the main text shows the NPV gains from beginning optimal management in different years, and the permanent orbit use value losses in 2040 from delaying optimal management. The permanent orbit use value is the discounted value of the satellite fleet over the long-run, accounting for losses and replacements.

As mentioned in the main text, the discontinuous jumps in NPV in Figure 2 reflect the immediate effect of reducing launch activity while the satellite and debris stocks are suboptimally high. The change in launch activity increases the NPV by reducing the cost outflows each year. Since open access launching results in excess satellites and debris, the benefits of optimal management continue to accrue gradually as the satellite and debris stocks draw down. Figure 3 in the main text shows these launch rate dynamics for a 2006 optimal management start.

The fact that the optimal OUF, shown in Figure 2 of the main text, rises over time is an indication of the restored value of orbit use due to optimal management (orbital rents). Intuitively, firms fully dissipate orbital rents in an open

access equilibrium, as those rents act as an incentive to enter. To counter the growth in the incentive to enter due to restored orbital rents, the OUF must also rise over time. A firm which enters the commons early on (say, 2020) will pay a lower initial fee than a firm which enters later on (say, 2030) because the early entrant is paying to use a more congested environment.

Though our modeling procedure abstracts from many economic and physical complications, our optimal OUF and long-run industry value estimates are likely robust to these limitations and on the correct order of magnitude with the correct qualitative features. On the OUF side, given that the projected optimal launch path beginning from the projected 2020 state of LEO involves cessation of launch activity, the estimated OUF in 2020 needs only to be large enough to deter launches, particularly those likely to generate large and suboptimal increases in collision risk. While satellite operators in LEO are heterogeneous and represent diverse interests, we estimate an optimal averaged-across-LEO-use-cases per-satellite-year fee beginning at approximately \$13,500 USD and growing over time. Letizia et al. (2017) identify that the majority of collision risk increases in the near future will be driven by satellite constellation operators. For an entity planning to launch a constellation of 600 satellites, our OUF amounts to an additional yearly expenditure beginning at \$8.1 million USD — likely enough to prompt serious reconsideration of the size and nature of the constellation. On the industry value side, the global satellite sector currently produces approximately \$0.15 trillion USD in revenues, and is anticipated to grow to \$2 trillion USD in revenues and costs by 2040 (Jonas et al., 2017). A decrease in collision risk which leads to a 10% increase in the per-year economic value of the satellite sector would immediately add 10% to the sector’s NPV. Our model projects that collision risk in 2040 will decrease by roughly 33% (compared to BAU) under an optimal management plan beginning in 2020. This reduction in collision risk would cut collision-related replacement costs, increase the expected lifespan of satellites in orbit, and reduce collision-related disruptions in the stream of satellite-related economic returns. Given this, long-run industry value in 2040 on the order of single-digit trillions of USD seems plausible.

### **1.8.2 Open access and active debris removal**

The introduction of debris removal makes both open access firms and the planner launch additional satellites. However, the planner launches considerably fewer additional satellites than open access firms. The immediate decrease in debris when removal becomes available induces new launches under open access until the collision risk is once again equated with the excess return on a satellite. Figure 4 of the main text shows the effects of debris removal beginning in 2029 on satellite and debris accumulation under open access and a range of optimal management paths. The removal-induced additional launching leads to a higher steady-state satellite stock and a lower steady-state debris stock under open access and optimal management. The amount of debris removed in our “50% removal” scenario is substantially larger under open access than the optimal plan as there is more debris in orbit under open access.

Though debris removal allows open access to sustain more satellites in orbit, over time collision risk returns to the equilibrium level. The costs of additional launches erodes some of the NPV gains due to reduced risk. Figure 4e in the main text shows the percentage change in open-access NPV loss due to the introduction of debris removal in 2029, as a fraction of the potential gain from implementing the optimal plan in 2020. Figure 4f in the main text summarizes the minimum, mean, and maximum changes in long-run industry value losses due to open access with removal beginning in each year in 2021–2034. On average, debris removal beginning in the 2021–2034 window reduces open access NPV losses by about 1.65% relative to a counterfactual world without removal. When optimal management begins ahead of debris removal, the NPV losses tend to be reduced (negative changes). When optimal management begins after debris removal, the NPV losses tend to be increased (positive changes). The increase in NPV losses in the latter scenario is driven by the costs of additional launching under open access, both immediately following the initial debris reduction and to sustain the larger satellite population. Due to open access, firms will take advantage of lower collision risk due to debris removal by launching satellites until there are no more profits from launching new satellites. When the planner is in charge of launching before debris removal begins, they do not waste resources by launching to dissipate the gains from debris removal.

### **1.8.3 Unmodeled physical and economic factors**

In addition to the limitations imposed by modeling spatially and temporally heterogeneous physical and economic processes at an aggregated level, there are three main analytical limitations pertaining to unobservables in the past

and present and unknowables in the future: launch market frictions, constellations (coordinated systems of satellites intended to serve a common purpose), and satellite placement. These limitations may make our OUF estimates lower bounds on the true values required to induce optimal orbit use, as we describe below.

Our conclusions about the suboptimality of open access to orbit and the necessity of a globally-coordinated OUF (or policies equivalent to one) are robust to these limitations. The fundamental problem creating the need for policies equivalent to a globally-coordinated OUF is the lack of legally-enforceable property rights over orbits.<sup>17</sup> The lack of property rights prevents satellite owners from internalizing the costs they impose on others through collision risk and debris creation. The same issue manifests in other common-resource settings, such as fisheries (Gordon, 1954).

Our economic model is founded on the assumption that all agents who want to launch satellites are able to do so with no frictions. In practice, there are factors other than orbital property rights and willingness-to-pay which limit agents' access to orbit, such as limited availability of launch windows and rockets. These factors constrain humanity's ability to launch satellites. To ensure that our simulations do not violate this launch constraint in observed years, we calculate the launch constraint in each observed period as the cumulative maximum number of launches observed so far. The shadow value of the launch constraint is recovered in the economic parameter calibration process, but the individual factors are not identifiable from the data. We then fit a linear time trend to the observed launch constraint, and project it into the future. To the extent that the launch constraint will be relaxed faster than a linear trend would predict, our estimates are economically conservative, i.e. we assume fewer launches than may occur, which biases our estimated OUF downward.

Our economic model is also founded on the simplifying assumption of "one satellite per firm". In practice, there are a number of firms which own constellations or fleets of satellites. However, unless a single firm owned all satellites in orbit, orbit users would not internalize the full scope of the externality they impose on others. To the extent that the observed data reflects agents internalizing those externalities due to ownership of multiple satellites, our economic parameter estimates would entangle those factors with the estimated launch constraint shadow value. Our projections of single-satellite-owning firms' responses to increases in satellite profitability would therefore be attenuated toward zero, making our projections environmentally conservative, i.e. closer to an environmental "worst-case" analysis. However, the same assumption also increases the optimal OUF we estimate.

Lastly, our model abstracts away entirely from the question of satellite placement. That is, two orbital objects within a given volume shell can be placed in orbits such that at one extreme they are guaranteed to collide, or at the other extreme they will never collide. Our projections are based on collision rate estimates which are calculated using historical placement patterns. Thus, our projections assume that the systematic factors which resulted in current object placements will continue into the future. While technology and constellation ownership are likely to lead to improvements in placement patterns our collision risk projections would be biased for both the open access and optimal launch paths. However, the magnitude of the gap between open access and optimal collision risk may actually be understated by this issue. To the extent that economic agents have the placement margin available to them it induces another externality, similar in spirit but different in detail to the orbit use externality we describe in this article, wherein firms do not account for the full magnitude of orbital-use efficiency losses due to their placement. A fleet planner who coordinated all satellites in orbit would account for such placement-related externalities. By taking advantage of any efficiencies in placement, the planner would be able to reduce collision rates below what open access satellite owners would have an incentive to consider. Thus, while the inclusion of a placement margin may reduce levels of collision risk, the differences in collision risk between open access and optimal use may increase, which would make our OUF estimates a lower bound on average.<sup>18</sup>

#### **1.8.4 Consequences of measurement error in satellite and debris counts**

Limitations of sensor technology suggest that the debris counts we use are lower-bound estimates. To the extent that this biases the collision probability and debris counts downward, it will bias the estimated decay rate, collision

---

<sup>17</sup>The geostationary belt is the exception to this statement. In general, however, there is no globally-coordinated procedure for allocating orbital paths, or even a globally-agreed-upon definition of an orbital path property right.

<sup>18</sup>While some regimes in a spatially-differentiated orbit model may have lower OUF values than the ones we calculate here, others may have higher OUF values.



probability parameters, fragmentation parameters, and launch debris weakly downwards. Since downward bias in the physical parameters makes collisions and missile tests appear to cause less congestion than they actually do, the open access and optimal launch rates will be inflated.

Downward bias in the collision probability data will bias the economic parameter estimates weakly downwards as well. This will to some degree offset the inflation in the launch rate caused by the physical parameter underestimation, though the exact extent of the offset is not clear.

In general, measurement error in the collision probability data also causes the nonnegativity constraint on the collision probability parameters ( $\alpha_{SS}$  and  $\alpha_{SD}$ ) to bind in some bootstrap replications. This causes issues of the type described in Ketz (2018) in obtaining asymptotic standard errors.

### **1.8.5 Consequences of collision probability model misspecification**

We assume that the collision probability model has constant parameters. Changes in patterns of satellite placement, construction, and ownership structures lead to changes over time in the physical primitives reflected in  $\alpha_{SS}$  and  $\alpha_{SD}$ . The “net” convexity or concavity of the time path of the primitives will determine whether the constant approximations over or understate the true time-varying parameters in any period on average. A convex time path — low values initially and high values later on — will be overestimated on average, while a concave time path — high values initially, with slow increases over time — will be underestimated on average.

The misspecification causes two problems with simulation and inference. First, underestimation will inflate launch rate projections and overestimation will deflate them. However, because the misspecification affects both open access and optimal launch rates in the same way, the simulated optimal OUF will not be affected. Second, underestimation may cause the nonnegativity constraint on the collision probability parameters to bind in some bootstrap replications, causing the same types of asymptotic issues as measurement error.

### **1.8.6 Consequences of measurement error in returns and costs**

We take the returns and costs of satellite ownership from the data used in Wienzierl (2018), which aggregate revenues from all commercial satellites in orbit. By including more than just LEO satellites, the direct returns and costs data overstate the returns to LEO paths. The economic parameter estimates therefore reflect a “LEO share” coefficient on the revenue data between 0 and 1. The LEO share coefficient attenuates the estimates of  $a_{L1}$ ,  $a_{L2}$ , and  $a_{L3}$ .




Figure S1: Calibrated physical model fits to historical data. The black dots show historical data points (1957–2014) for debris counts, active satellite counts, and estimated collision rate. The blue dashed lines show the fits of the calibrated models.




Figure S2: Projected launch constraint model. To represent the various factors constraining the number of rocket launches possible each year – e.g. limited rocket availability, weather-induced launch cancellations, limited launch slots – we model the aggregate launch constraint as a linear time trend based on observed launch behavior from 2006–2015. We use this constraint in our open access and optimal management model simulations to avoid implausibly large launch rates in any given year. Panel a shows the observed and projected launch constraint. The solid black line from 2006–2015 indicates the observed cumulative maximum number of rocket launches in any given year, the dashed blue line from 2016–2050 shows the projected maximum number of rocket launches possible in each year, and the vertical dashed black line marks the end of the in-sample period (2006–2015). Panel b shows the estimated intercept and time trend with standard errors.




Figure S3: Sensitivity of physical model calibration to parameter uncertainty. Panels a-b show a sensitivity analysis of the model calibrations for the collision rate function and debris law of motion. The black lines show the underlying collision rate and debris count data from ESA, 1957–2014. The thin darker blue lines show the fit of the models calibrated to the black lines (the same model fits shown in figure 3 of the main text). The light blue lines show model fits from residual bootstrap-resampled data, indicating the effects of uncertainty in the model calibration. The same procedure was used to generate Fig. 2d. We describe the residual bootstrap resampling procedure in more detail in section 1.4 of the Supporting Information.




Figure S4: Sensitivity of open-access paths to physical parameter uncertainty. Panels a-d show a sensitivity analysis of open-access model projections beginning in 2020 to uncertainty in the physical parameters. The thick black lines show the projection from the main model estimated on the full 1957–2014 time series of object counts and collision rate estimates. The gray lines show projections using alternate sets of parameter estimates generated from 50 draws of the residual bootstrap-resampling procedure used in main text figure 2d.




Figure S5: Sensitivity of optimal paths to physical parameter uncertainty. Panels a-d show a sensitivity analysis of optimal model projections beginning in 2020 to uncertainty in the physical parameters. The thick black lines show the projection from the main model, where the physical parameters were estimated on the 1957–2014 calibration time series constructed from ESA and UCS count data. The gray lines show projections using alternate sets of parameter estimates generated from 50 draws of the residual bootstrap-resampling procedure used in main text figure 2d.




Figure S6: Sensitivity of optimal OUF to physical parameter uncertainty. The thick black line shows the projection from the main model, where the physical parameters were estimated on the 1957–2014 calibration time series constructed from ESA and UCS count data. The gray lines show projections using alternate sets of parameter estimates, generated from 50 draws of the residual bootstrap-resampling procedure used in main text figure 2d.

## References

- Bradley, Andrew M. and Lawrence M. Wein. 2009. "Space debris: Assessing risk and responsibility." *Advances in Space Research* 43:1372–1390.
- Combined Space Operations Center. 2018. "Space-Track.org Satellite Catalog." From Space-Track.org at <https://www.space-track.org/>.
- European Space Agency. 2018. "DISCOS database." From ESA at <https://discosweb.esoc.esa.int/web/guest/home>.
- Gordon, H. Scott. 1954. "The Economic Theory of a Common-Property Resource: The Fishery." *Journal of Political Economy* 62.
- IADC. 2007. "IADC Space Debris Mitigation Guidelines." Tech. Rep. IADC-02-01.
- Jonas, Adam, Arminas Sinkevicius, Simon Flannery, Benjamin Swinburne, Patrick Wellington, Terence Tsui, Rajeev Lalwani, James E Faucette, Brian Nowak, Ravi Shanker, Kai Pan, and Eva Zlotnicka. 2017. "Space: Investment Implications of the Final Frontier." Tech. rep., Morgan Stanley. URL [https://fa.morganstanley.com/griffithwheelwrightgroup/mediahandler/media/106686/Space\\_%20Investment%20Implications%20of%20the%20Final%20Frontier.pdf](https://fa.morganstanley.com/griffithwheelwrightgroup/mediahandler/media/106686/Space_%20Investment%20Implications%20of%20the%20Final%20Frontier.pdf).
- Kelly, David L and Charles D Kolstad. 1999. "Integrated assessment models for climate change control." *International yearbook of environmental and resource economics* 2000:171–197.
- Ketz, Philipp. 2018. "Subvector inference when the true parameter vector may be near or at the boundary." *Journal of Econometrics* 207:285–306.
- Letizia, F., S. Lemmens, and H. Krag. 2018. "Application of a debris index for global evaluation of mitigation strategies." 69th International Astronautical Congress.
- Letizia, Francesca, Camilla Colombo, Hugh Lewis, and Holger Krag. 2017. "Extending the ECOB space debris index with fragmentation risk estimation."
- Nordhaus, William. 2013. "Chapter 16 - Integrated Economic and Climate Modeling." In *Handbook of Computable General Equilibrium Modeling SET, Vols. 1A and 1B, Handbook of Computable General Equilibrium Modeling*, vol. 1, edited by Peter B. Dixon and Dale W. Jorgenson. Elsevier, 1069 – 1131. URL <http://www.sciencedirect.com/science/article/pii/B978044459568300016X>.
- Rao, Akhil. 2019. *The Economics of Orbit Use: Theory, Policy, and Measurement*. Ph.D. thesis, University of Colorado at Boulder.
- Union of Concerned Scientists. 2018. "UCS Satellite Database." From UCS at <https://www.ucsusa.org/nuclear-weapons/space-weapons/satellite-database>. Accessed April 2019.
- Wienzierl, Matthew. 2018. "Space, the Final Economic Frontier." *Journal of Economic Perspectives* 32:173–192.
- Wilkerson, Jordan, Benjamin Leibowicz, Delavane Diaz, and John Weyant. 2015. "Comparison of Integrated Assessment Models: Carbon Price Impacts on U.S. Energy." *Energy Policy* 76:18–31.