Ketoprofen for Daniel Okoro

Order ord_2f154cce · AWAITING_REVIEW · prescribed by Dr. Erik Lindqvist

Rx: 10%, 3×/day, 30 days, topical | 55.0 y · 90.0 kg | Silver Star Medicare Part D

Conditions

- Renal impairment
- Hypertension
- Osteoarthritis
- Atrial fibrillation

Allergies / avoids

- Soy-derived

Current medicines

- Lisinopril
- Warfarin

Proposed formulation

Strength	Make	Beyond-use date
10%	60 g	35 days

Topical ointment in **Hydrophilic ointment base** · Apply three times daily

Label: Apply a thin layer to the affected area three times daily. Apply to right knee. Store at room temperature.

Beyond-use date source: No product-specific study: USP <795> default beyond-use date applies

Bases ruled out

- Pluronic lecithin organogel (PLO): Contains Soy lecithin; patient allergic to Soy-derived.

Insurance coverage

Silver Star Medicare Part D · member SSD-430-88 · synthetic demo plan

Recommended	Patient copay	Prior authorization
Diclofenac transdermal gel	$9	Not needed

Covered options

- Diclofenac transdermal gel: $9 copay (ARTHRITIS PAIN RELIEVER (H E B), Arthritis Pain Reliever (Family Dollar (FAMILY WELLNESS)))
- compounded Ketoprofen 10% topical ointment: $47 copay, prior auth

Denied or unsuitable

None.

Message to prescriber

```
Dr. Erik Lindqvist, about Daniel Okoro (member ID SSD-430-88, Silver Star Medicare Part
D):
You prescribed Ketoprofen 10%, 3x/day.
Silver Star Medicare Part D does not cover a suitable form of Ketoprofen, but it covers
Diclofenac transdermal gel at $9 copay, no prior auth. Rx diclofenac 1% gel.
Would you consider switching? If yes, please send a new prescription for Diclofenac
transdermal gel.
I checked these products against the patient's allergies and conditions.
```

Compounding procedure

Ingredients

- Ketoprofen USP 6 g
- Hydrophilic ointment base 54 g

Equipment

- Analytical balance
- Ointment slab and spatulas
- Ointment mill / electronic mortar
- Jar or tube

Steps

1. Calculate: 10% × 60 g = 6 g Ketoprofen.
2. Weigh 6 g Ketoprofen USP powder and 54 g Hydrophilic ointment base.
3. Levigate the drug powder with a few drops of a compatible levigating agent until smooth.
4. Incorporate the base in portions (geometric dilution) on an ointment slab until uniform.
5. Pass through an ointment mill (or mix in an electronic mortar) for a smooth, grit-free product.
6. Fill into a 60 g jar or tube, label and store: room temperature.

Quality checks

- Smooth, no grit or particles
- Final weight checked
- Second person verifies calculation

Label text

- Apply a thin layer to the affected area three times daily. Apply to right knee. Store at room temperature.
- Discard 35 days after the compounding date (beyond-use date).
- Compounded medication. Keep out of reach of children.

Demo only. A licensed pharmacist must verify every step.

Safety findings

WARN · interaction — Ketoprofen + Warfarin: Increased bleeding risk.
`Ketoprofen is a NSAID; Warfarin is a Vitamin K antagonist anticoagulant`

WARN · condition — Renal impairment: NSAIDs can worsen kidney function (lower with topical use).
`Ketoprofen → NSAID`


WARN · coverage — Best covered option on Silver Star Medicare Part D: Diclofenac transdermal gel at $9 copay, no prior auth.

INFO · interaction — Ketoprofen + Lisinopril: NSAIDs can blunt the antihypertensive effect and stress kidney function.
`Ketoprofen is a NSAID; Lisinopril is a ACE inhibitor`

INFO · regulatory — No FDA-labeled Ketoprofen product found in an acceptable form (openFDA snapshot). Compounding is justified on availability.

Evidence graph

What the agents looked at in FalkorDB.



Patient Drug DrugClass Base Excluded base Excipient ExcipientGroup Condition DosageForm

Agent trace

Intake · *validate_prescription*
Ketoprofen 10%, 3x/day for 30 days (topical) for Daniel Okoro.

History · *load_profile*
Daniel Okoro, 55.0 y, 90.0 kg. Conditions: Renal impairment, Hypertension, Osteoarthritis, Atrial fibrillation. Allergies: Soy-derived. Current meds: Lisinopril, Warfarin.

Formulation · *screen_bases*
2 candidate base(s); 1 excluded. Pluronic lecithin organogel (PLO): Contains Soy lecithin; patient allergic to Soy-derived.

Formulation · *propose_formulation*
Ketoprofen 10% topical ointment in Hydrophilic ointment base (USP <795> default BUD, BUD 35 days). Apply three times daily. Make 60 g.

Safety · *safety_review*
0 blocking, 2 warning, 2 info finding(s).

Coverage · *check_coverage*
Silver Star Medicare Part D: 2 covered option(s), 0 denied or unsuitable. Recommend: Diclofenac transdermal gel.

Generated by Compounder (FalkorDB + agents). Demo only, not for clinical use. All patients and insurance plans are synthetic.