

Systematic Review: Recent Development of Magnetic Nanoparticles for DNA Extraction

Stanley Evander Emeltan Tjoa ^{1), 4)}, Mudasir Mudasir²⁾, Edi Suharyadi ³⁾, Budi Setiadi Daryono ^{*1)}

**Corresponding author*

*ORCHID IDs: <https://orcid.org/0000-0002-0703-2123>

¹⁾ Department of Biology, Universitas Gadjah Mada, Yogyakarta, Indonesia

²⁾ Department of Chemistry, Universitas Gadjah Mada, Yogyakarta, Indonesia

³⁾ Department of Physics, Universitas Gadjah Mada, Yogyakarta, Indonesia

⁴⁾ Konimex Diagnostic Center, PT Konimex, Solo, Indonesia

**email: bs_daryono@mail.ugm.ac.id*

Abstract

DNA is a research object for various molecular fields of study and DNA extraction is a basic procedure in a molecular biology laboratories. DNA extraction methods are evolving, from several steps using organic solvents to recent technology using magnetic nanoparticles as adsorbent for its purification step. Using magnetic nanoparticles for DNA extraction is magnetizing because it is simple without heavy machinery such as centrifugation and organic solvents. The application of this method is very wide and diverse. This systematic review summarises current developmental trends for some aspects of this DNA extraction system from nanoparticle synthesis method, extraction protocol, and different types of samples used. The main purpose of this review is to give a big picture of magnetic nanoparticle utilization in DNA extraction for the researcher to start their own exploration regarding this safe, simple, and rapid technique.

Keywords: *Magnetic nanoparticle, Nucleic acid, DNA isolation, Product development*

1. Introduction

Deoxyribonucleic Acid or DNA is a strand of polymer that carries genetic material stored in cells. DNA is an object of research for various technological fields, such as molecular diagnostic technologies such as PCR [1], bioinformatics [2] and genetic engineering [3]. The technique of obtaining DNA is an important factor in the development of this technology.

DNA extraction is a method for taking DNA from cells and separating it from other impurities. The principle of DNA extraction is cell lysis and DNA purification. Cell lysis is the stage where the cell is broken down so that the DNA comes out. In this stage, not only DNA comes out but also other molecules mix with the target DNA. Therefore, there is the next stage, DNA purification.

DNA purification techniques are very varied, starting from extraction with organic solvents such as phenol: chloroform extraction, DNA will be extracted in the water phase and impurities in the organic phase [4]; chromatography method using a column, DNA will be

bound to the column and impurities will flush out and then the DNA will be eluted with an appropriate solvent [5]; to modern methods that use adsorption using particles that can bind DNA [6]. One method of DNA adsorption is the use of magnetic nanoparticles.

Magnetic nanoparticles are nano-sized materials that have unique properties that can react to magnetic fields. This property is a typical property of the transition metal group, one of which is Fe or iron. Iron was chosen as one of the materials for making magnetic nanoparticles because of its non-toxic and biocompatible properties [7]. Apart from that, it is also very abundant in nature, in the form of iron oxide [8]. Magnetite or Fe_3O_4 is a form of iron oxide that exists in nature. Magnetite has the strongest magnetic properties among all other iron oxides [9].

The use and development of magnetic nanoparticles for DNA extraction is quite common. This review will collect information on the fabrication of magnetic nanoparticles that have been carried out and the DNA extraction protocol. We hope that this review can be used to broaden the horizon of knowledge regarding the use of magnetic nanoparticles in the DNA extraction process and can become a basis for further development and innovation. This review can also be a starting point for readers to start their in-house DNA extraction using magnetic nanoparticles.

2. Method

The search method uses Boolean operators and quotation marks to determine specific keywords. The keywords used “magnetic nanoparticles” AND “DNA extraction” OR “DNA isolation” were entered into the entry fields in four databases including JSTOR (<https://www.jstor.org/>); DOAJ (<https://doaj.org/>); Mendeley (<https://www.mendeley.com/search/>); and Scopus (<https://www.scopus.com/search/form.uri?display=basic#basic>). The article selection process is displayed in the PRISMA flow diagram to make it easier to observe the selection process at each stage so that the articles that will be reviewed are selected. Articles included in the inclusion are articles that contain how to make nanoparticles and their use in DNA extraction.

Search results with the keywords "magnetic nanoparticle" AND "DNA extraction" OR "DNA isolation" on JSTOR, DOAJ, Mendeley, and Scopus, display 1, 190, 10, and 166 articles respectively. Next, initial screening was carried out by reading the titles and abstracts of articles that discussed DNA isolation or extraction using magnetic nanoparticles, and then the articles were collected. Apart from screening the title and abstract, this stage also eliminates articles whose full text is not available. Obtained from the JSTOR, DOAJ, Mendeley, and Scopus databases respectively, leaving 1, 1, 8, and 62 articles.


Then, the article skimming process is carried out to see the contents of the article at a glance. Articles that are eliminated from this stage are articles that do not explain how to make magnets and/or use magnets for DNA extraction.

In the DOAJ database, all articles were eliminated from the previous stage because DNA extraction was carried out using commercial magnetic nanoparticles, for the same reason one article was also eliminated from the Mendeley database and 12 articles from the Scopus database. There were two and 12 articles from Mendeley and Scopus that were eliminated because the studies were conducted using commercial DNA or DNA extracted using other

methods (not using magnetic nanoparticles) to conduct DNA interactions study. In the Scopus database, four articles were eliminated because they discussed the development of automated DNA extraction machines with magnetic nanoparticles, and four articles were eliminated because they did not include the magnetic nanoparticle synthesis process.

Next, the selection is carried out using the Mendeley reference manager to mark duplicate articles. There are eight additional articles from manual searches using Google Scholar regarding the use of magnetic nanoparticles for DNA extraction. In total, 37 articles will be reviewed.


3. PRISMA Statement




4. Magnetic Nanoparticle Synthesis


Co-precipitation is the most common method used in making magnetic nanoparticles for DNA extraction. This is shown in Figure 1 (A). Co-precipitation is a very simple method. This method can be carried out at room temperature and does not produce toxic byproducts. The advantage of this method is that, with fixed parameters, the resulting product will always have similar quality. In addition, this method is very easy to scale up [10,11].

In producing magnetic nanoparticles, the precursors used are transition metals. For iron oxide, the precursor is Fe. In Figure 1 (B) it is shown that the precursors that are widely used are FeCl_2 : FeCl_3 and FeSO_4 : FeCl_3 . It does not rule out the possibility of using other Fe salts such as $\text{Fe}(\text{acac})_3$ [12]; $\text{Fe}(\text{NO}_3)_2$ [13]; $(\text{NH}_4)_2\text{Fe}(\text{SO}_4)_2$; and $\text{NH}_4\text{Fe}(\text{SO}_4)_2$ [14,15]. The commonly used Fe^{2+} and Fe^{3+} ratio is 1:2, this is related to the amount of impurities or byproducts produced [16].





C.



D.

Figure 1. Recapitulation of Methods for Making Magnetic Nanoparticles Used for DNA Extraction. (A) Kind of Methods (B) Kind of Precursor (C) Kind of Base (D) Fe²⁺ and Fe³⁺ ratio used in co-precipitation

Iron oxide synthesis requires alkaline conditions, optimal at pH > 8 [17]. The base used is quite uniform. As seen in Figure 1 (C) there are 3 types of bases used, NaOH; NH₄OH, and KOH. According to research by Mascolo et al [18], the effect of the difference in base is the size of the nanoparticles produced. This could be related to the nature of the base. Strong base or weak base.

For their use in DNA extraction, magnetic nanoparticles can be modified or not. This can be seen in Figure 2. The most frequently used coating agent is tetraorthosilicate (TEOS). Magnetic nanoparticles are coated with the aim of protecting them from agglomeration [19,20]. Apart from that, it also protects it from oxidation [21]. However, the main goal is for magnetic nanoparticles to specifically bind DNA and separate it from impurities during DNA isolation [22].




Figure 2. Coating Agent Used in Magnetic Nanoparticle Fabrication for DNA Extraction

5. DNA Extraction

DNA extraction is divided into two steps. The first step, lysis, is the stage of breaking down cells. This step aims to make DNA come out of the cells or matrix that confines it. The second step is purification. This stage aims to purify DNA from impurities in the lysate mixture from the first step. In general, the DNA purification step with magnetic nanoparticles is divided into three. Binding, washing, and elution. These three steps are important because they relate to the interactions of DNA, impurities, and magnetic nanoparticles. ingredients that are often or commonly used in DNA extraction using magnetic nanoparticles can be seen in Figure 3.

Binding buffer is expected to provide conditions for DNA and magnetic nanoparticle materials to bind. From Figure 3 (A), it can be seen that generally, a combination of PEG and NaCl is used for the binding stage. This is possible because of the nature of PEG and NaCl which will cause DNA to precipitate and be adsorbed on the magnetic material. Other ingredients used will also have more or less the same purpose, such as isopropanol. Other impurity particles remain in the supernatant to be removed.

Washing buffer is used to dissolve impurities that are bound to the magnetic nanoparticles. The washing buffer that is often used is 70% ethanol, as seen in Figure 3 (B). Ethanol is used for impurity elimination, the same as the pellet washing stage in conventional DNA extraction methods [23]. The majority of other ingredients in the washing buffer are alcohol with some additional salt. This is unique because the function of 70% ethanol is to dissolve impurity salts and minimize DNA solubility, but here, these salts are instead added to the washing buffer.

Elution buffer is used to dissolve DNA that is still bound to the magnetic nanoparticles. The solution that is often used as an elution buffer is Tris EDTA. This can be seen in Figure 3 (C). The materials used as elution buffers are materials that can dissolve DNA. The nature of DNA is polar, so the solutions for elution are selected that are polar. The addition of salt, such as NaCl or potassium acetate is aimed to maintain the stability of the eluted DNA.




Figure 3. Regular Reagent used in DNA Extraction Using Magnetic Nanoparticle. (A) Binding Buffer; (B) Washing Buffer; (C) Elution Buffer.

The types of samples used in extraction vary greatly. As seen in Table 1. sample types range from human specimens, animals, plants, bacteria, environmental samples, and food. Further development can be carried out for other types of species or other human specimens,

such as urine and sputum. These various sample types can be integrated into the next process of DNA research, this is discussed further in the next point, about DNA quality.

Table 1. Various Kind of Sample That Can Be Processed by Magnetic Nanoparticle DNA Extraction

Samples	Source	Samples	Source
Human specimen		Animal	
Saliva	[24,25]	Animal tissue	[49]
Blood	[12,25-31]	animal blood	[45]
Serum	[28]	Cattle	[43]
FFPE tissue	[32]	Plant	
Tissue	[31]	Cottonseed	[48]
Bacteria		Clover	[49]
Plasmid [13,26,33-39,56]		Daisy	[49]
<i>E. coli</i> [13,28,40-43,50]		Ryegrass	[49]
<i>L. monocytogenes</i>	[44]	Leaf (<i>A. littoralis</i> , Rice)	[43]
<i>S. enteriditis</i>	[44]	Petal <i>R. hybrida</i>	[43]
<i>Arthrospira platensis</i>	[45]	Environment	
<i>P. viridiflava</i>	[43]	Soil	[51]
<i>S. aureus</i>	[43]	Pond water	[52]
<i>B. subtilis</i>	[43]	Food	
<i>X. campestris</i>	[43]	Milk [44,48,52]	
<i>O. anthropi</i>	[46]	Juice	[52]
Fungi		Any other Sample	
<i>R. oryzae</i>	[14]	Virus	[28,53]
<i>Aspergillus niger</i>	[47]	Gel electrophoresis	[54]
Yeast	[28]	Cell culture	[49]
Fungus	[48]	HeLa cell	[55]
		Synthetic	[49]

6. DNA Quality

The quality of DNA extraction can be seen from the yield, purity, and downstream process. The yield value shows the amount of DNA that can be extracted. Purity is indicated by the absorbance value at a wavelength of 260 nm divided by absorbance at 280 nm. The third quality can be seen from the exploitation of DNA in advanced processes. The results of DNA extraction using magnetic nanoparticles can be seen in Table 2. The results of DNA extraction are quite varied. From a range of 0.03 µg to over 100 µg. This may be affected by many factors. From the type of sample, availability of the sample, and the total of cells extracted.

The purity of the extracted DNA can be seen in Figure 4. The purity ratio is categorized by a group, plant, animal, human, plasmid, bacteria, and any other sample. The purity range varies from 0.8 – 2.6 (in bacterial samples). DNA extract is said to be pure if it has a ratio in the range of 1.8. A value below 1.8 indicates that the isolate contains a lot of impurities in the form of protein. If the ratio gives a value above 2, it indicates that the DNA isolated may still

be contaminated by RNA [57]. This data may be further analyzed using meta-analysis to determine what factors influence the purity of the DNA extract.



Table 2. DNA Yield for Various Sample

Study	Yield (μg)	Study	Yield (μg)
Animal sample		Bacteria sample	
Rat liver tissue [54]	1.8 ± 0	<i>X. campestris</i> [43]	0.58 ± 0.01
Cattle [43]	1.8 ± 0.14	<i>S. aureus</i> [43]	0.6 ± 0.01
Rat brain tissue [31]	2 ± 0	<i>P. viridiflava</i> [43]	0.62 ± 0.13
Blood [23]	3 ± 0.91	<i>E. coli</i> [43]	0.71 ± 0.11
Plant samples		<i>B. subtilis</i> [43]	0.73 ± 0.11
Plant sample [49]	0.03 ± 0.01	<i>E. coli</i> [28]	1.75 ± 0.08
Leaf of Rice (Nemat cultivars) [43]	2.37 ± 0.1	<i>A. platensis</i> [45]	2.38 ± 0.039
<i>Aeluropus littoralis</i> [43]	2.44 ± 0.12	<i>E. coli</i> [42]	3.27 ± 1.23
Petals of Rosa hybrid [43]	2.5 ± 0.16	Bacteria sample [49]	6.17 ± 1.75
Human specimen		Bacteria sample [41]	7.29 ± 2.98
Blood [31]	1.2 ± 0	<i>E. coli</i> [50]	19 ± 1.41
Blood [28]	1.31 ± 0.07	Any other samples	
Blood [26]	1.57 ± 0.37	Environment/ Soil [51]	0.49 ± 0
Human specimen [43]	1.95 ± 0.19	Virus [28]	0.69 ± 0.07
Saliva [24]	8.2 ± 2.71	Yeast [28]	0.95 ± 0.01
Blood [27]	62.93 ± 4.46	Gel Electrophoresis [54]	10 ± 0
Plasmid		Environment/ water [52]	16.5 ± 2.12
Plasmid [35]	6.9 ± 0	FFPE tissue [32]	17.5 ± 10.61
Plasmid [56]	8.57 ± 0.13	Food [52]	19 ± 1.41
Plasmid [33]	10.9 ± 0.49	Cell/ HeLa cell [55]	116.63 ± 45.36
Plasmid [38]	11.69 ± 6.67		
Plasmid [49]	14.92 ± 5.76		
Plasmid [26]	76.1 ± 32.2		
Plasmid [13]	91.1 ± 4.08		


Table 3. Downstream Process Applying Extracted DNA Using Magnetic Nanoparticle

Downstream Process	Source
Electrophoresis	[24,28,29,33,31,38,39,42,45,47,50,51,54]
PCR (conventional, Multiplex, qPCR)	[12,14,24-28,30-32,34-35,38,40,44,47,49,50-53,55]
Restriction Analysis	[14,33,37,50,56]
Reverse Transcriptase	[43]
Sequencing	[25,46,49]
Cloning and Expression	[37,49]
Bisulphite conversion	[49]
Mass Spectrophotometry	[49]


The third quality is seen from the availability of the extracted DNA to be used in advanced stages of research. This list is shown in Table 3. Processes that exploit this DNA are very diverse, ranging from electrophoresis to mass spectrophotometry.



A260/280 for Plant Sample



A260/280 for Bacteria Sample






Figure 4. Forest Plot for A260/280 for Different Kind of Sample, Such as Human, Animal, Plant, Bacteria, Plasmid, and Any Other Sample

7. Conclusion

The adoption of magnetic nanoparticles as adsorbents for DNA extraction has become very common for various types of samples. From human, animal, plant, and bacterial samples. For diverse purposes such as health, environment, and food. The DNA results from the extraction can also be widely used for numerous molecular biology laboratory processes such as PCR,

cloning expression, and sequencing. Hopefully, this review can give a big picture as a starting point to carry out product development or further applications using magnetic nanoparticles for DNA extraction.

References

- [1] Debnath, M; Prasad, GBKS; & Bisen, PS. 2010. *Molecular Diagnostics: Promises and Possibilities*. New York: Springer
- [2] Claverie, J & Notredame, C. 2007. *Bioinformatics For Dummies*. 2nd Edition. New Jersey: Wiley Publishing
- [3] Hudge, Russ. 2009. *Genetic Engineering: Manipulating the Mechanism of Life*. New York: Facts On File
- [4] Butler, JM. 2012. *Advanced Topics in Forensic DNA Typing: Methodology*. Maryland: Elsevier
- [5] Budelier, K & Schorr, J. 1998. *Purification of DNA by Anion-Exchange Chromatography*. Current Protocols in Molecular Biology. 42(1).
- [6] Gupta, N. 2019. *DNA Extraction and Polymerase Chain Reaction*. Journal of Cytology. 36(2):116-117
- [7] Kandpal, ND; Sah, N; Loshali, R; Joshi, R; & Prasad, J. 2014. *Co-precipitation method of synthesis and characterization of iron oxide nanoparticles*. Journal of Scientific & Industrial Research. 73: 87-90
- [8] Voloshina, E. 2018. *Hematite, Its Stable Surface Terminations and Their Reactivity To Water*. Encyclopedia of Interfacial Chemistry :115-121
- [9] Teja, AS & Koh, PY. 2009. *Synthesis, Properties, and Application of Magnetic Iron Oxide Nanoparticles*. Progress in Crystal Growth and Characterization of Materials. 55: 22-45
- [10] Lu, A; Salabas, EL; & Schuth, F. 2007. *Magnetic Nanoparticles: Synthesis, Protection, Functionalization, and Application*. Angewandte Chemie. 46: 1222-1244
- [11] Ahn, T; Kim, JH; Yang, H; Lee, JW; & Kim, J. 2011. *Formation Pathways of Magnetite Nanoparticles by Coprecipitation Method*. Journal of Physical Chemical C. 116: 6069-6076
- [12] Min, JH; Woo, M; Yoon, HY; Jang, JW; Wu, JH; Lim, C; & Kim, YK. 2014. *Isolation of DNA using magnetic nanoparticles coated with dimercaptosuccinic acid*. Analytical Biochemistry. 447: 114–118
- [13] Gerzsenyi, TB; Ilosvai, AM; Szilágyi, G; Szóri, M; Várad, C; Viskolcz, B; Vanyorek, L; & Szóri-Dorogházi, E. 2023. *A Simplified and Efficient Method for Production of Manganese Ferrite Magnetic Nanoparticles and Their Application in DNA Isolation*. International Journal of Molecular Science. 24: 2156
- [14] Alghuthaymi, M. 2020. *Magnetic-silica nanoshell for extraction of fungal genomic DNA from Rhizopus oryzae*. Biointerface Research in Applied Chemistry. 10 (2): 4972 - 4976
- [15] Danthanarayana, AN; Manatunga, DC; De Silva, RM; Chandrasekharan, NV; & De Silva, MN. 2018. *Magnetofection and isolation of DNA using polyethyleneimine functionalized magnetic iron oxide nanoparticles*. Royal Society Publishing. 5: 181369
- [16] Jiang, W; Lai, K; Hu, H; Zeng, X; Lan, F; Liu, K; Wu, Y; & Gu, Z. 2010. *The effect of [Fe³⁺]/[Fe²⁺] molar ratio and iron salts concentration on the properties of superparamagnetic iron oxide nanoparticles in the water/ethanol/toluene system*. Journal of Nanoparticle Research. 13: 5135–5145
- [17] Ramadan, W; Kareem, M; Hannoyer, B; & Saha, S. 2011. Effect of pH on the structural and

- magnetic properties of magnetite nanoparticles synthesised by co-precipitation. *Advanced Materials Research*. 324: 129-132
- [18] Mascolo, MC; Pei, Y; & Ring, TA. 2013. *Room Temperature Co-Precipitation Synthesis of Magnetite Nanoparticles in a Large pH Window with Different Bases*. *Materials*. 6: 5549-5567
- [19] Favela-Camacho, SE; Samaniego-Benitez, EJ; Godinez-Garcia, A; Aviles-Arellano, L; & Perez-Robles, JF. 2009. *How to decrease the agglomeration of magnetite nanoparticles and increase their stability using surface properties*. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*. 574:29-35
- [20] de Mendonca, ESDT; de Faria, ACB; Dias, SCL; Aragon, FFH; Mantila, JC; Coaquira, JAH; & Dias, JA. 2019. *Effects of silica coating on the magnetic properties of magnetite nanoparticles*. *Surfaces and Interfaces*: 34-43
- [21] Dzeranov, A; Bondarenko, L; Pankratov, D; Dzhardimalieva, G; Jorobekova, S; Saman, D; & Kydralieva, K. 2015. *Impact of Silica-Modification and Oxidation on the Crystal Structure of Magnetite Nanoparticles*. *Magnetochemistry*. 2023. 9. 18
- [22] Shi, B; Shin, YK; Hassanali, AA; Singer, SJ. *DNA Binding to the Silica Surface*. *The Journal of Physical Chemistry*. 119. 34
- [23] Sambrook, J & Russell, DW. 2001. *Molecular Cloning A Laboratory Manual*. Cold Spring Harbor laboratory Press: New York
- [24] Bhati, A; Varghese, A; Rajan, G; Sridhar, V; Mohan, Y; Pradeep, S; Babu, S; Kaikkolante, N; Sarma, M; Arun, S; Sekar, AP; Iype, T; Santhosh, S; & Ramchand, CN. 2020. *An effective method for saliva stabilization and magnetic nanoparticles based DNA extraction for genomic applications*. *Analytical Biochemistry*. 624: 114182
- [25] Vangala, RK; Sridevi, G; & Hugar, V. 2010. *Instrument-free, automation and multi-platform ready unmodified iron Oxide based DNA isolation system*. *Journal Biochemical Technic*. 3 (4): 384-388
- [26] Rahnama, H; Sattarzadeh, A; Kazemi, F; Ahmadi, N; Sanjarian, F; & Zand, Z. 2016. *Comparative study of three magnetic nano-particles (FeSO₄, FeSO₄/ SiO₂, FeSO₄/SiO₂/TiO₂) in plasmid DNA extraction*. *Analytical Biochemistry*. 513: 68e76
- [27] Zhang, M; Li, L; Li, B; Tian, N; Yang, M; Zhang, H; You, C; & Zhang, J. 2019. *Adsorption of DNA by using polydopamine modified magnetic nanoparticles based on solid-phase extraction*. *Analytical Biochemistry*. 579: 9–17
- [28] Ma, C; Li, C; Wang, F; Ma, N; Li, X; Li, Z; Deng, Y; Wang, Z; Xi, Z; Tang, Y; & He, N. 2013. *Magnetic Nanoparticles-Based Extraction and Verification of Nucleic Acids from Different Sources*. *Journal of Biomedical Nanotechnology*. 9: 703–709
- [29] Ali, TH; Mandal, AM; Heidelberg, T; Hussena, RSD; & Goh, EW. 2020. *Ionic magnetic core–shell nanoparticles for DNA Extraction*. *RSC Advances*. 10: 38818
- [30] Stolyar, SV; Komogortsev, SV; Gorbenko, AS; Knyazev, YV; Yaroslavtsev, RN; Neznakhin, DS; Tyumentseva, AV; Bayukov, OA; & Iskhakov, RS. 2022. *Maghemite Nanoparticles for DNA Extraction: Performance and Blocking Temperature*. *Journal of Superconductivity and Novel Magnetism*. 35: 1929–1936
- [31] Saiyed, ZM; Ramchand, CN; & Telang, SD. 2008. *Isolation of genomic DNA using magnetic nanoparticles as a solid-phase support*. *Journal of Physics: Condensed Matter*. 20: 204153
- [32] Hieu, NM; Nam, NH; Huyen, NT; Anh, NTV; Nghia, PTN; Khoa, NB; Toan, NL; & Luong, NH. 2017. *Synthesis of SiO₂-Coated Fe₃O₄ Nanoparticles Using Ultrasound and Its Application in DNA Extraction from Formalin-Fixed, Paraffin-Embedded Human Cancer Tissues*. *Journal of Electronic Materials*. 46 (6)
- [33] Shan, Z; Li, C; Zhang, X; Oakes, KD; Servos, MR; Wu, Q; Chen, H; Wang, X; Huang, Q; Zhou,

- Y; & Yang, W . 2011. *Temperature-dependent selective purification of plasmid DNA using magnetic nanoparticles in an RNase-free process*. Analytical Biochemistry. 412: 117–119
- [34] Saraji, M; Yousefia, S; & Talebi, M. 2017. *Plasmid DNA purification by zirconia magnetic nanocomposite*. Analytical Biochemistry. 539: 33–38
- [35] da Silva, RJ; Maciela, BG; Medina-Llamas, JC; Chávez-Guajardoc, AE; Alcaraz-Espinoza, JJ; & de Melo, CP. 2019. *Extraction of plasmid DNA by use of a magnetic maghemite-polyaniline nanocomposite*. Analytical Biochemistry. 575: 27–35
- [36] Fan, Q; Guan, Y; Zhang, Z; Xu, G; Yang, Y; & Guo, C. 2019. *A new method of synthesis well-dispersion and dense Fe₃O₄@SiO₂ magnetic nanoparticles for DNA extraction*. Chemical Physics Letters. 715: 7–13
- [37] He, X; Huo, H; Wang, K; Tan, W; Gong, P; & Ge, J. 2007. *Plasmid DNA isolation using amino-silica coated magnetic nanoparticles (ASMNPs)*. Talanta. 73: 764–769
- [38] Sheng, HY; Yuan, JL; & Liu, CX. 2019. *Fabrication of superparamagnetic Fe₃O₄ particles in a dynamic reaction kettle and their application in DNA extraction*. Journal of Beijing University of Chemical Technology (Natural Science). 46 (6)
- [39] Gai, L; Li, Z; Hou, Y; Jiang, H; Han, X; & Ma, W. 2010. *Preparation of core-shell Fe₃O₄/SiO₂ microspheres as adsorbents for purification of DNA*. Journal of Physic. D: Applied Physic. 43: 44500
- [40] Firoozeh, F; Neshan, A; Khaledi, A; Zibaei, M, Amiri, A; Sobhani, A; Badmasti, F; & Nikbin, VS. 2023. *Evaluation of the effect of magnetic nanoparticles on extraction of genomic DNA of Escherichia coli*. Polymer Bulletin. 80: 3153–3163
- [41] Pansini, M; Dell’Agli, G; Marocco, A; Netti, PA; Battista, E; Lettera, V; Vergara, P; Allia, P; Bonelli, B; Tiberto, P; Barrera, G; Alberto, G; Martra, G; Arletti, R; & Esposito, S. 2013. *Preparation and Characterization of Magnetic and Porous Metal-Ceramic Nanocomposites from a Zeolite Precursor and Their Application for DNA Separation*. Journal of Biomedical Nanotechnology. 13: 337–348
- [42] Torres-Rodríguez, J; Soto, G; Medina, JL; Portillo-López, A; Hernández-López, EL; Viveros, EV; Galindo, JTE; Tiznado, H; Flores, D; & Muñoz-Muñoz, F. 2019. *Cobalt-zinc ferrite and magnetite SiO₂ nanocomposite powder for magnetic extraction of DNA*. Journal of Sol-Gel Science and Technology. 91:33–43
- [43] Ghahari, S; Ghahari, S; & Nematzadeh, GA. 2018. *Magnetic nano fluids for isolation of genomic DNA and total RNA from various prokaryote and eukaryote cells*. Journal of Chromatography B 1102–1103: 125–134
- [44] Bai, Y; Song, M; Cui, Y; Shi, C; Wang, D; Paoli, GC; Shi, X. 2013. *A rapid method for the detection of foodborne pathogens by extraction of a trace amount of DNA from raw milk based on amino-modified silica-coated magnetic nanoparticles and polymerase chain reaction*. Analytica Chimica Acta. 787: 93–101
- [45] Nguyen, LT; Le, NT; Ta, HKT; & Nguyen, KD. 2022. *Isolation of DNA from Arthrospira platensis and whole blood using magnetic nanoparticles (Fe₃O₄@OA and Fe₃O₄@OA@SiO₂)*. Journal of Analytical Science and Technology. 13(1):28
- [46] Tiwari, AP; Rohiwal, SS; Suryavanshi, MV; Ghosh, SJ; & Pawar, SH. 2016. *Detection of the genomic DNA of pathogenic α -proteobacterium Ochrobactrum anthropi via magnetic DNA enrichment using pH responsive BSA@Fe₃O₄ nanoparticles prior to in-situ PCR and electrophoretic separation*. Microchimica Acta. 183(2):675–681
- [47] Hou, Y; Han, X; Chen, J; Li, Z; Chen, X; & Gai, L. 2013. *Isolation of PCR-ready genomic DNA from Aspergillus niger cells with Fe₃O₄/SiO₂ microspheres*. Separation and Purification Technology. 116: 101–106
- [48] Bai, Y; Roncancio, D; Suoa, Y; Shao, Y; Zhang, D; & Zhou, C. 2019. *A method based on amino-modified magnetic nanoparticles to extract DNA for PCR-based analysis*. Colloids and Surfaces B: Biointerfaces. 179: 87–93

- [49] Oberacker, P; Stepper, P; Bond, DM; Höhn, S; Focken, J; Meyer, V; Schelle, L; Sugrue, VJ; Jeunen, G; Moser, T; Hore, SR; von Meyenn, F; Hipp, K; Hore, TA; & Jurkowski, TP. 2019. *Bio-On-Magnetic-Beads (BOMB): Open platform for high-throughput nucleic acid extraction and manipulation*. PLOS Biology
- [50] Bandyopadhyay, A & Chatterjee, S. 2011. *Rapid isolation of genomic DNA from E. coli XL1 Blue strain approaching bare magnetic nanoparticles*. Current Science. 101 (2): 210-214
- [51] Paul, T; Basu, S; & Sarkar, K. 2014. *SPION-mediated soil DNA extraction and comparative analysis with conventional and commercial kit-based protocol*. 3 Biotech. 4: 669-677
- [52] Basu, S; Chatterjee, S; Bandyopadhyay, A; & Sarkar, K. 2012. *Potential application of superparamagnetic nanoparticles for extraction of bacterial genomic DNA from contaminated food and environmental samples*. Journal of The Science of Food and Agriculture
- [53] Quy,DV; Hieu, QM; Tra, PT; Nam, NH; Hai, NH; Son, NT; Nghia, PT; Anh, NTV; Hong, TT; & Luong, NH. 2013. *Synthesis of Silica-Coated Magnetic Nanoparticles and Application in the Detection of Pathogenic Viruses*. Journal of Nanomaterials. 2013: 603940
- [54] Saiyed, ZM; Parasramka, M; Telang, SD; & Ramchand, CN. 2007. *Extraction of DNA from agarose gel using magnetic nanoparticles (magnetite or Fe₃O₄)*. Analytical Biochemistry 363: 288–290
- [55] Zhou, Z; Kadam, US; & Irudayaraj, J. 2013. *One-stop genomic DNA extraction by salicylic acid-coated magnetic nanoparticles*. Analytical Biochemistry. 442: 249–252
- [56] Shan, Z; Wu, Q; Wang, X; Zhou, Z; Oakes, K; Zhang, X; Huang, Q; & Yang, W. 2010. *Bacteria capture, lysate clearance, and plasmid DNA extraction using pH-sensitive multifunctional magnetic nanoparticles*. Analytical Biochemistry. 398: 120–122
- [57] Lucena-Aguillar, G; Sánchez-López, AM; Barberán-Aceituno, C; Carrillo-Ávila, JA; López-Guerrero, JA; & Aguilar-Quesada, R. 2016. *DNA Source Selection for Downstream Applications Based on DNA Quality Indicators Analysis*. Biopreservation and Biobanking. 14(4): 264-270