- 1. (18pts) (a) If an m by n matrix Q has orthonormal columns is the matrix Q necessarily invertible? Give a reason or a counterexample.
  - (b) What is the nullspace of Q (and WHY)?
  - (c) What is the projection matrix onto the column space of Q? Avoid inverses where possible.
- 2. (30pts) We look for the line y = C + Dt closest to 3 points (t, y) = (0, -1) and (1, 2) and (2, -1).
  - (a) If the line went through those points (it doesn't), what three equations would be solved?
  - (b) Find the best C and D by the least squares method.
  - (c) Explain the result you get for C and D: How is the vector b = (-1, 2, -1) related to the plane you are projecting onto?
  - (d) What is the length of the error vector e (= distance to plane =  $||b A\bar{x}||$ ).
- 3. (22pts) The problem is to find the determinants of

- (a) Find  $\det A$  and give a reason.
- (b) Find  $\det B$  using elimination.
- (c) Find  $\det C$  for any value of x. For this you could use Property 1 of the determinant.
- 4. (30pts) (a) Decide if A is singular or invertible.  $A = \begin{bmatrix} 1 & 2 & -3 \\ 2 & 1 & -3 \\ 2 & 2 & -4 \end{bmatrix}.$ 
  - (b) Find an orthonormal basis for its column space (if such a basis exists).
  - (c) Why does  $P = A(A^TA)^{-1}A^T$  not give the projection matrix onto the column space of A?

Find that projection matrix somehow.