

Exercises on orthogonal matrices and Gram-Schmidt

Problem 17.1: (4.4 #10.b *Introduction to Linear Algebra*: Strang)

Orthonormal vectors are automatically linearly independent.

Matrix Proof: Show that $Q\mathbf{x} = \mathbf{0}$ implies $\mathbf{x} = \mathbf{0}$. Since Q may be rectangular, you can use Q^T but not Q^{-1} .

Solution: By definition, Q is a matrix whose columns are orthonormal, and so we know that $Q^T Q = I$ (where Q may be rectangular). Then:

$$Q\mathbf{x} = \mathbf{0} \implies Q^T Q\mathbf{x} = Q^T \mathbf{0} \implies I\mathbf{x} = \mathbf{0} \implies \mathbf{x} = \mathbf{0}.$$

Thus the nullspace of Q is the zero vector, and so the columns of Q are linearly independent. There are no non-zero linear combinations of the columns that equal the zero vector. Thus, orthonormal vectors are automatically linearly independent.

Problem 17.2: (4.4 #18) Given the vectors \mathbf{a} , \mathbf{b} and \mathbf{c} listed below, use the Gram-Schmidt process to find orthogonal vectors \mathbf{A} , \mathbf{B} , and \mathbf{C} that span the same space.

$$\mathbf{a} = (1, -1, 0, 0), \mathbf{b} = (0, 1, -1, 0), \mathbf{c} = (0, 0, 1, -1).$$

Show that $\{\mathbf{A}, \mathbf{B}, \mathbf{C}\}$ and $\{\mathbf{a}, \mathbf{b}, \mathbf{c}\}$ are bases for the space of vectors perpendicular to $\mathbf{d} = (1, 1, 1, 1)$.

Solution: We apply Gram-Schmidt to $\mathbf{a}, \mathbf{b}, \mathbf{c}$. First, we set

$$\mathbf{A} = \mathbf{a} = (1, -1, 0, 0).$$

Next we find \mathbf{B} :

$$\mathbf{B} = \mathbf{b} - \frac{\mathbf{A}^T \mathbf{b}}{\mathbf{A}^T \mathbf{A}} \mathbf{A} = (0, 1, -1, 0) + \frac{1}{2}(1, -1, 0, 0) = \left(\frac{1}{2}, \frac{1}{2}, -1, 0\right).$$

And then we find \mathbf{C} :

$$\mathbf{C} = \mathbf{c} - \frac{\mathbf{A}^T \mathbf{c}}{\mathbf{A}^T \mathbf{A}} \mathbf{A} - \frac{\mathbf{B}^T \mathbf{c}}{\mathbf{B}^T \mathbf{B}} \mathbf{B} = (0, 0, 1, -1) + \frac{2}{3} \left(\frac{1}{2}, \frac{1}{2}, -1, 0\right) = \left(\frac{1}{3}, \frac{1}{3}, \frac{1}{3}, -1\right).$$

We know from the first problem that the elements of the set $\{\mathbf{A}, \mathbf{B}, \mathbf{C}\}$ are linearly independent, and each vector is orthogonal to $(1,1,1,1)$. The space of vectors perpendicular to \mathbf{d} is three dimensional (since the row space of $(1,1,1,1)$ is one-dimensional, and the number of dimensions of the row space added to the number of dimensions of the nullspace add to 4). Therefore $\{\mathbf{A}, \mathbf{B}, \mathbf{C}\}$ forms a basis for the space of vectors perpendicular to \mathbf{d} .

Similarly, $\{\mathbf{a}, \mathbf{b}, \mathbf{c}\}$ is a basis for the space of vectors perpendicular to \mathbf{d} because the vectors are linearly independent, orthogonal to $(1,1,1,1)$, and because there are three of them.

MIT OpenCourseWare
<http://ocw.mit.edu>

18.06SC Linear Algebra
Fall 2011

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.