

These are the slides of the lecture

Pattern Recognition
Winter term 2020/21
Friedrich-Alexander University of Erlangen-Nuremberg.

These slides are released under Creative Commons License Attribution CC BY 4.0.

Please feel free to reuse any of the figures and slides, as long as you keep a reference to the source of these slides at <https://lme.tf.fau.de/teaching/> acknowledging the authors Niemann, Hornegger, Hahn, Steidl, Nöth, Seitz, Rodriguez, Das and Maier.

Erlangen, January 8, 2021
Prof. Dr.-Ing. Andreas Maier

Pattern Recognition (PR)

Prof. Dr.-Ing. Andreas Maier

Pattern Recognition Lab (CS 5), Friedrich-Alexander-Universität Erlangen-Nürnberg

Winter Term 2020/21



Duality in Optimization



The Primal Problem

- Consider the *primal optimization problem*:

$$\begin{array}{ll}\text{minimize} & f_0(\mathbf{x}) \\ \text{subject to} & f_i(\mathbf{x}) \leq 0, \quad i = 1, 2, \dots, m \\ & h_i(\mathbf{x}) = 0, \quad i = 1, 2, \dots, p\end{array}$$

with variable $\mathbf{x} \in \mathbb{R}^n$.

- The function $f_0(\mathbf{x})$ is **not** required to be **convex**.

The Lagrangian

Lagrangian

The *Lagrangian* L of the aforementioned problem is defined as

$$L(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\nu}) = f_0(\mathbf{x}) + \sum_{i=1}^m \lambda_i f_i(\mathbf{x}) + \sum_{i=1}^p \nu_i h_i(\mathbf{x})$$

The Lagrangian

Lagrangian

The *Lagrangian* L of the aforementioned problem is defined as

$$L(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\nu}) = f_0(\mathbf{x}) + \sum_{i=1}^m \lambda_i f_i(\mathbf{x}) + \sum_{i=1}^p \nu_i h_i(\mathbf{x})$$

- λ_i is the Lagrange multipliers associated with the i -th *inequality* constraint $f_i(\mathbf{x}) \leq 0$.

The Lagrangian

Lagrangian

The *Lagrangian* L of the aforementioned problem is defined as

$$L(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\nu}) = f_0(\mathbf{x}) + \sum_{i=1}^m \lambda_i f_i(\mathbf{x}) + \sum_{i=1}^p \nu_i h_i(\mathbf{x})$$

- λ_i is the Lagrange multipliers associated with the i -th **inequality** constraint $f_i(\mathbf{x}) \leq 0$.
- ν_i is the Lagrange multiplier associated with the i -th **equality** constraint $h_i(\mathbf{x}) = 0$.

The Lagrangian

Lagrangian

The *Lagrangian* L of the aforementioned problem is defined as

$$L(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\nu}) = f_0(\mathbf{x}) + \sum_{i=1}^m \lambda_i f_i(\mathbf{x}) + \sum_{i=1}^p \nu_i h_i(\mathbf{x})$$

- λ_i is the Lagrange multipliers associated with the i -th **inequality** constraint $f_i(\mathbf{x}) \leq 0$.
- ν_i is the Lagrange multiplier associated with the i -th **equality** constraint $h_i(\mathbf{x}) = 0$.
- The vectors $\boldsymbol{\lambda}$ and $\boldsymbol{\nu}$ are called *Lagrange multiplier vectors* or simply **dual variables**.

Lagrange Dual Function

Lagrange dual function

The *Lagrange dual function* is defined as the infimum of the Lagrangian over \mathbf{x}

$$g(\boldsymbol{\lambda}, \nu) = \inf_{\mathbf{x}} L(\mathbf{x}, \boldsymbol{\lambda}, \nu)$$

Lagrange Dual Function

Lagrange dual function

The *Lagrange dual function* is defined as the infimum of the Lagrangian over \mathbf{x}

$$\begin{aligned} g(\boldsymbol{\lambda}, \boldsymbol{\nu}) &= \inf_{\mathbf{x}} L(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\nu}) \\ &= \inf_{\mathbf{x}} \left(f_0(\mathbf{x}) + \sum_{i=1}^m \lambda_i f_i(\mathbf{x}) + \sum_{i=1}^p \nu_i h_i(\mathbf{x}) \right) \end{aligned}$$

Langrange Dual Function

Lagrange dual function

The *Lagrange dual function* is defined as the infimum of the Lagrangian over \mathbf{x}

$$\begin{aligned} g(\boldsymbol{\lambda}, \boldsymbol{\nu}) &= \inf_{\mathbf{x}} L(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\nu}) \\ &= \inf_{\mathbf{x}} \left(f_0(\mathbf{x}) + \sum_{i=1}^m \lambda_i f_i(\mathbf{x}) + \sum_{i=1}^p \nu_i h_i(\mathbf{x}) \right) \end{aligned}$$

Note:

- The Lagrange dual function is a **pointwise affine function** in the dual variables.

Langrange Dual Function

Lagrange dual function

The *Lagrange dual function* is defined as the infimum of the Lagrangian over \mathbf{x}

$$\begin{aligned} g(\boldsymbol{\lambda}, \boldsymbol{\nu}) &= \inf_{\mathbf{x}} L(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\nu}) \\ &= \inf_{\mathbf{x}} \left(f_0(\mathbf{x}) + \sum_{i=1}^m \lambda_i f_i(\mathbf{x}) + \sum_{i=1}^p \nu_i h_i(\mathbf{x}) \right) \end{aligned}$$

Note:

- The Lagrange dual function is a **pointwise affine function** in the dual variables.
- The **Lagrange dual function is concave** (even if the original problem is not convex).


Optimal Value and Lower Bound

Lemma


Let p^ be the optimal value of the optimization problem.
For any $\lambda \succeq 0$ and any ν the following bound is valid:*

$$g(\lambda, \nu) \leq p^*$$


Optimal Value and Lower Bound (cont.)




Optimal Value and Lower Bound (cont.)




Optimal Value and Lower Bound (cont.)




Optimal Value and Lower Bound (cont.)




Optimal Value and Lower Bound (cont.)




Optimal Value and Lower Bound (cont.)




Optimal Value and Lower Bound (cont.)




Optimal Value and Lower Bound (cont.)




Optimal Value and Lower Bound (cont.)




Optimal Value and Lower Bound (cont.)




Optimal Value and Lower Bound (cont.)




Optimal Value and Lower Bound (cont.)



Optimal Value and Lower Bound (cont.)



Optimal Value and Lower Bound (cont.)



- Neither $f_0(x)$ nor $f_1(x)$ is convex,
- but the dual function $g(\lambda)$ is concave!

Optimal Value and Lower Bound (cont.)

Let $\tilde{\mathbf{x}}$ be a **feasible point** of the optimization problem.

If $\lambda \succeq 0$, we have due to the **m inequality** and **p equality** constraints:

$$\sum_{i=1}^m \lambda_i f_i(\tilde{\mathbf{x}}) + \sum_{i=1}^p v_i h_i(\tilde{\mathbf{x}}) \leq 0 ,$$

Optimal Value and Lower Bound (cont.)

Let $\tilde{\mathbf{x}}$ be a **feasible point** of the optimization problem.

If $\boldsymbol{\lambda} \succeq 0$, we have due to the **m inequality** and **p equality** constraints:

$$\sum_{i=1}^m \lambda_i f_i(\tilde{\mathbf{x}}) + \sum_{i=1}^p \nu_i h_i(\tilde{\mathbf{x}}) \leq 0 ,$$

Thus we have

$$L(\tilde{\mathbf{x}}, \boldsymbol{\lambda}, \boldsymbol{\nu}) = f_0(\tilde{\mathbf{x}}) + \sum_{i=1}^m \lambda_i f_i(\tilde{\mathbf{x}}) + \sum_{i=1}^p \nu_i h_i(\tilde{\mathbf{x}}) \leq f_0(\tilde{\mathbf{x}}) .$$

Optimal Value and Lower Bound (cont.)

Using the definition of the dual function we get:

$$g(\boldsymbol{\lambda}, \nu) = \inf_{\mathbf{x}} L(\mathbf{x}, \boldsymbol{\lambda}, \nu) \leq L(\tilde{\mathbf{x}}, \boldsymbol{\lambda}, \nu) \leq f_0(\tilde{\mathbf{x}})$$

Optimal Value and Lower Bound (cont.)

Using the definition of the dual function we get:

$$g(\boldsymbol{\lambda}, \nu) = \inf_{\mathbf{x}} L(\mathbf{x}, \boldsymbol{\lambda}, \nu) \leq L(\tilde{\mathbf{x}}, \boldsymbol{\lambda}, \nu) \leq f_0(\tilde{\mathbf{x}})$$

The inequality $g(\boldsymbol{\lambda}, \nu) \leq f_0(\tilde{\mathbf{x}})$ holds for every feasible point $\tilde{\mathbf{x}}$.

Optimal Value and Lower Bound (cont.)

Using the definition of the dual function we get:

$$g(\boldsymbol{\lambda}, \nu) = \inf_{\mathbf{x}} L(\mathbf{x}, \boldsymbol{\lambda}, \nu) \leq L(\tilde{\mathbf{x}}, \boldsymbol{\lambda}, \nu) \leq f_0(\tilde{\mathbf{x}})$$

The inequality $g(\boldsymbol{\lambda}, \nu) \leq f_0(\tilde{\mathbf{x}})$ holds for every feasible point $\tilde{\mathbf{x}}$.

Consequently, the dual function $g(\boldsymbol{\lambda}, \nu)$ is also smaller or equal to the optimal value p^* :

$$g(\boldsymbol{\lambda}, \nu) \leq p^*$$



The Lagrange Dual Problem

Problem: how to find the best lower bound for the primal problem

The Lagrange dual problem is given by the optimization problem:

$$\text{maximize} \quad g(\lambda, \nu)$$

$$\text{subject to} \quad \lambda \succeq 0$$

The Lagrange Dual Problem (cont.)

Optimal duality gap

Let p^* be the optimal value of the primal problem and d^* the optimal value of the Lagrange dual problem.

The Lagrange Dual Problem (cont.)

Optimal duality gap

Let p^* be the optimal value of the primal problem and d^* the optimal value of the Lagrange dual problem.

- The difference $p^* - d^*$ is the *optimal duality gap*.

The Lagrange Dual Problem (cont.)

Optimal duality gap

Let p^* be the optimal value of the primal problem and d^* the optimal value of the Lagrange dual problem.

- The difference $p^* - d^*$ is the *optimal duality gap*.
- If $p^* = d^*$, the duality gap is zero.
In this case we talk about *strong duality*.

The Lagrange Dual Problem (cont.)

Optimal duality gap

Let p^* be the optimal value of the primal problem and d^* the optimal value of the Lagrange dual problem.

- The difference $p^* - d^*$ is the *optimal duality gap*.
- If $p^* = d^*$, the duality gap is zero.
In this case we talk about *strong duality*.
- If $p^* > d^*$, we have *weak duality*.

Slater's Condition

Theorem

Given a *convex* primal optimization problem:

$$\begin{aligned} &\text{minimize} && f_0(\mathbf{x}) \\ &\text{subject to} && f_i(\mathbf{x}) \leq 0, \quad i = 1, 2, \dots, m \\ & && \mathbf{Ax} = \mathbf{b} \end{aligned}$$

with f_0, f_1, \dots, f_m being convex.

If there exists an $\mathbf{x} \in \text{relint} \left\{ \mathcal{D} = \bigcap_{i=1}^m \text{dom}(f_i) \right\}$ with

$$\begin{aligned} &f_i(\mathbf{x}) < 0, \quad i = 1, \dots, m \\ &\mathbf{Ax} = \mathbf{b} \end{aligned}$$

then *strong duality* holds.

Refinement of Slater's Condition

Theorem

Given a *convex* primal optimization problem.

If the first k constraint functions f_1, \dots, f_k are *affine*, and
if there exists an $\mathbf{x} \in \text{relint } \mathcal{D}$ with

$$f_i(\mathbf{x}) \leq 0, \quad i = 1, \dots, k \quad (\text{affine constraints})$$

$$f_i(\mathbf{x}) < 0, \quad i = k+1, \dots, m \quad (\text{convex constraints})$$

$$\mathbf{Ax} = \mathbf{b}$$

then *strong duality* holds.

Refinement of Slater's Condition

Theorem

Given a *convex* primal optimization problem.

If the first k constraint functions f_1, \dots, f_k are *affine*, and if there exists an $\mathbf{x} \in \text{relint } \mathcal{D}$ with

$$f_i(\mathbf{x}) \leq 0, \quad i = 1, \dots, k \quad (\text{affine constraints})$$

$$f_i(\mathbf{x}) < 0, \quad i = k + 1, \dots, m \quad (\text{convex constraints})$$

$$\mathbf{Ax} = \mathbf{b}$$

then *strong duality* holds.

Note: the refined Slater's condition reduces to *feasibility* when the constraints are all linear equalities and inequalities, and $\text{dom}(f_0)$ is open.

Karush-Kuhn-Tucker Optimality Conditions

Let \mathbf{x}^* be a primal and $(\boldsymbol{\lambda}^*, \boldsymbol{\nu}^*)$ dual optimal points with zero duality gap.

For the primal optimal point \mathbf{x}^* , the gradient with respect to \mathbf{x} of $L(\mathbf{x}, \boldsymbol{\lambda}^*, \boldsymbol{\nu}^*)$ is 0:

$$\nabla L(\mathbf{x}^*, \boldsymbol{\lambda}^*, \boldsymbol{\nu}^*) = \nabla f_0(\mathbf{x}^*) + \sum_{i=1}^m \lambda_i^* \nabla f_i(\mathbf{x}^*) + \sum_{i=1}^p \nu_i^* \nabla h_i(\mathbf{x}^*) = 0$$

Karush-Kuhn-Tucker Optimality Conditions

The following four conditions are called KKT conditions:

Karush-Kuhn-Tucker Optimality Conditions

The following four conditions are called KKT conditions:

1. Primal constraints:

- $f_i(\mathbf{x}) \leq 0, \quad i = 1, 2, \dots, m$
- $h_i(\mathbf{x}) = 0, \quad i = 1, 2, \dots, p$

Karush-Kuhn-Tucker Optimality Conditions

The following four conditions are called KKT conditions:

1. Primal constraints:

- $f_i(\mathbf{x}) \leq 0, \quad i = 1, 2, \dots, m$
- $h_i(\mathbf{x}) = 0, \quad i = 1, 2, \dots, p$

2. Dual constraints: $\lambda \succeq 0$

Karush-Kuhn-Tucker Optimality Conditions

The following four conditions are called KKT conditions:

1. Primal constraints:

- $f_i(\mathbf{x}) \leq 0, \quad i = 1, 2, \dots, m$
- $h_i(\mathbf{x}) = 0, \quad i = 1, 2, \dots, p$

2. Dual constraints: $\lambda \succeq 0$

3. Complementary slackness: $\lambda_i f_i(\mathbf{x}) = 0$

Karush-Kuhn-Tucker Optimality Conditions

The following four conditions are called KKT conditions:

1. Primal constraints:

- $f_i(\mathbf{x}) \leq 0, \quad i = 1, 2, \dots, m$
- $h_i(\mathbf{x}) = 0, \quad i = 1, 2, \dots, p$

2. Dual constraints: $\lambda \succeq 0$

3. Complementary slackness: $\lambda_i f_i(\mathbf{x}) = 0$

4. Gradient of the Lagrangian L is zero:

$$\nabla L(\mathbf{x}, \lambda, \nu) = \nabla f_0(\mathbf{x}) + \sum_{i=1}^m \lambda_i \nabla f_i(\mathbf{x}) + \sum_{i=1}^p \nu_i \nabla h_i(\mathbf{x}) = 0$$

Karush-Kuhn-Tucker Optimality Conditions

The following four conditions are called KKT conditions:

1. Primal constraints:

- $f_i(\mathbf{x}) \leq 0, \quad i = 1, 2, \dots, m$
- $h_i(\mathbf{x}) = 0, \quad i = 1, 2, \dots, p$

2. Dual constraints: $\lambda \succeq 0$

3. Complementary slackness: $\lambda_i f_i(\mathbf{x}) = 0$

4. Gradient of the Lagrangian L is zero:

$$\nabla L(\mathbf{x}, \lambda, \nu) = \nabla f_0(\mathbf{x}) + \sum_{i=1}^m \lambda_i \nabla f_i(\mathbf{x}) + \sum_{i=1}^p \nu_i \nabla h_i(\mathbf{x}) = 0$$

If strong duality holds and if \mathbf{x}^* and (λ^*, ν^*) are optimal points, then the KKT conditions hold.

Karush-Kuhn-Tucker Optimality Conditions

Complementary slackness

$$f_0(\mathbf{x}^*)$$

Karush-Kuhn-Tucker Optimality Conditions

Complementary slackness

$$f_0(\mathbf{x}^*) = g(\lambda^*, \nu^*)$$

Karush-Kuhn-Tucker Optimality Conditions

Complementary slackness

$$\begin{aligned} f_0(\mathbf{x}^*) &= g(\boldsymbol{\lambda}^*, \boldsymbol{\nu}^*) \\ &= \inf_{\mathbf{x}} \left(f_0(\mathbf{x}) + \sum_{i=1}^m \lambda_i^* f_i(\mathbf{x}) + \sum_{i=1}^p \nu_i^* h_i(\mathbf{x}) \right) \end{aligned}$$

Karush-Kuhn-Tucker Optimality Conditions

Complementary slackness

$$\begin{aligned} f_0(\mathbf{x}^*) &= g(\boldsymbol{\lambda}^*, \boldsymbol{\nu}^*) \\ &= \inf_{\mathbf{x}} \left(f_0(\mathbf{x}) + \sum_{i=1}^m \lambda_i^* f_i(\mathbf{x}) + \sum_{i=1}^p \nu_i^* h_i(\mathbf{x}) \right) \\ &\leq f_0(\mathbf{x}^*) + \sum_{i=1}^m \lambda_i^* f_i(\mathbf{x}^*) + \sum_{i=1}^p \nu_i^* h_i(\mathbf{x}^*) \end{aligned}$$

Karush-Kuhn-Tucker Optimality Conditions

Complementary slackness

$$\begin{aligned} f_0(\mathbf{x}^*) &= g(\boldsymbol{\lambda}^*, \boldsymbol{\nu}^*) \\ &= \inf_{\mathbf{x}} \left(f_0(\mathbf{x}) + \sum_{i=1}^m \lambda_i^* f_i(\mathbf{x}) + \sum_{i=1}^p \nu_i^* h_i(\mathbf{x}) \right) \\ &\leq f_0(\mathbf{x}^*) + \sum_{i=1}^m \lambda_i^* f_i(\mathbf{x}^*) + \sum_{i=1}^p \nu_i^* h_i(\mathbf{x}^*) \end{aligned}$$

Karush-Kuhn-Tucker Optimality Conditions

Complementary slackness

$$\begin{aligned}
 f_0(\mathbf{x}^*) &= g(\boldsymbol{\lambda}^*, \boldsymbol{\nu}^*) \\
 &= \inf_{\mathbf{x}} \left(f_0(\mathbf{x}) + \sum_{i=1}^m \lambda_i^* f_i(\mathbf{x}) + \sum_{i=1}^p \nu_i^* h_i(\mathbf{x}) \right) \\
 &\leq f_0(\mathbf{x}^*) + \sum_{i=1}^m \lambda_i^* f_i(\mathbf{x}^*) + \underbrace{\sum_{i=1}^p \nu_i^* h_i(\mathbf{x}^*)}_{=0}
 \end{aligned}$$

Karush-Kuhn-Tucker Optimality Conditions

Complementary slackness

$$\begin{aligned}
 f_0(\mathbf{x}^*) &= g(\boldsymbol{\lambda}^*, \boldsymbol{\nu}^*) \\
 &= \inf_{\mathbf{x}} \left(f_0(\mathbf{x}) + \sum_{i=1}^m \lambda_i^* f_i(\mathbf{x}) + \sum_{i=1}^p \nu_i^* h_i(\mathbf{x}) \right) \\
 &\leq f_0(\mathbf{x}^*) + \underbrace{\sum_{i=1}^m \lambda_i^* f_i(\mathbf{x}^*) + \sum_{i=1}^p \nu_i^* h_i(\mathbf{x}^*)}_{=0}
 \end{aligned}$$

Karush-Kuhn-Tucker Optimality Conditions

Complementary slackness

$$\begin{aligned}
 f_0(\mathbf{x}^*) &= g(\boldsymbol{\lambda}^*, \boldsymbol{\nu}^*) \\
 &= \inf_{\mathbf{x}} \left(f_0(\mathbf{x}) + \sum_{i=1}^m \lambda_i^* f_i(\mathbf{x}) + \sum_{i=1}^p \nu_i^* h_i(\mathbf{x}) \right) \\
 &\leq f_0(\mathbf{x}^*) + \underbrace{\sum_{i=1}^m \lambda_i^* f_i(\mathbf{x}^*)}_{\leq 0} + \underbrace{\sum_{i=1}^p \nu_i^* h_i(\mathbf{x}^*)}_{= 0}
 \end{aligned}$$

Karush-Kuhn-Tucker Optimality Conditions

Complementary slackness

$$\begin{aligned}
 f_0(\mathbf{x}^*) &= g(\boldsymbol{\lambda}^*, \boldsymbol{\nu}^*) \\
 &= \inf_{\mathbf{x}} \left(f_0(\mathbf{x}) + \sum_{i=1}^m \lambda_i^* f_i(\mathbf{x}) + \sum_{i=1}^p \nu_i^* h_i(\mathbf{x}) \right) \\
 &\leq f_0(\mathbf{x}^*) + \underbrace{\sum_{i=1}^m \lambda_i^* f_i(\mathbf{x}^*)}_{\leq 0} + \underbrace{\sum_{i=1}^p \nu_i^* h_i(\mathbf{x}^*)}_{= 0} \\
 &\leq f_0(\mathbf{x}^*)
 \end{aligned}$$

Karush-Kuhn-Tucker Optimality Conditions

Complementary slackness

$$\begin{aligned}
 f_0(\mathbf{x}^*) &= g(\boldsymbol{\lambda}^*, \boldsymbol{\nu}^*) \\
 &= \inf_{\mathbf{x}} \left(f_0(\mathbf{x}) + \sum_{i=1}^m \lambda_i^* f_i(\mathbf{x}) + \sum_{i=1}^p \nu_i^* h_i(\mathbf{x}) \right) \\
 &\leq f_0(\mathbf{x}^*) + \underbrace{\sum_{i=1}^m \lambda_i^* f_i(\mathbf{x}^*)}_{\leq 0} + \underbrace{\sum_{i=1}^p \nu_i^* h_i(\mathbf{x}^*)}_{= 0} \\
 &\leq f_0(\mathbf{x}^*)
 \end{aligned}$$

Karush-Kuhn-Tucker Optimality Conditions

Complementary slackness

$$\begin{aligned}
 f_0(\mathbf{x}^*) &= g(\boldsymbol{\lambda}^*, \boldsymbol{\nu}^*) \\
 &= \inf_{\mathbf{x}} \left(f_0(\mathbf{x}) + \sum_{i=1}^m \lambda_i^* f_i(\mathbf{x}) + \sum_{i=1}^p \nu_i^* h_i(\mathbf{x}) \right) \\
 &\leq f_0(\mathbf{x}^*) + \underbrace{\sum_{i=1}^m \lambda_i^* f_i(\mathbf{x}^*)}_{\leq 0} + \underbrace{\sum_{i=1}^p \nu_i^* h_i(\mathbf{x}^*)}_{= 0} \\
 &\stackrel{=}{\leq} f_0(\mathbf{x}^*)
 \end{aligned}$$

Karush-Kuhn-Tucker Optimality Conditions

Complementary slackness

$$\begin{aligned}
 f_0(\mathbf{x}^*) &= g(\boldsymbol{\lambda}^*, \boldsymbol{\nu}^*) \\
 &= \inf_{\mathbf{x}} \left(f_0(\mathbf{x}) + \sum_{i=1}^m \lambda_i^* f_i(\mathbf{x}) + \sum_{i=1}^p \nu_i^* h_i(\mathbf{x}) \right) \\
 &\stackrel{=}{\neq} f_0(\mathbf{x}^*) + \underbrace{\sum_{i=1}^m \lambda_i^* f_i(\mathbf{x}^*)}_{\leq 0} + \underbrace{\sum_{i=1}^p \nu_i^* h_i(\mathbf{x}^*)}_{=0} \\
 &\stackrel{=}{\neq} f_0(\mathbf{x}^*)
 \end{aligned}$$

Karush-Kuhn-Tucker Optimality Conditions

Complementary slackness

$$\begin{aligned}
 f_0(\mathbf{x}^*) &= g(\boldsymbol{\lambda}^*, \boldsymbol{\nu}^*) \\
 &= \inf_{\mathbf{x}} \left(f_0(\mathbf{x}) + \sum_{i=1}^m \lambda_i^* f_i(\mathbf{x}) + \sum_{i=1}^p \nu_i^* h_i(\mathbf{x}) \right) \\
 &\stackrel{=}{\neq} f_0(\mathbf{x}^*) + \underbrace{\sum_{i=1}^m \lambda_i^* f_i(\mathbf{x}^*)}_{\stackrel{=}{\neq} 0} + \underbrace{\sum_{i=1}^p \nu_i^* h_i(\mathbf{x}^*)}_{= 0} \\
 &\stackrel{=}{\neq} f_0(\mathbf{x}^*)
 \end{aligned}$$

Karush-Kuhn-Tucker Optimality Conditions

Complementary slackness

$$\begin{aligned}
 f_0(\mathbf{x}^*) &= g(\boldsymbol{\lambda}^*, \boldsymbol{\nu}^*) \\
 &= \inf_{\mathbf{x}} \left(f_0(\mathbf{x}) + \sum_{i=1}^m \lambda_i^* f_i(\mathbf{x}) + \sum_{i=1}^p \nu_i^* h_i(\mathbf{x}) \right) \\
 &\stackrel{=}{\neq} f_0(\mathbf{x}^*) + \underbrace{\sum_{i=1}^m \lambda_i^* f_i(\mathbf{x}^*)}_{\stackrel{=}{\neq} 0} + \underbrace{\sum_{i=1}^p \nu_i^* h_i(\mathbf{x}^*)}_{= 0} \\
 &\stackrel{=}{\neq} f_0(\mathbf{x}^*)
 \end{aligned}$$

Karush-Kuhn-Tucker Optimality Conditions

Complementary slackness: $\lambda_i^* \cdot f_i(\mathbf{x}^*) = 0$

$$\begin{aligned}
 f_0(\mathbf{x}^*) &= g(\lambda^*, \nu^*) \\
 &= \inf_{\mathbf{x}} \left(f_0(\mathbf{x}) + \sum_{i=1}^m \lambda_i^* f_i(\mathbf{x}) + \sum_{i=1}^p \nu_i^* h_i(\mathbf{x}) \right) \\
 &\stackrel{=}{\neq} f_0(\mathbf{x}^*) + \underbrace{\sum_{i=1}^m \underbrace{\lambda_i^* f_i(\mathbf{x}^*)}_{=0}}_{\stackrel{=}{\neq} 0} + \underbrace{\sum_{i=1}^p \nu_i^* h_i(\mathbf{x}^*)}_{=0} \\
 &\stackrel{=}{\neq} f_0(\mathbf{x}^*)
 \end{aligned}$$



Karush-Kuhn-Tucker Optimality Conditions (cont.)

Conclusions (Boyd 2004, Sec. 5.5.3)

- For *any* optimization problem with differentiable objective and constraint functions for which strong duality obtains, any pair of primal and dual optimal points must satisfy the KKT conditions.

Karush-Kuhn-Tucker Optimality Conditions (cont.)

Conclusions (Boyd 2004, Sec. 5.5.3)

- For *any optimization problem* with differentiable objective and constraint functions for which strong duality obtains, any pair of primal and dual optimal points must satisfy the KKT conditions.
- For *any convex optimization problem* with differentiable objective and and constraint functions, any points that satisfy the KKT conditions are primal and dual optimal, and have zero duality gap.

Karush-Kuhn-Tucker Optimality Conditions (cont.)

Conclusions (Boyd 2004, Sec. 5.5.3)

- For *any optimization problem* with differentiable objective and constraint functions for which strong duality obtains, any pair of primal and dual optimal points must satisfy the KKT conditions.
- For *any convex optimization problem* with differentiable objective and and constraint functions, any points that satisfy the KKT conditions are primal and dual optimal, and have zero duality gap.
- If a *convex optimization problem* with differentiable objective and constraint functions satisfies Slater's condition, then the KKT conditions provide necessary and sufficient conditions for optimality.

Lessons Learned

- Formalization of the primal problem using the Lagrangian
- Lagrange dual function
- Duality gap
- Karush-Kuhn-Tucker optimality conditions



**Pattern
Recognition
Lab**



**FRIEDRICH-ALEXANDER
UNIVERSITÄT
ERLANGEN-NÜRNBERG**


TECHNISCHE FAKULTÄT

Next Time in

Pattern Recognition



Further Readings

- S. Boyd, L. Vandenberghe:
[Convex Optimization](#),
Cambridge University Press, 2004.
 <http://www.stanford.edu/~boyd/cvxbook/>
- Jorge Nocedal, Stephen Wright:
[Numerical Optimization](#),
Springer, New York, 1999.

Comprehensive Questions

- What is the Lagrangian of a constrained objective function?
- What is the Lagrange dual function?
- What is the duality gap?
- What are the Karush-Kuhn-Tucker optimality conditions?