

Arctic Termination ... Below Zero

Adam Koprowski¹ Johannes Waldmann²

¹Eindhoven University of Technology

²Hochschule für Technik, Wirtschaft und Kultur (FH) Leipzig

15 July 2008
RTA, Hagenberg

Outline

- 1 Introduction
- 2 Monotone Algebras
- 3 Polynomial and Matrix Interpretations
- 4 Arctic Interpretations
- 5 Arctic Below Zero Interpretations
- 6 Certification
- 7 Evaluation
- 8 Conclusions

- 1 Introduction
- 2 Monotone Algebras
- 3 Polynomial and Matrix Interpretations
- 4 Arctic Interpretations
- 5 Arctic Below Zero Interpretations
- 6 Certification
- 7 Evaluation
- 8 Conclusions

Termination of rewriting

- Termination of rewriting is an important property of TRSs.

Termination of rewriting

- Termination of rewriting is an important property of TRSs.
- There are many methods to prove termination of rewriting

Termination of rewriting

- Termination of rewriting is an important property of TRSs.
- There are many methods to prove termination of rewriting
- and recently the emphasis is on automation.

Termination of rewriting

- Termination of rewriting is an important property of TRSs.
- There are many methods to prove termination of rewriting
- and recently the emphasis is on automation.
- There exists a number of tools for proving termination

Termination of rewriting

- Termination of rewriting is an important property of TRSs.
- There are many methods to prove termination of rewriting
- and recently the emphasis is on automation.
- There exists a number of tools for proving termination
- and the new developments are stimulated by an annual termination competition.

Termination of rewriting

- Termination of rewriting is an important property of TRSs.
- There are many methods to prove termination of rewriting
- and recently the emphasis is on automation.
- There exists a number of tools for proving termination
- and the new developments are stimulated by an annual termination competition.
- Tools, and the proofs they produce, are getting more and more complex,

Termination of rewriting

- Termination of rewriting is an important property of TRSs.
- There are many methods to prove termination of rewriting
- and recently the emphasis is on automation.
- There exists a number of tools for proving termination
- and the new developments are stimulated by an annual termination competition.
- Tools, and the proofs they produce, are getting more and more complex,
- hence the recent work on certification of termination.

Termination of rewriting

- Termination of rewriting is an important property of TRSs.
- There are many methods to prove termination of rewriting
- and recently the emphasis is on automation.
- There exists a number of tools for proving termination
- and the new developments are stimulated by an annual termination competition.
- Tools, and the proofs they produce, are getting more and more complex,
- hence the recent work on certification of termination.

⇒ This talk concerns a **new method for proving termination**, its **automation** and **certification**.

Outline

- 1 Introduction
- 2 Monotone Algebras**
- 3 Polynomial and Matrix Interpretations
- 4 Arctic Interpretations
- 5 Arctic Below Zero Interpretations
- 6 Certification
- 7 Evaluation
- 8 Conclusions

Given TRSs \mathcal{R} and \mathcal{S} define:

- *top rewrite relation*: $t \xrightarrow{\text{top}}_{\mathcal{R}} u$ if and only if there is a rewrite rule $\ell \rightarrow r \in \mathcal{R}$ and a substitution $\sigma : \mathcal{V} \rightarrow \mathcal{T}(\Sigma, \mathcal{V})$ such that $t = \ell\sigma$ and $u = r\sigma$.

Given TRSs \mathcal{R} and \mathcal{S} define:

- *top rewrite relation*: $t \xrightarrow{\text{top}}_{\mathcal{R}} u$ if and only if there is a rewrite rule $\ell \rightarrow r \in \mathcal{R}$ and a substitution $\sigma : \mathcal{V} \rightarrow \mathcal{T}(\Sigma, \mathcal{V})$ such that $t = \ell\sigma$ and $u = r\sigma$.
- *rewrite relation*: $\rightarrow_{\mathcal{R}}$ is the smallest relation such that $\xrightarrow{\text{top}}_{\mathcal{R}} \subseteq \rightarrow_{\mathcal{R}}$ and $\rightarrow_{\mathcal{R}}$ is context-closed.

Given TRSs \mathcal{R} and \mathcal{S} define:

- *top rewrite relation*: $t \xrightarrow{\text{top}}_{\mathcal{R}} u$ if and only if there is a rewrite rule $\ell \rightarrow r \in \mathcal{R}$ and a substitution $\sigma : \mathcal{V} \rightarrow \mathcal{T}(\Sigma, \mathcal{V})$ such that $t = \ell\sigma$ and $u = r\sigma$.
- *rewrite relation*: $\rightarrow_{\mathcal{R}}$ is the smallest relation such that $\xrightarrow{\text{top}}_{\mathcal{R}} \subseteq \rightarrow_{\mathcal{R}}$ and $\rightarrow_{\mathcal{R}}$ is context-closed.
- *relation modulo*: $\rightarrow_1 / \rightarrow_2 \equiv \rightarrow_2^* \cdot \rightarrow_1$.

Basic definitions

Given TRSs \mathcal{R} and \mathcal{S} define:

- *top rewrite relation*: $t \xrightarrow{\text{top}}_{\mathcal{R}} u$ if and only if there is a rewrite rule $\ell \rightarrow r \in \mathcal{R}$ and a substitution $\sigma : \mathcal{V} \rightarrow \mathcal{T}(\Sigma, \mathcal{V})$ such that $t = \ell\sigma$ and $u = r\sigma$.
- *rewrite relation*: $\rightarrow_{\mathcal{R}}$ is the smallest relation such that $\xrightarrow{\text{top}}_{\mathcal{R}} \subseteq \rightarrow_{\mathcal{R}}$ and $\rightarrow_{\mathcal{R}}$ is context-closed.
- *relation modulo*: $\rightarrow_1 / \rightarrow_2 \equiv \rightarrow_2^* \cdot \rightarrow_1$.
- *termination*: $\text{SN}(\rightarrow_{\mathcal{R}})$.

Basic definitions

Given TRSs \mathcal{R} and \mathcal{S} define:

- *top rewrite relation*: $t \xrightarrow{\text{top}}_{\mathcal{R}} u$ if and only if there is a rewrite rule $\ell \rightarrow r \in \mathcal{R}$ and a substitution $\sigma : \mathcal{V} \rightarrow \mathcal{T}(\Sigma, \mathcal{V})$ such that $t = \ell\sigma$ and $u = r\sigma$.
- *rewrite relation*: $\rightarrow_{\mathcal{R}}$ is the smallest relation such that $\xrightarrow{\text{top}}_{\mathcal{R}} \subseteq \rightarrow_{\mathcal{R}}$ and $\rightarrow_{\mathcal{R}}$ is context-closed.
- *relation modulo*: $\rightarrow_1 / \rightarrow_2 \equiv \rightarrow_2^* \cdot \rightarrow_1$.
- *termination*: $\text{SN}(\rightarrow_{\mathcal{R}})$.
- *relative termination*: $\text{SN}(\rightarrow_{\mathcal{R}} / \rightarrow_{\mathcal{S}})$.

Basic definitions

Given TRSs \mathcal{R} and \mathcal{S} define:

- *top rewrite relation*: $t \xrightarrow{\text{top}}_{\mathcal{R}} u$ if and only if there is a rewrite rule $\ell \rightarrow r \in \mathcal{R}$ and a substitution $\sigma : \mathcal{V} \rightarrow \mathcal{T}(\Sigma, \mathcal{V})$ such that $t = \ell\sigma$ and $u = r\sigma$.
- *rewrite relation*: $\rightarrow_{\mathcal{R}}$ is the smallest relation such that $\xrightarrow{\text{top}}_{\mathcal{R}} \subseteq \rightarrow_{\mathcal{R}}$ and $\rightarrow_{\mathcal{R}}$ is context-closed.
- *relation modulo*: $\rightarrow_1 / \rightarrow_2 \equiv \rightarrow_2^* \cdot \rightarrow_1$.
- *termination*: $\text{SN}(\rightarrow_{\mathcal{R}})$.
- *relative termination*: $\text{SN}(\rightarrow_{\mathcal{R}} / \rightarrow_{\mathcal{S}})$.
- *relative top termination*: $\text{SN}(\xrightarrow{\text{top}}_{\mathcal{R}} / \rightarrow_{\mathcal{S}})$ (important in the dependency pairs setting).

Definition (Monotonicity)

An operation $[f] : A \times \dots \times A \rightarrow A$ is *monotone* with respect to a binary relation \triangleright on A if

$$a_i \triangleright a'_i \implies [f](a_1, \dots, a_i, \dots, a_n) \triangleright [f](a_1, \dots, a'_i, \dots, a_n).$$

Monotone algebras

Definition (Monotonicity)

An operation $[f] : A \times \dots \times A \rightarrow A$ is *monotone* with respect to a binary relation \triangleright on A if

$$a_i \triangleright a'_i \implies [f](a_1, \dots, a_i, \dots, a_n) \triangleright [f](a_1, \dots, a'_i, \dots, a_n).$$

Definition (Monotone Σ -algebras)

A *weakly monotone Σ -algebra* $(A, [\cdot], >, \gtrsim)$ is a Σ -algebra $(A, [\cdot])$ equipped with two binary relations $>, \gtrsim$ on A such that

- $>$ is well-founded;
- $> \cdot \gtrsim \subseteq >$;
- for every $f \in \Sigma$ the operation $[f]$ is monotone with respect to \gtrsim .

An *extended monotone Σ -algebra* $(A, [\cdot], >, \gtrsim)$ is a weakly monotone Σ -algebra $(A, [\cdot], >, \gtrsim)$ in which moreover for every $f \in \Sigma$ the operation $[f]$ is monotone with respect to $>$.

Theorem

Let $\mathcal{R}, \mathcal{R}', S, S'$ be TRSs over a signature Σ , $(A, [\cdot], >, \succeq)$ be an extended monotone Σ -algebra such that:

- $\forall_{\alpha} [\ell]_{\alpha} \succeq [r]_{\alpha}$ for every rule $\ell \rightarrow r$ in $\mathcal{R} \cup S$ and
- $\forall_{\alpha} [\ell]_{\alpha} > [r]_{\alpha}$ for every rule $\ell \rightarrow r$ in $\mathcal{R}' \cup S'$

Then $\text{SN}(\rightarrow_{\mathcal{R}} / \rightarrow_S)$ implies $\text{SN}(\rightarrow_{\mathcal{R}} \cup \rightarrow_{\mathcal{R}'} / \rightarrow_S \cup \rightarrow_{S'})$.

Theorem

Let $\mathcal{R}, \mathcal{R}', S, S'$ be TRSs over a signature Σ , $(A, [\cdot], >, \gtrsim)$ be an extended monotone Σ -algebra such that:

- $\forall_{\alpha} [\ell]_{\alpha} \gtrsim [r]_{\alpha}$ for every rule $\ell \rightarrow r$ in $\mathcal{R} \cup S$ and
- $\forall_{\alpha} [\ell]_{\alpha} > [r]_{\alpha}$ for every rule $\ell \rightarrow r$ in $\mathcal{R}' \cup S'$

Then $\text{SN}(\rightarrow_{\mathcal{R}} / \rightarrow_S)$ implies $\text{SN}(\rightarrow_{\mathcal{R}} \cup \rightarrow_{\mathcal{R}'} / \rightarrow_S \cup \rightarrow_{S'})$.

Theorem

Let $\mathcal{R}, \mathcal{R}', S, S'$ be TRSs over a signature Σ , let $(A, [\cdot], >, \gtrsim)$ be a weakly monotone Σ -algebra such that:

- $\forall_{\alpha} [\ell]_{\alpha} \gtrsim [r]_{\alpha}$ for every rule $\ell \rightarrow r$ in $\mathcal{R} \cup S$ and
- $\forall_{\alpha} [\ell]_{\alpha} > [r]_{\alpha}$ for every rule $\ell \rightarrow r$ in \mathcal{R}' ,

Then $\text{SN}(\overset{\text{top}}{\rightarrow}_{\mathcal{R}} / \rightarrow_S)$ implies $\text{SN}(\overset{\text{top}}{\rightarrow}_{\mathcal{R}} \cup \overset{\text{top}}{\rightarrow}_{\mathcal{R}'} / \rightarrow_S)$.

Outline

- 1 Introduction
- 2 Monotone Algebras
- 3 Polynomial and Matrix Interpretations**
- 4 Arctic Interpretations
- 5 Arctic Below Zero Interpretations
- 6 Certification
- 7 Evaluation
- 8 Conclusions

Polynomial interpretations

- Interpretation domain: \mathbb{N} .

Polynomial interpretations

- Interpretation domain: \mathbb{N} .
- Interpretations: polynomials over \mathbb{N} .

Polynomial interpretations

- Interpretation domain: \mathbb{N} .
- Interpretations: polynomials over \mathbb{N} .

Example

$$x * (y + z) \rightarrow x * y + x * z$$

Polynomial interpretations

- Interpretation domain: \mathbb{N} .
- Interpretations: polynomials over \mathbb{N} .

Example

$$x * (y + z) \rightarrow x * y + x * z$$

$$[x + y] = x + y + 2, \quad [x * y] = 2x + 2y + 2xy + 1$$

Polynomial interpretations

- Interpretation domain: \mathbb{N} .
- Interpretations: polynomials over \mathbb{N} .

Example

$$x * (y + z) \rightarrow x * y + x * z$$

$$[x + y] = x + y + 2, \quad [x * y] = 2x + 2y + 2xy + 1$$

$$[x * (y + z)] = 2x + 2(y + z + 2) + 2x(y + z + 2) + 1$$

$$[x * y + x * z] = (2x + 2y + 2xy + 1) + (2x + 2z + 2xz + 1) + 2$$

Polynomial interpretations

- Interpretation domain: \mathbb{N} .
- Interpretations: polynomials over \mathbb{N} .

Example

$$x * (y + z) \rightarrow x * y + x * z$$

$$[x + y] = x + y + 2, \quad [x * y] = 2x + 2y + 2xy + 1$$

$$[x * (y + z)] = 2x + 2y + 2z + 4 + 2xy + 2xz + 4x + 1$$

$$[x * y + x * z] = 2x + 2y + 2xy + 1 + 2x + 2z + 2xz + 1 + 2$$

Polynomial interpretations

- Interpretation domain: \mathbb{N} .
- Interpretations: polynomials over \mathbb{N} .

Example

$$x * (y + z) \rightarrow x * y + x * z$$

$$[x + y] = x + y + 2, \quad [x * y] = 2x + 2y + 2xy + 1$$

$$[x * (y + z)] = 6x + 2y + 2z + 2xy + 2xz + 5$$

$$[x * y + x * z] = 4x + 2y + 2z + 2xy + 2xz + 4$$

Polynomial interpretations

- Interpretation domain: \mathbb{N} .
- Interpretations: polynomials over \mathbb{N} .

Example

$$x * (y + z) \rightarrow x * y + x * z$$

$$[x + y] = x + y + 2, \quad [x * y] = 2x + 2y + 2xy + 1$$

$$[x * (y + z)] = 6x + 2y + 2z + 2xy + 2xz + 5$$

$$[x * y + x * z] = 4x + 2y + 2z + 2xy + 2xz + 4$$

- To obtain strict monotonicity we require that for every interpretation $[f(x_1, \dots, x_n)]$, $\forall_i \exists_{c>0} cx_i \in [f(x_1, \dots, x_n)]$.

Matrix interpretations

- Interpretation domain: \mathbb{N}^d , for some fixed d .

Matrix interpretations

- Interpretation domain: \mathbb{N}^d , for some fixed d .
- $\vec{u} \geq \vec{v}$ iff $\forall_i \vec{u}_i \geq \vec{v}_i$.

Matrix interpretations

- Interpretation domain: \mathbb{N}^d , for some fixed d .
- $\vec{u} \geq \vec{v}$ iff $\forall_i \vec{u}_i \geq \vec{v}_i$.
- $\vec{u} > \vec{v}$ iff $\vec{u} \geq \vec{v} \wedge \vec{u}_1 > \vec{v}_1$.

Matrix interpretations

- Interpretation domain: \mathbb{N}^d , for some fixed d .
- $\vec{u} \geq \vec{v}$ iff $\forall_i \vec{u}_i \geq \vec{v}_i$.
- $\vec{u} > \vec{v}$ iff $\vec{u} \geq \vec{v} \wedge \vec{u}_1 > \vec{v}_1$.

Example

$$a(a(x)) \rightarrow a(b(a(x))).$$

Matrix interpretations

- Interpretation domain: \mathbb{N}^d , for some fixed d .
- $\vec{u} \geq \vec{v}$ iff $\forall_i \vec{u}_i \geq \vec{v}_i$.
- $\vec{u} > \vec{v}$ iff $\vec{u} \geq \vec{v} \wedge \vec{u}_1 > \vec{v}_1$.

Example

$$a(a(x)) \rightarrow a(b(a(x))).$$

$$[a(x)] = \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix} x + \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad [b(x)] = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} x + \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

Matrix interpretations

- Interpretation domain: \mathbb{N}^d , for some fixed d .
- $\vec{u} \geq \vec{v}$ iff $\forall_i \vec{u}_i \geq \vec{v}_i$.
- $\vec{u} > \vec{v}$ iff $\vec{u} \geq \vec{v} \wedge \vec{u}_1 > \vec{v}_1$.

Example

$$a(a(x)) \rightarrow a(b(a(x))).$$

$$[a(x)] = \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix} x + \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad [b(x)] = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} x + \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$[a(a(x))] = \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix} \left(\begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix} x + \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right) + \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

$$[a(b(a(x)))] = \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix} \left(\begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \left(\begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix} x + \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right) + \begin{pmatrix} 0 \\ 0 \end{pmatrix} \right) + \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

Matrix interpretations

- Interpretation domain: \mathbb{N}^d , for some fixed d .
- $\vec{u} \geq \vec{v}$ iff $\forall_i \vec{u}_i \geq \vec{v}_i$.
- $\vec{u} > \vec{v}$ iff $\vec{u} \geq \vec{v} \wedge \vec{u}_1 > \vec{v}_1$.

Example

$$a(a(x)) \rightarrow a(b(a(x))).$$

$$[a(x)] = \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix} x + \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad [b(x)] = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} x + \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$[a(a(x))] = \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix} x + \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

$$[a(b(a(x)))] = \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix} x + \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

Matrix interpretations

- Interpretation domain: \mathbb{N}^d , for some fixed d .
- $\vec{u} \geq \vec{v}$ iff $\forall_i \vec{u}_i \geq \vec{v}_i$.
- $\vec{u} > \vec{v}$ iff $\vec{u} \geq \vec{v} \wedge \vec{u}_1 > \vec{v}_1$.

Example

$$a(a(x)) \rightarrow a(b(a(x))).$$

$$[a(x)] = \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix} x + \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad [b(x)] = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} x + \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$[a(a(x))] = \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix} x + \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

$$[a(b(a(x)))] = \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix} x + \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

- Now we need to restrict to linear interpretations.

Matrix interpretations

- Interpretation domain: \mathbb{N}^d , for some fixed d .
- $\vec{u} \geq \vec{v}$ iff $\forall_i \vec{u}_i \geq \vec{v}_i$.
- $\vec{u} > \vec{v}$ iff $\vec{u} \geq \vec{v} \wedge \vec{u}_1 > \vec{v}_1$.

Example

$$a(a(x)) \rightarrow a(b(a(x))).$$

$$[a(x)] = \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix} x + \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad [b(x)] = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} x + \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$[a(a(x))] = \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix} x + \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

$$[a(b(a(x)))] = \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix} x + \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

- Now we need to restrict to linear interpretations.
- Strict monotonicity ensured if for every interpretation $[f(x_1, \dots, x_n)] = F_1 x_1 + \dots F_n x_n + \vec{f}$ we have $\forall_i (F_i)_{1,1} > 0$.

Outline

- 1 Introduction
- 2 Monotone Algebras
- 3 Polynomial and Matrix Interpretations
- 4 Arctic Interpretations**
- 5 Arctic Below Zero Interpretations
- 6 Certification
- 7 Evaluation
- 8 Conclusions

Arctic interpretations

- $\mathbb{A}_{\mathbb{N}} \equiv \{-\infty\} \cup \mathbb{N}$.

Arctic interpretations

- $\mathbb{A}_{\mathbb{N}} \equiv \{-\infty\} \cup \mathbb{N}$.
- We say that $a \in \mathbb{A}_{\mathbb{N}}$ is *finite* iff $a \neq -\infty$.

Arctic interpretations

- $\mathbb{A}_{\mathbb{N}} \equiv \{-\infty\} \cup \mathbb{N}$.
- We say that $a \in \mathbb{A}_{\mathbb{N}}$ is *finite* iff $a \neq -\infty$.
- Interpretation domain: $\mathbb{A}_{\mathbb{N}}^d$, for some fixed d .

Arctic interpretations

- $\mathbb{A}_{\mathbb{N}} \equiv \{-\infty\} \cup \mathbb{N}$.
- We say that $a \in \mathbb{A}_{\mathbb{N}}$ is *finite* iff $a \neq -\infty$.
- Interpretation domain: $\mathbb{A}_{\mathbb{N}}^d$, for some fixed d .
- Now we compute in the $\langle \mathbb{A}_{\mathbb{N}}, \max, + \rangle$ semi-ring.

Arctic interpretations

- $\mathbb{A}_{\mathbb{N}} \equiv \{-\infty\} \cup \mathbb{N}$.
- We say that $a \in \mathbb{A}_{\mathbb{N}}$ is *finite* iff $a \neq -\infty$.
- Interpretation domain: $\mathbb{A}_{\mathbb{N}}^d$, for some fixed d .
- Now we compute in the $\langle \mathbb{A}_{\mathbb{N}}, \max, + \rangle$ semi-ring.
- $a \gg b \equiv a > b \vee (a = b = -\infty)$.

Arctic interpretations

- $\mathbb{A}_{\mathbb{N}} \equiv \{-\infty\} \cup \mathbb{N}$.
- We say that $a \in \mathbb{A}_{\mathbb{N}}$ is *finite* iff $a \neq -\infty$.
- Interpretation domain: $\mathbb{A}_{\mathbb{N}}^d$, for some fixed d .
- Now we compute in the $\langle \mathbb{A}_{\mathbb{N}}, \max, + \rangle$ semi-ring.
- $a \gg b \equiv a > b \vee (a = b = -\infty)$.
- $\vec{u} \geq \vec{v}$ iff $\forall_i \vec{u}_i \geq \vec{v}_i$.

Arctic interpretations

- $\mathbb{A}_{\mathbb{N}} \equiv \{-\infty\} \cup \mathbb{N}$.
- We say that $a \in \mathbb{A}_{\mathbb{N}}$ is *finite* iff $a \neq -\infty$.
- Interpretation domain: $\mathbb{A}_{\mathbb{N}}^d$, for some fixed d .
- Now we compute in the $\langle \mathbb{A}_{\mathbb{N}}, \max, + \rangle$ semi-ring.
- $a \gg b \equiv a > b \vee (a = b = -\infty)$.
- $\vec{u} \geq \vec{v}$ iff $\forall_i \vec{u}_i \geq \vec{v}_i$.
- $\vec{u} > \vec{v}$ iff $\forall_i \vec{u}_i \gg \vec{v}_i$.

Arctic interpretations

- $\mathbb{A}_{\mathbb{N}} \equiv \{-\infty\} \cup \mathbb{N}$.
- We say that $a \in \mathbb{A}_{\mathbb{N}}$ is *finite* iff $a \neq -\infty$.
- Interpretation domain: $\mathbb{A}_{\mathbb{N}}^d$, for some fixed d .
- Now we compute in the $\langle \mathbb{A}_{\mathbb{N}}, \max, + \rangle$ semi-ring.
- $a \gg b \equiv a > b \vee (a = b = -\infty)$.
- $\vec{u} \geq \vec{v}$ iff $\forall_i \vec{u}_i \geq \vec{v}_i$.
- $\vec{u} > \vec{v}$ iff $\forall_i \vec{u}_i \gg \vec{v}_i$.
- **Problem:** arctic addition is not strictly monotonic in single arguments, ie. $5 > 3$ but $5 \oplus 6 = 6 \not> 6 = 3 \oplus 6$. We cannot get strict monotonicity for symbols of arity > 1 .

Arctic interpretations

- $\mathbb{A}_{\mathbb{N}} \equiv \{-\infty\} \cup \mathbb{N}$.
- We say that $a \in \mathbb{A}_{\mathbb{N}}$ is *finite* iff $a \neq -\infty$.
- Interpretation domain: $\mathbb{A}_{\mathbb{N}}^d$, for some fixed d .
- Now we compute in the $\langle \mathbb{A}_{\mathbb{N}}, \max, + \rangle$ semi-ring.
- $a \gg b \equiv a > b \vee (a = b = -\infty)$.
- $\vec{u} \geq \vec{v}$ iff $\forall_i \vec{u}_i \geq \vec{v}_i$.
- $\vec{u} > \vec{v}$ iff $\forall_i \vec{u}_i \gg \vec{v}_i$.
- **Problem:** arctic addition is not strictly monotonic in single arguments, ie. $5 > 3$ but $5 \oplus 6 = 6 \not> 6 = 3 \oplus 6$. We cannot get strict monotonicity for symbols of arity > 1 .
- \Rightarrow full termination only for SRSs (as used in Matchbox 2007).

Arctic interpretations

- $\mathbb{A}_{\mathbb{N}} \equiv \{-\infty\} \cup \mathbb{N}$.
- We say that $a \in \mathbb{A}_{\mathbb{N}}$ is *finite* iff $a \neq -\infty$.
- Interpretation domain: $\mathbb{A}_{\mathbb{N}}^d$, for some fixed d .
- Now we compute in the $\langle \mathbb{A}_{\mathbb{N}}, \max, + \rangle$ semi-ring.
- $a \gg b \equiv a > b \vee (a = b = -\infty)$.
- $\vec{u} \geq \vec{v}$ iff $\forall_i \vec{u}_i \geq \vec{v}_i$.
- $\vec{u} > \vec{v}$ iff $\forall_i \vec{u}_i \gg \vec{v}_i$.
- **Problem:** arctic addition is not strictly monotonic in single arguments, ie. $5 > 3$ but $5 \oplus 6 = 6 \not> 6 = 3 \oplus 6$. We cannot get strict monotonicity for symbols of arity > 1 .
- \Rightarrow full termination only for SRSs (as used in Matchbox 2007).
- \Rightarrow for TRSs we can “only” prove top-termination.

Arctic interpretations

- $\mathbb{A}_{\mathbb{N}} \equiv \{-\infty\} \cup \mathbb{N}$.
- We say that $a \in \mathbb{A}_{\mathbb{N}}$ is *finite* iff $a \neq -\infty$.
- Interpretation domain: $\mathbb{A}_{\mathbb{N}}^d$, for some fixed d .
- Now we compute in the $\langle \mathbb{A}_{\mathbb{N}}, \max, + \rangle$ semi-ring.
- $a \gg b \equiv a > b \vee (a = b = -\infty)$.
- $\vec{u} \geq \vec{v}$ iff $\forall_i \vec{u}_i \geq \vec{v}_i$.
- $\vec{u} > \vec{v}$ iff $\forall_i \vec{u}_i \gg \vec{v}_i$.
- **Problem:** arctic addition is not strictly monotonic in single arguments, ie. $5 > 3$ but $5 \oplus 6 = 6 \not> 6 = 3 \oplus 6$. We cannot get strict monotonicity for symbols of arity > 1 .
- \Rightarrow full termination only for SRSs (as used in Matchbox 2007).
- \Rightarrow for TRSs we can “only” prove top-termination.
- **Problem:** well-foundedness of $>$ (as $-\infty \gg -\infty$)

Arctic interpretations

- $\mathbb{A}_{\mathbb{N}} \equiv \{-\infty\} \cup \mathbb{N}$.
- We say that $a \in \mathbb{A}_{\mathbb{N}}$ is *finite* iff $a \neq -\infty$.
- Interpretation domain: $\mathbb{N} \times \mathbb{A}_{\mathbb{N}}^{d-1}$, for some fixed d .
- Now we compute in the $\langle \mathbb{A}_{\mathbb{N}}, \max, + \rangle$ semi-ring.
- $a \gg b \equiv a > b \vee (a = b = -\infty)$.
- $\vec{u} \geq \vec{v}$ iff $\forall_i \vec{u}_i \geq \vec{v}_i$.
- $\vec{u} > \vec{v}$ iff $\forall_i \vec{u}_i \gg \vec{v}_i$.
- **Problem:** arctic addition is not strictly monotonic in single arguments, ie. $5 > 3$ but $5 \oplus 6 = 6 \not> 6 = 3 \oplus 6$. We cannot get strict monotonicity for symbols of arity > 1 .
- \Rightarrow full termination only for SRSs (as used in Matchbox 2007).
- \Rightarrow for TRSs we can “only” prove top-termination.
- **Problem:** well-foundedness of $>$ (as $-\infty \gg -\infty$)

Arctic interpretations

- $\mathbb{A}_{\mathbb{N}} \equiv \{-\infty\} \cup \mathbb{N}$.
- We say that $a \in \mathbb{A}_{\mathbb{N}}$ is *finite* iff $a \neq -\infty$.
- Interpretation domain: $\mathbb{N} \times \mathbb{A}_{\mathbb{N}}^{d-1}$, for some fixed d .
- Now we compute in the $\langle \mathbb{A}_{\mathbb{N}}, \max, + \rangle$ semi-ring.
- $a \gg b \equiv a > b \vee (a = b = -\infty)$.
- $\vec{u} \geq \vec{v}$ iff $\forall_i \vec{u}_i \geq \vec{v}_i$.
- $\vec{u} > \vec{v}$ iff $\forall_i \vec{u}_i \gg \vec{v}_i$.
- **Problem:** arctic addition is not strictly monotonic in single arguments, ie. $5 > 3$ but $5 \oplus 6 = 6 \not> 6 = 3 \oplus 6$. We cannot get strict monotonicity for symbols of arity > 1 .
- \Rightarrow full termination only for SRSs (as used in Matchbox 2007).
- \Rightarrow for TRSs we can “only” prove top-termination.
- **Problem:** well-foundedness of $>$ (as $-\infty \gg -\infty$)
- **Problem:** we need to ensure that we stay within the domain.

Arctic interpretations

- $\mathbb{A}_{\mathbb{N}} \equiv \{-\infty\} \cup \mathbb{N}$.
- We say that $a \in \mathbb{A}_{\mathbb{N}}$ is *finite* iff $a \neq -\infty$.
- Interpretation domain: $\mathbb{N} \times \mathbb{A}_{\mathbb{N}}^{d-1}$, for some fixed d .
- Now we compute in the $\langle \mathbb{A}_{\mathbb{N}}, \max, + \rangle$ semi-ring.
- $a \gg b \equiv a > b \vee (a = b = -\infty)$.
- $\vec{u} \geq \vec{v}$ iff $\forall_i \vec{u}_i \geq \vec{v}_i$.
- $\vec{u} > \vec{v}$ iff $\forall_i \vec{u}_i \gg \vec{v}_i$.
- **Problem:** arctic addition is not strictly monotonic in single arguments, ie. $5 > 3$ but $5 \oplus 6 = 6 \not> 6 = 3 \oplus 6$. We cannot get strict monotonicity for symbols of arity > 1 .
- \Rightarrow full termination only for SRSs (as used in Matchbox 2007).
- \Rightarrow for TRSs we can “only” prove top-termination.
- **Problem:** well-foundedness of $>$ (as $-\infty \gg -\infty$)
- **Problem:** we need to ensure that we stay within the domain.
- \Rightarrow for every interpretation $[f(x_1, \dots, x_n)] = F_1 x_1 + \dots F_n x_n + \vec{f}$ we require $\exists_i \text{finite}((F_i)_{1,1})$ or $\text{finite}(\vec{f}_1)$.

Example arctic proof

Example

$$\{cac \rightarrow \epsilon, aca \rightarrow a^4 \mid \epsilon \rightarrow c^4\}.$$

Example arctic proof

Example

$$\{\mathbf{c} \mathbf{a} \mathbf{c} \rightarrow \epsilon, \mathbf{a} \mathbf{c} \mathbf{a} \rightarrow \mathbf{a}^4 \mid \epsilon \rightarrow \mathbf{c}^4\}.$$

$$[\mathbf{a}](x) = \begin{pmatrix} 0 & 0 & -\infty \\ 0 & 0 & -\infty \\ 1 & 1 & 0 \end{pmatrix} x \oplus \begin{pmatrix} -\infty \\ -\infty \\ -\infty \end{pmatrix} \quad [\mathbf{c}](x) = \begin{pmatrix} 0 & -\infty & -\infty \\ -\infty & -\infty & 0 \\ -\infty & 0 & -\infty \end{pmatrix} x \oplus \begin{pmatrix} -\infty \\ -\infty \\ -\infty \end{pmatrix}$$

Example

$$\{\mathbf{c} \mathbf{a} \mathbf{c} \rightarrow \epsilon, \mathbf{a} \mathbf{c} \mathbf{a} \rightarrow \mathbf{a}^4 / \epsilon \rightarrow \mathbf{c}^4\}.$$

$$[\mathbf{a}](x) = \begin{pmatrix} 0 & 0 & -\infty \\ 0 & 0 & -\infty \\ 1 & 1 & 0 \end{pmatrix} x \oplus \begin{pmatrix} -\infty \\ -\infty \\ -\infty \end{pmatrix} \quad [\mathbf{c}](x) = \begin{pmatrix} 0 & -\infty & -\infty \\ -\infty & -\infty & 0 \\ -\infty & 0 & -\infty \end{pmatrix} x \oplus \begin{pmatrix} -\infty \\ -\infty \\ -\infty \end{pmatrix}$$

- $[\mathbf{c}]$ is a permutation (it swaps the second and third component), so $[\mathbf{c}]^2 = [\mathbf{c}]^4 = [\epsilon]$.

Example

$$\{\mathbf{c} \mathbf{a} \mathbf{c} \rightarrow \epsilon, \mathbf{a} \mathbf{c} \mathbf{a} \rightarrow \mathbf{a}^4 \mid \epsilon \rightarrow \mathbf{c}^4\}.$$

$$[\mathbf{a}](x) = \begin{pmatrix} 0 & 0 & -\infty \\ 0 & 0 & -\infty \\ 1 & 1 & 0 \end{pmatrix} x \oplus \begin{pmatrix} -\infty \\ -\infty \\ -\infty \end{pmatrix} \quad [\mathbf{c}](x) = \begin{pmatrix} 0 & -\infty & -\infty \\ -\infty & -\infty & 0 \\ -\infty & 0 & -\infty \end{pmatrix} x \oplus \begin{pmatrix} -\infty \\ -\infty \\ -\infty \end{pmatrix}$$

- $[\mathbf{c}]$ is a permutation (it swaps the second and third component), so $[\mathbf{c}]^2 = [\mathbf{c}]^4 = [\epsilon]$.
- $[\mathbf{a}]$ is idempotent, so $[\mathbf{a}] = [\mathbf{a}^4]$.

Example arctic proof

Example

$$\{\mathbf{c} \mathbf{a} \mathbf{c} \rightarrow \epsilon, \mathbf{a} \mathbf{c} \mathbf{a} \rightarrow \mathbf{a}^4 / \epsilon \rightarrow \mathbf{c}^4\}.$$

$$[\mathbf{a}](x) = \begin{pmatrix} 0 & 0 & -\infty \\ 0 & 0 & -\infty \\ 1 & 1 & 0 \end{pmatrix} x \oplus \begin{pmatrix} -\infty \\ -\infty \\ -\infty \end{pmatrix} \quad [\mathbf{c}](x) = \begin{pmatrix} 0 & -\infty & -\infty \\ -\infty & -\infty & 0 \\ -\infty & 0 & -\infty \end{pmatrix} x \oplus \begin{pmatrix} -\infty \\ -\infty \\ -\infty \end{pmatrix}$$

- $[\mathbf{c}]$ is a permutation (it swaps the second and third component), so $[\mathbf{c}]^2 = [\mathbf{c}]^4 = [\epsilon]$.
- $[\mathbf{a}]$ is idempotent, so $[\mathbf{a}] = [\mathbf{a}^4]$.

$$[\mathbf{c} \mathbf{a} \mathbf{c}](x) = \begin{pmatrix} 0 & -\infty & 0 \\ 1 & 0 & 1 \\ 0 & -\infty & 0 \end{pmatrix} x \geq \begin{pmatrix} -\infty & 0 & 0 \\ 0 & -\infty & 0 \\ 0 & 0 & -\infty \end{pmatrix} x = [\epsilon](x)$$

$$[\mathbf{a} \mathbf{c} \mathbf{a}](x) = \begin{pmatrix} 1 & 1 & 0 \\ 1 & 1 & 0 \\ 2 & 2 & 1 \end{pmatrix} x > \begin{pmatrix} 0 & 0 & -\infty \\ 0 & 0 & -\infty \\ 1 & 1 & 0 \end{pmatrix} x = [\mathbf{a}^4](x)$$

$$[\epsilon](x) = \begin{pmatrix} -\infty & 0 & 0 \\ 0 & -\infty & 0 \\ 0 & 0 & -\infty \end{pmatrix} x = \begin{pmatrix} -\infty & 0 & 0 \\ 0 & -\infty & 0 \\ 0 & 0 & -\infty \end{pmatrix} x = [\mathbf{c}^4](x)$$

Outline

- 1 Introduction
- 2 Monotone Algebras
- 3 Polynomial and Matrix Interpretations
- 4 Arctic Interpretations
- 5 Arctic Below Zero Interpretations**
- 6 Certification
- 7 Evaluation
- 8 Conclusions

- $\mathbb{A}_{\mathbb{Z}} \equiv \{-\infty\} \cup \mathbb{Z}.$

Arctic below zero interpretations

- $\mathbb{A}_{\mathbb{Z}} \equiv \{-\infty\} \cup \mathbb{Z}$.
- Interpretation domain: $\mathbb{N} \times \mathbb{A}_{\mathbb{Z}}^{d-1}$, for some fixed d .

Arctic below zero interpretations

- $\mathbb{A}_{\mathbb{Z}} \equiv \{-\infty\} \cup \mathbb{Z}$.
- Interpretation domain: $\mathbb{N} \times \mathbb{A}_{\mathbb{Z}}^{d-1}$, for some fixed d .
- \Rightarrow we restrict first component to \mathbb{N} to get well-foundedness.

Arctic below zero interpretations

- $\mathbb{A}_{\mathbb{Z}} \equiv \{-\infty\} \cup \mathbb{Z}$.
- Interpretation domain: $\mathbb{N} \times \mathbb{A}_{\mathbb{Z}}^{d-1}$, for some fixed d .
- \Rightarrow we restrict first component to \mathbb{N} to get well-foundedness.
- **Semi-ring structure:** $\langle \mathbb{A}_{\mathbb{Z}}, \text{max}, + \rangle$.

Arctic below zero interpretations

- $\mathbb{A}_{\mathbb{Z}} \equiv \{-\infty\} \cup \mathbb{Z}$.
- Interpretation domain: $\mathbb{N} \times \mathbb{A}_{\mathbb{Z}}^{d-1}$, for some fixed d .
- \Rightarrow we restrict first component to \mathbb{N} to get well-foundedness.
- Semi-ring structure: $\langle \mathbb{A}_{\mathbb{Z}}, \max, + \rangle$.
- $\geq, >$ as before.

Arctic below zero interpretations

- $\mathbb{A}_{\mathbb{Z}} \equiv \{-\infty\} \cup \mathbb{Z}$.
- Interpretation domain: $\mathbb{N} \times \mathbb{A}_{\mathbb{Z}}^{d-1}$, for some fixed d .
- \Rightarrow we restrict first component to \mathbb{N} to get well-foundedness.
- Semi-ring structure: $\langle \mathbb{A}_{\mathbb{Z}}, \max, + \rangle$.
- $\geq, >$ as before.
- **Problem:** we need to ensure that we stay within the domain.

Arctic below zero interpretations

- $\mathbb{A}_{\mathbb{Z}} \equiv \{-\infty\} \cup \mathbb{Z}$.
- Interpretation domain: $\mathbb{N} \times \mathbb{A}_{\mathbb{Z}}^{d-1}$, for some fixed d .
- \Rightarrow we restrict first component to \mathbb{N} to get well-foundedness.
- Semi-ring structure: $\langle \mathbb{A}_{\mathbb{Z}}, \max, + \rangle$.
- $\geq, >$ as before.
- **Problem:** we need to ensure that we stay within the domain.
- \Rightarrow for every interpretation $[f(x_1, \dots, x_n)] = F_1 x_1 + \dots F_n x_n + \vec{f}$ we require $\vec{f}_1 \geq 0$.

Example arctic below zero proof

Example

```
while x > y do x := x - 1;
```

Example arctic below zero proof

Example

`while x > y do x := x - 1;`

$\text{cond}(\text{true}, x, y) \rightarrow \text{cond}(\text{gr}(x, y), p(x), y),$	$\text{gr}(s(x), s(y)) \rightarrow \text{gr}(x, y),$
$\text{gr}(0, x) \rightarrow \text{false},$	$\text{gr}(s(x), 0) \rightarrow \text{true},$
$p(0) \rightarrow 0,$	$p(s(x)) \rightarrow x$

Example arctic below zero proof

Example

`while x > y do x := x - 1;`

$\text{cond}(\text{true}, x, y) \rightarrow \text{cond}(\text{gr}(x, y), p(x), y), \quad \text{gr}(s(x), s(y)) \rightarrow \text{gr}(x, y),$

$\text{gr}(0, x) \rightarrow \text{false},$

$\text{gr}(s(x), 0) \rightarrow \text{true},$

$p(0) \rightarrow 0,$

$p(s(x)) \rightarrow x$

$\text{cond}^\#(\text{true}, x, y) \rightarrow \text{cond}^\#(\text{gr}(x, y), p(x), y)$

Example arctic below zero proof

Example

while $x > y$ do $x := x - 1$;

$$\begin{aligned} \text{cond}(\text{true}, x, y) &\rightarrow \text{cond}(\text{gr}(x, y), p(x), y), & \text{gr}(s(x), s(y)) &\rightarrow \text{gr}(x, y), \\ \text{gr}(0, x) &\rightarrow \text{false}, & \text{gr}(s(x), 0) &\rightarrow \text{true}, \\ p(0) &\rightarrow 0, & p(s(x)) &\rightarrow x \end{aligned}$$

$$\text{cond}^\#(\text{true}, x, y) \rightarrow \text{cond}^\#(\text{gr}(x, y), p(x), y)$$

$$\begin{aligned} [\text{cond}^\#(x, y, z)] &= (0)x \oplus (0)y \oplus (-\infty)z \oplus (0), & [0] &= (0), \\ [\text{cond}(x, y, z)] &= (0)x \oplus (2)y \oplus (-\infty)z \oplus (0), & [\text{false}] &= (0), \\ [\text{gr}(x, y)] &= (-1)x \oplus (-\infty)y \oplus (0), & [\text{true}] &= (2), \\ [p(x)] &= (-1)x \oplus (0), & [s(x)] &= (2)x \oplus (3). \end{aligned}$$

Example arctic below zero proof

Example

while $x > y$ do $x := x - 1$;

$$\begin{aligned} \text{cond}(\text{true}, x, y) &\rightarrow \text{cond}(\text{gr}(x, y), p(x), y), & \text{gr}(s(x), s(y)) &\rightarrow \text{gr}(x, y), \\ \text{gr}(0, x) &\rightarrow \text{false}, & \text{gr}(s(x), 0) &\rightarrow \text{true}, \\ p(0) &\rightarrow 0, & p(s(x)) &\rightarrow x \end{aligned}$$

$$\text{cond}^\#(\text{true}, x, y) \rightarrow \text{cond}^\#(\text{gr}(x, y), p(x), y)$$

$$\begin{aligned} [\text{cond}^\#(x, y, z)] &= (0)x \oplus (0)y \oplus (-\infty)z \oplus (0), & [0] &= (0), \\ [\text{cond}(x, y, z)] &= (0)x \oplus (2)y \oplus (-\infty)z \oplus (0), & [\text{false}] &= (0), \\ [\text{gr}(x, y)] &= (-1)x \oplus (-\infty)y \oplus (0), & [\text{true}] &= (2), \\ [p(x)] &= (-1)x \oplus (0), & [s(x)] &= (2)x \oplus (3). \end{aligned}$$

$$[\text{cond}^\#(\text{true}, x, y)] = (0)x \oplus (-\infty)y \oplus (2)$$

$$[\text{cond}^\#(\text{gr}(x, y), p(x), y)] = (-1)x \oplus (-\infty)y \oplus (0)$$

Outline

- 1 Introduction
- 2 Monotone Algebras
- 3 Polynomial and Matrix Interpretations
- 4 Arctic Interpretations
- 5 Arctic Below Zero Interpretations
- 6 Certification**
- 7 Evaluation
- 8 Conclusions

CoLoR: certification of termination proofs

Certification of termination: formal verification (using a theorem prover/checker) of termination proofs produced by termination provers.

CoLoR: certification of termination proofs

Certification of termination: formal verification (using a theorem prover/checker) of termination proofs produced by termination provers.

CoLoR: Coq Library on Rewriting and Termination.

`http://color.loria.fr`

Goal: certification of termination proofs produced by various termination provers.

CoLoR: certification of termination proofs

Certification of termination: formal verification (using a theorem prover/checker) of termination proofs produced by termination provers.

CoLoR: Coq Library on Rewriting and Termination.

`http://color.loria.fr`

Goal: certification of termination proofs produced by various termination provers.

- TPG: common format for termination proofs (independent of termination tools and the certification back-end).

CoLoR: certification of termination proofs

Certification of termination: formal verification (using a theorem prover/checker) of termination proofs produced by termination provers.

CoLoR: Coq Library on Rewriting and Termination.

`http://color.loria.fr`

Goal: certification of termination proofs produced by various termination provers.

- **TPG**: common format for termination proofs (independent of termination tools and the certification back-end).
- **Tools output proofs in TPG format.**

CoLoR: certification of termination proofs

Certification of termination: formal verification (using a theorem prover/checker) of termination proofs produced by termination provers.

CoLoR: Coq Library on Rewriting and Termination.

`http://color.loria.fr`

Goal: certification of termination proofs produced by various termination provers.

- **TPG**: common format for termination proofs (independent of termination tools and the certification back-end).
- Tools output proofs in TPG format.
- **CoLoR**: a Coq library of results on termination.

CoLoR: certification of termination proofs

Certification of termination: formal verification (using a theorem prover/checker) of termination proofs produced by termination provers.


CoLoR: Coq Library on Rewriting and Termination.

`http://color.loria.fr`



Goal: certification of termination proofs produced by various termination provers.

- **TPG**: common format for termination proofs (independent of termination tools and the certification back-end).
- Tools output proofs in TPG format.
- **CoLoR**: a Coq library of results on termination.
- **Rainbow**: a tool for translation from proofs in TPG format to Coq proofs, using results from CoLoR.


CoLoR's architecture overview



CoLoR's architecture overview



CoLoR's architecture overview



Outline

- 1 Introduction
- 2 Monotone Algebras
- 3 Polynomial and Matrix Interpretations
- 4 Arctic Interpretations
- 5 Arctic Below Zero Interpretations
- 6 Certification
- 7 Evaluation**
- 8 Conclusions

Termination competition:

- in 2007 introduced a new category of certified termination,

Termination competition:

- in 2007 introduced a new category of certified termination,
- TPA+CoLoR was the winner.

Evaluation

Termination competition:

- in 2007 introduced a new category of certified termination,
- TPA+CoLoR was the winner.

The approach of arctic interpretations was:

- implemented in Matchbox using a propositional encoding into SAT.

Termination competition:

- in 2007 introduced a new category of certified termination,
- TPA+CoLoR was the winner.

The approach of arctic interpretations was:

- implemented in Matchbox using a propositional encoding into SAT.
- formalized within the CoLoR project allowing certification of arctic proofs.

Evaluation

Termination competition:

- in 2007 introduced a new category of certified termination,
- TPA+CoLoR was the winner.

The approach of arctic interpretations was:

- **implemented** in Matchbox using a propositional encoding into SAT.
- **formalized** within the CoLoR project allowing certification of arctic proofs.

problem set	time	s	sa	sz	saz	2007 winner
975 TRS	1 min	361	376	388	389	TPA: 354
	10 min	365	381	393	394	
517 SRS	1 min	178	312	298	320	Matchbox: 337
	10 min	185	349	323	354	

Outline

- 1 Introduction
- 2 Monotone Algebras
- 3 Polynomial and Matrix Interpretations
- 4 Arctic Interpretations
- 5 Arctic Below Zero Interpretations
- 6 Certification
- 7 Evaluation
- 8 Conclusions**

Conclusions

- We presented an extension of matrix interpretations method by replacing the usual semi-ring structure with the arctic semi-ring.

Conclusions

- We presented an extension of matrix interpretations method by replacing the usual semi-ring structure with the arctic semi-ring.
- The method can prove full termination for SRSs and (relative) top termination for TRSs

Conclusions

- We presented an extension of matrix interpretations method by replacing the usual semi-ring structure with the arctic semi-ring.
- The method can prove full termination for SRSs and (relative) top termination for TRSs
- ... and imposes linear derivational complexity (without DP).

Conclusions

- We presented an extension of matrix interpretations method by replacing the usual semi-ring structure with the arctic semi-ring.
- The method can prove full termination for SRSs and (relative) top termination for TRSs
- ... and imposes linear derivational complexity (without DP).
- (The admissibility problem of an arctic interpretation corresponds to a reachability problem for weighted tree automata).

Conclusions

- We presented an extension of matrix interpretations method by replacing the usual semi-ring structure with the arctic semi-ring.
- The method can prove full termination for SRSs and (relative) top termination for TRSs
- ... and imposes linear derivational complexity (without DP).
- (The admissibility problem of an arctic interpretation corresponds to a reachability problem for weighted tree automata).
- We extended this from naturals to integers, resulting in arctic below zero interpretations.

Conclusions

- We presented an extension of matrix interpretations method by replacing the usual semi-ring structure with the arctic semi-ring.
- The method can prove full termination for SRSs and (relative) top termination for TRSs
- ... and imposes linear derivational complexity (without DP).
- (The admissibility problem of an arctic interpretation corresponds to a reachability problem for weighted tree automata).
- We extended this from naturals to integers, resulting in arctic below zero interpretations.
- The whole method has been formalized in Coq within the CoLoR project.

Conclusions

- We presented an extension of matrix interpretations method by replacing the usual semi-ring structure with the arctic semi-ring.
- The method can prove full termination for SRSs and (relative) top termination for TRSs
- ... and imposes linear derivational complexity (without DP).
- (The admissibility problem of an arctic interpretation corresponds to a reachability problem for weighted tree automata).
- We extended this from naturals to integers, resulting in arctic below zero interpretations.
- The whole method has been formalized in Coq within the CoLoR project.
- It has also been implemented in Matchbox, by transforming the constraints to propositional satisfiability problem and running Minisat.



Thank you for your attention.