

1. GENERAL DISCUSSION

Let Σ be any smooth, compact surface (potentially with boundary), and let Δ be its Laplace operators. Let $(f_a \in L^2(\Sigma))_{a \in I}$ be a complete, orthonormal set of eigenfunctions (thus I is countable infinite), that is $\Delta f_a = \lambda_a f_a$. Without any loss of generality, assume that $0 \in I$ and f_0 is a constant. Then write

$$\varrho_t = \sum_{a \in I} \sum_{n \in \mathbb{N}} R_{n,a} t^n f_a.$$

Let

$$\varphi_{a,b,c} := \int_{\Sigma} f_a f_b f_c dA.$$

If ϱ_t solves the Keller–Segel equation, then and $a \neq 0$, then

$$R_{n+1,a} = \frac{R_{0,0} - \lambda_a}{n+1} R_{n,a} + \sum_{b,c \in I - \{0\}} \sum_{m=0}^n \frac{\lambda_a - \lambda_b + \lambda_c}{(n+1)\lambda_c} \varphi_{a,b,c} R_{b,m} R_{c,n-m}.$$

Remark 1.1. For all $\Lambda \in \mathbb{R}_+$, let

$$n(\Lambda) := |\{ a \in I \mid \lambda_a \leq \Lambda \}|.$$

Weyl's law, that states that

$$\lim_{\Lambda \rightarrow \infty} \frac{\Lambda}{n(\Lambda)} = \frac{2}{\text{Area}(\Sigma, g)}.$$

2. CONVERGENCE OF THE ITERATIVE METHOD ON THE STANDARD TORUS

Let us write

$$\varrho(x, t) = \sum_{\underline{k} \in \mathbb{Z}^2} \sum_{n \in \mathbb{N}} R_{n,\underline{k}} t^n \exp(2\pi i \underline{k} \cdot \underline{x}).$$

If ϱ solves the Keller–Segel equation, then and $\underline{k} \neq \underline{0}$, then

$$R_{n+1,\underline{k}} = \frac{R_{0,0} - 4\pi^2 |\underline{k}|^2}{n+1} R_{n,\underline{k}} + \sum_{\underline{l} \in \mathbb{Z}^2 - \{0,\underline{k}\}} \frac{\underline{k} \cdot \underline{l}}{|\underline{l}|^2} \sum_{m=0}^n \frac{1}{n+1} R_{m,\underline{l}} R_{n-m,\underline{k}-\underline{l}}. \quad (2.1)$$

Finally let

$$\forall \underline{k} \in \mathbb{Z}^2: \quad S_{\underline{k}}(t) := R_{\underline{k}}(t) \exp(4\pi^2 |\underline{k}|^2 t),$$

and write

$$S_{\underline{k}}(t) = \sum_{n \in \mathbb{N}} S_{n,\underline{k}} t^n.$$

Then, in terms of the $S_{n,\underline{k}}$ coefficients, equation (2.1) becomes

$$S_{n+1,\underline{k}} = \frac{1}{n+1} \sum_{\underline{l} \in \mathbb{Z}^2 - \{0,\underline{k}\}} \frac{\underline{k} \cdot \underline{l}}{|\underline{l}|^2} \sum_{a=0}^n \sum_{b=0}^{n-a} S_{a,\underline{l}} S_{b,\underline{k}-\underline{l}} \frac{(8\pi^2 (\underline{k}-\underline{l}) \cdot \underline{l})^{n-a-b}}{(n-a-b)!}. \quad (2.2)$$

If that the coefficients in equation (2.1) exist after the n^{th} step, then let

$$Z_n := \{ \underline{k} \in \mathbb{Z}^2 \mid S_{n,\underline{k}} \neq 0 \} \subseteq \mathbb{Z}^2.$$

Lemma 2.1. Let us assume that Z_0 is finite and symmetric, that is $-Z_0 = Z_0$, and let

$$d := \max(\{ |\underline{k}| \mid \underline{k} \in Z_0 \}) > 0.$$

Then for all $n \in \mathbb{N}_+$, the coefficients in equation (2.1) exist and Z_n is finite and symmetric, moreover

$$Z_n \subseteq Z_{n-1} + Z_{n-1},$$

and

$$\max(\{ |\underline{k}| \mid \underline{k} \in Z_n \}) = \max(\{ |\underline{k}| \mid S_{n,\underline{k}} \neq 0 \}) \leq 2^n d.$$

Proof. *Insert Adam's proof here.* □

For all $\underline{k} \in \mathbb{Z}^2 - \{\underline{0}\}$ let

$$N(\underline{k}) := \min(\{n \in \mathbb{N} \mid \underline{k} \in Z_n\}).$$

Lemma 2.2. *Under the assumptions above, we have that*

$$\forall \underline{k} \in \mathbb{Z}^2 - \{\underline{0}\} : \forall n \in \mathbb{N} : |S_{n,\underline{k}}| < \epsilon^n. \quad (2.3)$$

Proof. By Lemma 2.1, we can rewrite equation (2.2) as

$$S_{n+1,\underline{k}} = \frac{1}{n+1} \sum_{x=0}^n \sum_{\underline{l}: N(\underline{l})=x} \frac{\underline{k} \cdot \underline{l}}{|\underline{l}|^2} \sum_{a=x}^n \sum_{b=0}^{n-a} S_{a,\underline{l}} S_{b,\underline{k}-\underline{l}} \frac{(8\pi^2(\underline{k}-\underline{l}) \cdot \underline{l})^{n-a-b}}{(n-a-b)!}. \quad (2.4)$$

The inequality (2.3) is true for $n = 0$, so we can argue by induction.

Let us assume now that we have proved inequality (2.3) for every nonnegative integer up to $n \in \mathbb{N}$. Without any loss of generality, we can assume that $n > N(\underline{k})$, then inequality (2.3) holds. Now using equation (2.4) we have:

$$\begin{aligned} |S_{n+1,\underline{k}}| &\leq \frac{1}{n+1} \sum_{x=0}^n \sum_{\underline{l}: N(\underline{l})=x} \sum_{a=x}^n \sum_{b=0}^{n-a} \epsilon^{a+b} \frac{|8\pi^2(\underline{k}-\underline{l}) \cdot \underline{l}|^{n-a-b}}{(n-a-b)!} \frac{|\underline{k} \cdot \underline{l}|}{|\underline{l}|^2} \\ &\leq \frac{\epsilon^n}{n+1} \sum_{x=0}^n \sum_{\underline{l}: N(\underline{l})=x} \sum_{a=0}^{n-x} a \frac{|8\pi^2(\underline{k}-\underline{l}) \cdot \underline{l}/\epsilon|^{n-x-a}}{(n-x-a)!} \frac{|\underline{k} \cdot \underline{l}|}{|\underline{l}|^2} \\ &\leq \frac{\epsilon^n}{n+1} \sum_{x=0}^n \sum_{\underline{l}: N(\underline{l})=x} \sum_{a=0}^{n-x} (n-x-a) \frac{|8\pi^2(\underline{k}-\underline{l}) \cdot \underline{l}/\epsilon|^a}{a!} \frac{|\underline{k} \cdot \underline{l}|}{|\underline{l}|^2} \\ &\leq \frac{\epsilon^n}{n+1} \sum_{x=0}^n \sum_{a=0}^{n-x} (n-x-a) \sum_{\underline{l}: N(\underline{l})=x} \frac{|8\pi^2(\underline{k}-\underline{l}) \cdot \underline{l}/\epsilon|^a}{a!} \frac{|\underline{k} \cdot \underline{l}|}{|\underline{l}|^2} \end{aligned}$$

Thus

$$\begin{aligned} \left| \sum_{m=0}^n \sum_{\substack{\underline{l} \in Z_m \\ \underline{k}-\underline{l} \in Z_{n-m}}} \frac{1}{n+1} R_{m,\underline{l}} R_{n-m,\underline{k}-\underline{l}} \right| &\leq \sum_{m=0}^n \frac{C^{n+2}}{(n+1)!} \exp\left(-\frac{|\underline{k}|}{C}\right) \binom{n}{m} 2\pi 4^m 4^{n \ln(n)} \\ &\leq \frac{C^{n+2}}{(n+1)!} \exp\left(-\frac{|\underline{k}|}{C}\right) 4^{n \ln(n)} \sum_{m=0}^n \binom{n}{m} 2\pi 4^m \\ &= \frac{C^{n+2}}{(n+1)!} \exp\left(-\frac{|\underline{k}|}{C}\right) 4^{n \ln(n)} 2\pi \frac{5^{n+1} - 1}{5 - 1}. \end{aligned}$$

Inserting this to the outer sum, we get

$$\left| \sum_{\underline{l} \in \mathbb{Z}^2 - \{\underline{0}, \underline{k}\}} \frac{\underline{k} \cdot \underline{l}}{|\underline{l}|^2} \sum_{m=0}^n \frac{1}{n+1} R_{m,\underline{l}} R_{n-m,\underline{k}-\underline{l}} \right| \leq \frac{C^{n+2}}{(n+1)!} \exp\left(-\frac{|\underline{k}|^2}{C}\right) \sum_{\underline{l} \in \mathbb{Z}^2 - \{\underline{0}\}} \frac{|\underline{k} \cdot \underline{l}|}{|\underline{l}|^2} \left(|\underline{l}|^2 + |\underline{k}-\underline{l}|^2\right)^n \exp\left(\frac{2(\underline{k}-\underline{l}) \cdot \underline{l}}{C}\right)$$

□