

Reparametrisation Tricks  
or  
How to Differentiate Through Samples from  
Probability Distributions

Adam Kosiorek

June 8, 2017

# Table of Contents


## A Few Examples

Why do we need this?

Continuous Random Variables: One-Liners

Discrete Random Variables: REINFORCE

# Playing Starcraft



$$\theta_t = f_{\phi}(\mathbf{x}_{1:t})$$

compute parameters

$$a_t \sim p(a \mid \theta_t)$$

sample action from a distribution


$$R = R(\mathbf{a}_{1:T}, \mathbf{x}_{1:T})$$

compute reward based on actions and states


## Alex Kendall's work on uncertainty



Input image  $I$



Depth estimate  $d$






Uncertainty  $\sigma^2$

$$\begin{aligned}\mu, \sigma^2 &= f_{\phi}(I) \\ d &\sim \mathcal{N}(d \mid \mu, \sigma^2)\end{aligned}$$

# Stochastic Attractors of Chaotic Systems

Neil Dhir, Adam Kosiorek, Michael Osborne, Ingmar Posner



Dimensionality of the latent space?

$$d_E \sim q_\phi(d_E \mid \mathbf{x}_{1:T})$$

$$f_{d_E} : \mathbb{R}^N \rightarrow \mathbb{R}^{d_E}$$

$$f_{d_E} : \mathbf{x}_{1:T} \mapsto \mathbf{z}_{1:T}$$

# Table of Contents


A Few Examples

Why do we need this?

Continuous Random Variables: One-Liners

Discrete Random Variables: REINFORCE

# General Computation Graph



# Backprop Equations

Let  $\delta^{(l)} = \frac{\partial \mathcal{L}(\cdot)}{\partial \mathbf{z}^{(l)}}$ , then backprop is given by:


$$\delta^{(L)} = \nabla_{\mathbf{a}^{(L)}} \mathcal{L}(\mathbf{a}^{(L)}, \mathbf{y}) \odot \sigma'(\mathbf{z}^{(L)}), \quad (1)$$

$$\delta^{(l)} = \left( \left( \mathbf{W}^{(l+1)} \right)^T \delta^{(l+1)} \right) \odot \sigma'(\mathbf{z}^{(l)}), \quad (2)$$

$$\frac{\partial \mathcal{L}(\cdot)}{\partial \mathbf{W}^{(l)}} = \delta^{(l)} \left( \mathbf{a}^{(l-1)} \right)^T. \quad (3)$$



# Stochastic Computation Graph



## Forward pass

$$\mathbf{z}^{(0)} = \mathbf{W}^{(0)} \mathbf{x}$$

## Forward pass

$$\mathbf{z}^{(0)} = \mathbf{W}^{(0)} \mathbf{x}$$

$$\mathbf{a}^{(0)} = \sigma \left( \mathbf{z}^{(0)} \right)$$

## Forward pass

$$\mathbf{z}^{(0)} = \mathbf{W}^{(0)} \mathbf{x}$$

$$\mathbf{a}^{(0)} = \sigma \left( \mathbf{z}^{(0)} \right)$$

$$\mathbf{z}^{(1)} = \mathbf{W}^{(1)} \mathbf{a}^{(0)}$$

## Forward pass

$$\mathbf{z}^{(0)} = \mathbf{W}^{(0)} \mathbf{x}$$

$$\mathbf{a}^{(0)} = \sigma \left( \mathbf{z}^{(0)} \right)$$

$$\mathbf{z}^{(1)} = \mathbf{W}^{(1)} \mathbf{a}^{(0)}$$

$$\theta = \sigma \left( \mathbf{z}^{(1)} \right)$$

## Forward pass

$$\mathbf{z}^{(0)} = \mathbf{W}^{(0)} \mathbf{x}$$

$$\mathbf{a}^{(0)} = \sigma \left( \mathbf{z}^{(0)} \right)$$

$$\mathbf{z}^{(1)} = \mathbf{W}^{(1)} \mathbf{a}^{(0)}$$

$$\theta = \sigma \left( \mathbf{z}^{(1)} \right)$$

$$\mathbf{a}^{(1)} \sim p(\mathbf{a} \mid \theta)$$

## Forward pass

$$\mathbf{z}^{(0)} = \mathbf{W}^{(0)} \mathbf{x}$$

$$\mathbf{a}^{(0)} = \sigma \left( \mathbf{z}^{(0)} \right)$$

$$\mathbf{z}^{(1)} = \mathbf{W}^{(1)} \mathbf{a}^{(0)}$$

$$\theta = \sigma \left( \mathbf{z}^{(1)} \right)$$

$$\mathbf{a}^{(1)} \sim p(\mathbf{a} \mid \theta)$$

$$\mathbf{z}^{(2)} = \mathbf{W}^{(2)} \mathbf{a}^{(1)}$$

## Forward pass

$$\mathbf{z}^{(0)} = \mathbf{W}^{(0)} \mathbf{x}$$

$$\mathbf{a}^{(0)} = \sigma \left( \mathbf{z}^{(0)} \right)$$

$$\mathbf{z}^{(1)} = \mathbf{W}^{(1)} \mathbf{a}^{(0)}$$

$$\theta = \sigma \left( \mathbf{z}^{(1)} \right)$$

$$\mathbf{a}^{(1)} \sim p(\mathbf{a} \mid \theta)$$

$$\mathbf{z}^{(2)} = \mathbf{W}^{(2)} \mathbf{a}^{(1)}$$

$$\mathbf{a}^{(2)} = \mathbf{z}^{(2)}$$



## Forward pass

$$\mathbf{z}^{(0)} = \mathbf{W}^{(0)} \mathbf{x}$$

$$\mathbf{a}^{(0)} = \sigma \left( \mathbf{z}^{(0)} \right)$$

$$\mathbf{z}^{(1)} = \mathbf{W}^{(1)} \mathbf{a}^{(0)}$$

$$\theta = \sigma \left( \mathbf{z}^{(1)} \right)$$

$$\mathbf{a}^{(1)} \sim p(\mathbf{a} \mid \theta)$$

$$\mathbf{z}^{(2)} = \mathbf{W}^{(2)} \mathbf{a}^{(1)}$$

$$\mathbf{a}^{(2)} = \mathbf{z}^{(2)}$$

$$\mathcal{L}(\cdot) = \frac{1}{2} \left( \mathbf{y} - \mathbf{a}^{(2)} \right)^2$$

## Backward pass

$$\mathbf{z}^{(0)} = \mathbf{W}^{(0)} \mathbf{x}$$

$$\mathbf{a}^{(0)} = \sigma \left( \mathbf{z}^{(0)} \right)$$

$$\mathbf{z}^{(1)} = \mathbf{W}^{(1)} \mathbf{a}^{(1)}$$

$$\theta = \sigma \left( \mathbf{z}^{(1)} \right)$$

$$\mathbf{a}^{(1)} \sim p(\mathbf{a} \mid \theta)$$

$$\mathbf{z}^{(2)} = \mathbf{W}^{(2)} \mathbf{a}^{(1)}$$

$$\mathbf{a}^{(2)} = \mathbf{z}^{(2)}$$

$$\delta^{(2)} = (\mathbf{y} - \mathbf{a}^{(2)})$$

$$\mathcal{L}(\cdot) = \frac{1}{2} \left( \mathbf{y} - \mathbf{a}^{(2)} \right)^2$$

## Backward pass

$$\mathbf{z}^{(0)} = \mathbf{W}^{(0)} \mathbf{x}$$

$$\mathbf{a}^{(0)} = \sigma(\mathbf{z}^{(0)})$$

$$\mathbf{z}^{(1)} = \mathbf{W}^{(1)} \mathbf{a}^{(1)}$$

$$\theta = \sigma(\mathbf{z}^{(1)})$$

$$\mathbf{a}^{(1)} \sim p(\mathbf{a} \mid \theta)$$

$$\mathbf{z}^{(2)} = \mathbf{W}^{(2)} \mathbf{a}^{(1)}$$

$$\mathbf{a}^{(2)} = \mathbf{z}^{(2)}$$

$$\delta^{(2)} = (\mathbf{y} - \mathbf{a}^{(2)})$$

$$\mathcal{L}(\cdot) = \frac{1}{2} \left( \mathbf{y} - \mathbf{a}^{(2)} \right)^2$$

## Backward pass

$$\mathbf{z}^{(0)} = \mathbf{W}^{(0)} \mathbf{x}$$

$$\mathbf{a}^{(0)} = \sigma(\mathbf{z}^{(0)})$$

$$\mathbf{z}^{(1)} = \mathbf{W}^{(1)} \mathbf{a}^{(1)} \quad \vdots$$

$$\theta = \sigma(\mathbf{z}^{(1)})$$

$$\mathbf{a}^{(1)} \sim p(\mathbf{a} \mid \theta) \quad \delta^{(1)} = \left( \left( \mathbf{W}^{(2)} \right)^T \delta^{(2)} \right) \odot \frac{\partial \mathbf{a}^{(1)}}{\partial \mathbf{z}^{(1)}}$$

$$\mathbf{z}^{(2)} = \mathbf{W}^{(2)} \mathbf{a}^{(1)}$$

$$\mathbf{a}^{(2)} = \mathbf{z}^{(2)} \quad \delta^{(2)} = (\mathbf{y} - \mathbf{a}^{(2)})$$

$$\mathcal{L}(\cdot) = \frac{1}{2} \left( \mathbf{y} - \mathbf{a}^{(2)} \right)^2$$

## Gradient of a sample?

$$\begin{aligned}\frac{\partial \mathbf{a}^{(1)}}{\partial \mathbf{z}^{(1)}} &= \frac{\mathbf{a}^{(1)}}{\partial p(\mathbf{a}^{(1)} \mid \theta)} \frac{\partial p(\mathbf{a}^{(1)} \mid \theta)}{\partial \mathbf{z}^{(1)}} \\ &= \frac{\mathbf{a}^{(1)}}{\partial p(\mathbf{a}^{(1)} \mid \theta)} \frac{\partial p(\mathbf{a}^{(1)} \mid \theta)}{\partial \theta} \frac{\partial \theta}{\partial \mathbf{z}^{(1)}}\end{aligned}\tag{4}$$

## Gradient of a sample?

$$\begin{aligned}\frac{\partial \mathbf{a}^{(1)}}{\partial \mathbf{z}^{(1)}} &= \frac{\mathbf{a}^{(1)}}{\partial p(\mathbf{a}^{(1)} | \theta)} \frac{\partial p(\mathbf{a}^{(1)} | \theta)}{\partial \mathbf{z}^{(1)}} \\ &= \frac{\mathbf{a}^{(1)}}{\partial p(\mathbf{a}^{(1)} | \theta)} \frac{\partial p(\mathbf{a}^{(1)} | \theta)}{\partial \theta} \frac{\partial \theta}{\partial \mathbf{z}^{(1)}}\end{aligned}\tag{4}$$

but...

$$\mathbf{a}^{(1)} \sim p(\mathbf{a} | \theta)\tag{5}$$

## Gradient of a sample?

$$\begin{aligned}\frac{\partial \mathbf{a}^{(1)}}{\partial \mathbf{z}^{(1)}} &= \frac{\mathbf{a}^{(1)}}{\partial p(\mathbf{a}^{(1)} | \theta)} \frac{\partial p(\mathbf{a}^{(1)} | \theta)}{\partial \mathbf{z}^{(1)}} \\ &= \frac{\mathbf{a}^{(1)}}{\partial p(\mathbf{a}^{(1)} | \theta)} \frac{\partial p(\mathbf{a}^{(1)} | \theta)}{\partial \theta} \frac{\partial \theta}{\partial \mathbf{z}^{(1)}}\end{aligned}\tag{4}$$

but...

$$\begin{aligned}\mathbf{a}^{(1)} &\sim p(\mathbf{a} | \theta) \\ \frac{\mathbf{a}^{(1)}}{\partial p(\mathbf{a}^{(1)} | \theta)} &= \nexists\end{aligned}\tag{5}$$

# Table of Contents

A Few Examples

Why do we need this?

Continuous Random Variables: One-Liners

Discrete Random Variables: REINFORCE



## (Continuous) Reparametrisation Trick

Let  $\mathbf{x} \sim p(\mathbf{x} \mid \theta)$  be a random variable. Perform change of variables such that

$$\begin{aligned}\epsilon &\sim p(\epsilon), \\ \mathbf{x} &= g(\epsilon, \theta)\end{aligned}\tag{6}$$

and treat  $\epsilon$  as if it was a constant.

## (Continuous) Reparametrisation Trick

Let  $\mathbf{x} \sim p(\mathbf{x} \mid \theta)$  be a random variable. Perform change of variables such that

$$\begin{aligned}\epsilon &\sim p(\epsilon), \\ \mathbf{x} &= g(\epsilon, \theta)\end{aligned}\tag{6}$$

and treat  $\epsilon$  as if it was a constant.

Example:

$$\epsilon \sim \mathcal{N}(0, 1)$$

$$x \sim \mathcal{N}\left(2, \frac{1}{4}\right) = \mathcal{N}(\mu, \sigma^2)$$

$$x = g(\epsilon, \mu, \sigma^2) = \mu + \sigma\epsilon = 2 + \frac{1}{2}\epsilon$$

## (Continuous) Reparametrisation Trick

Let  $\mathbf{x} \sim p(\mathbf{x} \mid \theta)$  be a random variable. Perform change of variables such that

$$\begin{aligned}\epsilon &\sim p(\epsilon), \\ \mathbf{x} &= g(\epsilon, \theta)\end{aligned}\tag{6}$$

and treat  $\epsilon$  as if it was a constant.

Example:

$$\epsilon \sim \mathcal{N}(0, 1)$$

$$x \sim \mathcal{N}\left(2, \frac{1}{4}\right) = \mathcal{N}(\mu, \sigma^2)$$

$$x = g(\epsilon, \mu, \sigma^2) = \mu + \sigma\epsilon = 2 + \frac{1}{2}\epsilon$$

$$\frac{dx}{d\mu} = 1 \qquad \frac{dx}{d\sigma} = \frac{1}{2}\epsilon \qquad \frac{dx}{d\epsilon} \stackrel{!}{=} 0$$

Why can we do this?

We are NOT computing  $\mathcal{L}(x)$ .

## Why can we do this?

We are NOT computing  $\mathcal{L}(x)$ .

We are computing  $\nabla_{\theta} \mathbb{E}_{p_{\theta}(x|\mathcal{D})}[\mathcal{L}(x)] = \nabla_{\theta} \int p_{\theta}(x|\mathcal{D}) \mathcal{L}(x) \mathrm{d}x$ .

## Why can we do this?

We are computing  $\nabla_{\theta} \mathbb{E}_{p_{\theta}(x|\mathcal{D})}[\mathcal{L}(x)] = \nabla_{\theta} \int p_{\theta}(x|\mathcal{D}) \mathcal{L}(x) dx$ .

We get the gradient by the change of variables:

$$p(x) = \left| \frac{d\epsilon}{dx} \right| p(\epsilon) \quad \text{Change of variables,}$$

$$\implies |p(x) dx| = |p(\epsilon) d\epsilon| \quad \text{Mass conservation.}$$

## Why can we do this?

We are computing  $\nabla_{\theta} \mathbb{E}_{p_{\theta}(x|\mathcal{D})}[\mathcal{L}(x)] = \nabla_{\theta} \int p_{\theta}(x|\mathcal{D}) \mathcal{L}(x) dx$ .

We get the gradient by the change of variables:

$$p(x) = \left| \frac{d\epsilon}{dx} \right| p(\epsilon) \quad \text{Change of variables,}$$

$$\implies |p(x) dx| = |p(\epsilon) d\epsilon| \quad \text{Mass conservation.}$$

$$\nabla_{\theta} \mathbb{E}_{p_{\theta}(x|\mathcal{D})}[\mathcal{L}(x)] = \nabla_{\theta} \int p_{\theta}(x|\mathcal{D}) \mathcal{L}(x) dx$$

## Why can we do this?

We are computing  $\nabla_{\theta} \mathbb{E}_{p_{\theta}(x|\mathcal{D})}[\mathcal{L}(x)] = \nabla_{\theta} \int p_{\theta}(x|\mathcal{D}) \mathcal{L}(x) dx$ .

We get the gradient by the change of variables:

$$\begin{aligned} p(x) &= \left| \frac{d\epsilon}{dx} \right| p(\epsilon) && \text{Change of variables,} \\ \implies |p(x) dx| &= |p(\epsilon) d\epsilon| && \text{Mass conservation.} \end{aligned}$$

$$\begin{aligned} \nabla_{\theta} \mathbb{E}_{p_{\theta}(x|\mathcal{D})}[\mathcal{L}(x)] &= \nabla_{\theta} \int p_{\theta}(x|\mathcal{D}) \mathcal{L}(x) dx \\ &= \nabla_{\theta} \int p(\epsilon) \mathcal{L}(x) d\epsilon \end{aligned}$$



## Why can we do this?

We are computing  $\nabla_{\theta} \mathbb{E}_{p_{\theta}(x|\mathcal{D})}[\mathcal{L}(x)] = \nabla_{\theta} \int p_{\theta}(x|\mathcal{D}) \mathcal{L}(x) dx$ .

We get the gradient by the change of variables:

$$\begin{aligned} p(x) &= \left| \frac{d\epsilon}{dx} \right| p(\epsilon) && \text{Change of variables,} \\ \implies |p(x) dx| &= |p(\epsilon) d\epsilon| && \text{Mass conservation.} \end{aligned}$$

$$\begin{aligned} \nabla_{\theta} \mathbb{E}_{p_{\theta}(x|\mathcal{D})}[\mathcal{L}(x)] &= \nabla_{\theta} \int p_{\theta}(x|\mathcal{D}) \mathcal{L}(x) dx \\ &= \nabla_{\theta} \int p(\epsilon) \mathcal{L}(x) d\epsilon = \nabla_{\theta} \int p(\epsilon) \mathcal{L}(g(\epsilon, \theta)) d\epsilon \end{aligned}$$

## Why can we do this?

We are computing  $\nabla_{\theta} \mathbb{E}_{p_{\theta}(x|\mathcal{D})}[\mathcal{L}(x)] = \nabla_{\theta} \int p_{\theta}(x|\mathcal{D}) \mathcal{L}(x) dx$ .

We get the gradient by the change of variables:

$$\begin{aligned} p(x) &= \left| \frac{d\epsilon}{dx} \right| p(\epsilon) && \text{Change of variables,} \\ \implies |p(x) dx| &= |p(\epsilon) d\epsilon| && \text{Mass conservation.} \end{aligned}$$

$$\begin{aligned} \nabla_{\theta} \mathbb{E}_{p_{\theta}(x|\mathcal{D})}[\mathcal{L}(x)] &= \nabla_{\theta} \int p_{\theta}(x|\mathcal{D}) \mathcal{L}(x) dx \\ &= \nabla_{\theta} \int p(\epsilon) \mathcal{L}(x) d\epsilon = \nabla_{\theta} \int p(\epsilon) \mathcal{L}(g(\epsilon, \theta)) d\epsilon \\ &= \int p(\epsilon) \nabla_{\theta} \mathcal{L}(g(\epsilon, \theta)) d\epsilon \end{aligned}$$

## Why can we do this?

We are computing  $\nabla_{\theta} \mathbb{E}_{p_{\theta}(x|\mathcal{D})}[\mathcal{L}(x)] = \nabla_{\theta} \int p_{\theta}(x|\mathcal{D}) \mathcal{L}(x) dx$ .

We get the gradient by the change of variables:

$$\begin{aligned} p(x) &= \left| \frac{d\epsilon}{dx} \right| p(\epsilon) && \text{Change of variables,} \\ \implies |p(x) dx| &= |p(\epsilon) d\epsilon| && \text{Mass conservation.} \end{aligned}$$

$$\begin{aligned} \nabla_{\theta} \mathbb{E}_{p_{\theta}(x|\mathcal{D})}[\mathcal{L}(x)] &= \nabla_{\theta} \int p_{\theta}(x|\mathcal{D}) \mathcal{L}(x) dx \\ &= \nabla_{\theta} \int p(\epsilon) \mathcal{L}(x) d\epsilon = \nabla_{\theta} \int p(\epsilon) \mathcal{L}(g(\epsilon, \theta)) d\epsilon \\ &= \int p(\epsilon) \nabla_{\theta} \mathcal{L}(g(\epsilon, \theta)) d\epsilon = \mathbb{E}_{p(\epsilon)}[\nabla_{\theta} \mathcal{L}(g(\epsilon, \theta))] \end{aligned}$$

## MC approximation

How to compute  $\mathbb{E}_{p(\epsilon)}[\nabla_{\theta} \mathcal{L}(g(\epsilon, \theta))]$ ?

1. Sample  $\epsilon^{(s)} \sim p(\epsilon)$ .
2.  $\nabla_{\theta} \mathcal{L}(x) \simeq \frac{1}{S} \sum_{s=1}^S \nabla_{\theta} \mathcal{L}(g(\epsilon^{(s)}, \theta))$

## MC approximation

How to compute  $\mathbb{E}_{p(\epsilon)}[\nabla_{\theta} \mathcal{L}(g(\epsilon, \theta))]$ ?

1. Sample  $\epsilon^{(s)} \sim p(\epsilon)$ .
2.  $\nabla_{\theta} \mathcal{L}(x) \simeq \frac{1}{S} \sum_{s=1}^S \nabla_{\theta} \mathcal{L}(g(\epsilon^{(s)}, \theta))$

What happens to the multiplication by  $p(\cdot)$ ?

# MC approximation

How to compute  $\mathbb{E}_{p(\epsilon)}[\nabla_{\theta} \mathcal{L}(g(\epsilon, \theta))]$ ?

1. Sample  $\epsilon^{(s)} \sim p(\epsilon)$ .
2.  $\nabla_{\theta} \mathcal{L}(x) \simeq \frac{1}{S} \sum_{s=1}^S \nabla_{\theta} \mathcal{L}(g(\epsilon^{(s)}, \theta))$

What happens to the multiplication by  $p(\cdot)$ ?

$$\mathbb{E}_{p(x)}[f(x)] = \int p(x) f(x) \, dx \simeq \frac{1}{S} \sum_{s=1}^S f(x^{(s)}) \quad (7)$$

# Table of Contents

A Few Examples

Why do we need this?

Continuous Random Variables: One-Liners

Discrete Random Variables: REINFORCE

## REINFORCE or the log-derivative trick

We can use REINFORCE for continuous variables, too, but one-liners are typically easier and implemented by default.

Recall that

$$\nabla \log f(x) = \frac{\nabla f(x)}{f(x)}.$$



# REINFORCE

Again, we're computing the expectation of the gradient:

$$\nabla_{\theta} \mathbb{E}_{p_{\theta}(x|\mathcal{D})}[\mathcal{L}(x)] = \int \nabla_{\theta} p_{\theta}(x|\mathcal{D}) \mathcal{L}(x) \mathrm{d}x$$

# REINFORCE

Again, we're computing the expectation of the gradient:

$$\begin{aligned}\nabla_{\theta} \mathbb{E}_{p_{\theta}(x|\mathcal{D})}[\mathcal{L}(x)] &= \int \nabla_{\theta} p_{\theta}(x|\mathcal{D}) \mathcal{L}(x) \mathrm{d}x \\ &= \int \frac{p_{\theta}(x|\mathcal{D})}{p_{\theta}(x|\mathcal{D})} \nabla_{\theta} p_{\theta}(x|\mathcal{D}) \mathcal{L}(x) \mathrm{d}x\end{aligned}$$

# REINFORCE

Again, we're computing the expectation of the gradient:

$$\begin{aligned}\nabla_{\theta} \mathbb{E}_{p_{\theta}(x|\mathcal{D})}[\mathcal{L}(x)] &= \int \nabla_{\theta} p_{\theta}(x|\mathcal{D}) \mathcal{L}(x) \, dx \\ &= \int \frac{p_{\theta}(x|\mathcal{D})}{p_{\theta}(x|\mathcal{D})} \nabla_{\theta} p_{\theta}(x|\mathcal{D}) \mathcal{L}(x) \, dx \\ &= \int p_{\theta}(x|\mathcal{D}) \nabla_{\theta} \log p_{\theta}(x|\mathcal{D}) \mathcal{L}(x) \, dx\end{aligned}$$

# REINFORCE

Again, we're computing the expectation of the gradient:

$$\begin{aligned}\nabla_{\theta} \mathbb{E}_{p_{\theta}(x|\mathcal{D})}[\mathcal{L}(x)] &= \int \nabla_{\theta} p_{\theta}(x|\mathcal{D}) \mathcal{L}(x) \, dx \\&= \int \frac{p_{\theta}(x|\mathcal{D})}{p_{\theta}(x|\mathcal{D})} \nabla_{\theta} p_{\theta}(x|\mathcal{D}) \mathcal{L}(x) \, dx \\&= \int p_{\theta}(x|\mathcal{D}) \nabla_{\theta} \log p_{\theta}(x|\mathcal{D}) \mathcal{L}(x) \, dx \\&= \mathbb{E}_{p_{\theta}(x|\mathcal{D})}[\nabla_{\theta} \log p_{\theta}(x|\mathcal{D}) \mathcal{L}(x)]\end{aligned}$$

# REINFORCE

Again, we're computing the expectation of the gradient:

$$\begin{aligned}\nabla_{\theta} \mathbb{E}_{p_{\theta}(x|\mathcal{D})}[\mathcal{L}(x)] &= \int \nabla_{\theta} p_{\theta}(x|\mathcal{D}) \mathcal{L}(x) \mathrm{d}x \\&= \int \frac{p_{\theta}(x|\mathcal{D})}{p_{\theta}(x|\mathcal{D})} \nabla_{\theta} p_{\theta}(x|\mathcal{D}) \mathcal{L}(x) \mathrm{d}x \\&= \int p_{\theta}(x|\mathcal{D}) \nabla_{\theta} \log p_{\theta}(x|\mathcal{D}) \mathcal{L}(x) \mathrm{d}x \\&= \mathbb{E}_{p_{\theta}(x|\mathcal{D})}[\nabla_{\theta} \log p_{\theta}(x|\mathcal{D}) \mathcal{L}(x)] \\&\simeq \frac{1}{S} \sum_{s=1}^S \nabla_{\theta} \log p_{\theta}(x|\mathcal{D}) \mathcal{L}(x)\end{aligned}$$