

## **A. Identification of PIs:**

Raymond W. Arritt  
Associate Professor  
Iowa State University  
Department of Agronomy  
3010 Agronomy Hall, Iowa State University, Ames, IA 50011-1010  
rwarritt@iastate.edu  
Tel: (515) 294-9870  
Fax: (515) 294-3163

Project Title: Comparative Regional Climate Simulations

Eugene S. Takle  
Professor  
Iowa State University  
Dept. of Geological and Atmospheric Sciences, and Department of Agronomy  
3010 Agronomy Hall, Iowa State University, Ames, IA 50011-1010  
gstakle@iastate.edu  
Tel: (515) 294-4758  
Fax: (515) 294-3163

William J. Gutowski, Jr.  
Associate Professor  
Iowa State University  
Department of Geological and Atmospheric Sciences, and Department of Agronomy  
3010 Agronomy Hall, Iowa State University, Ames, IA 50011-1010  
gutowski@iastate.edu  
Tel: (515) 294-4758  
Fax: (515) 294-3163

**B. Research Sponsor:**

Name: Electric Power Research Institute

Address: 3412 Hillview Ave., PO Box 10412, Palo Alto, CA 94303

Project manager: Dr. Charles Hakkarinen

Contract no: WO 4446-03

Date effective: 15 December 1996

Date of expiration/renewal: 14 December 1998

PIs: (same as CSL proposal)

Administrating officer: Richard Hasbrook, Contracts and Grants Administrator

Address: 213 Beardshear Hall, Iowa State Univ., Ames, IA 50011

Tel: 515-294-5225

E-mail: grants@iastate.edu

## **C. Programmers:**

Raymond W. Arritt  
Associate Professor  
Tel: (515) 294-9870  
Fax: (515) 294-3163  
3010 Agronomy Hall, Iowa State University, Ames, IA 50011-1010  
rwarritt@iastate.edu

Eugene S. Takle  
Professor  
Tel: (515) 294-9871  
Fax: (515) 294-3163  
Dept. of Geological and Atmospheric Sciences, and Dept. of Agronomy  
3010 Agronomy Hall, Iowa State University, Ames, IA 50011-1010  
gstakle@iastate.edu

William J. Gutowski, Jr.  
Associate Professor  
Tel: (515) 294-5632  
Fax: (515) 294-3163  
Department of Geological and Atmospheric Sciences, and Department of Agronomy  
3010 Agronomy Hall, Iowa State University, Ames, IA 50011-1010  
gutowski@iastate.edu

Moti Segal  
Scientist  
Tel: (515) 294-2804  
Fax: (515) 294-3163  
Department of Agronomy  
3010 Agronomy Hall, Iowa State University, Ames, IA 50011-1010  
segal@iastate.edu

Dr. Zaitao Pan  
Tel: (515) 294-4758  
Fax: (515) 294-3163  
Department of Agronomy  
3010 Agronomy Hall, Iowa State University, Ames, IA 50011-1010  
panz@iastate.edu

## **D. Description of Calculations**

### ***D.1. Appropriateness for CSL***

The proposed research adheres to the “Criteria for Use of the CSL.” It is part of a collaborative effort with impacts-assessment scientists, and builds upon the recommendations of a workshop, “Interfacing Mesoscale Climate Models with Impact Assessment,” that we hosted on 5–6 December 1996 at NCAR. Our work will lead ultimately to regional climate change impacts assessments in support of the IPCC Third Assessment Report (TAR). The long-term runs to be performed here require intensive, dedicated supercomputing resources such as the CSL.

In terms of the specific criteria for use of the CSL:

1. The research is being conducted by U.S. researchers at a U.S. institution.
2. The calculations require dedicated supercomputing. Regional climate models are to be run for multiple simulations of ten years or more, a task that is feasible with CSL resources but would not be practicable without dedicated high-performance computing.
3. The project will produce regional climate projections to be used in the IPCC Third Assessment Report. As indicated above, the project is being conducted in close collaboration with climate impacts scientists.
4. The proposed research is a critical part of our EPRI peer-reviewed research project.

## **D.2. Discussion of the Science**

### **D.2.1. Introduction**

The goal of this project is to evaluate regional implications of global climate change by a suite of comparative simulations using mesoscale models. In the first stage we will run several regional climate models for an extended period (at least 10 years) on a common domain driven by NCEP/NCAR reanalysis boundary conditions for comparison with observed climate conditions. We will then use these same models to perform simulations driven by GCM simulations for current climate and future climate perturbed by enhanced CO<sub>2</sub> and aerosol effects.

Atmospheric mesoscale models have been used to study and forecast the behavior of the atmosphere over periods usually of one to several days. In recent years, scientists have begun applying these models to regional climate (continental scale or smaller) in simulations lasting several weeks to several years (e.g., Giorgi et al. 1993; Jones et al. 1995; Hiriakuchi and Giorgi 1995; Copeland et al. 1996; Christensen et al. 1996). These models can be desirable tools for providing output for climate impact assessment because natural and human impacts of climate change are typically felt on the regional scales.

Climate simulation with mesoscale models is a relatively new approach and its range of validity must be documented. Biases that are of little consequence for the usual time scales of mesoscale simulations may become important when much longer integrations are performed. The adequacy of GCM-supplied boundary conditions for the regional model may also be a significant source of error (Pielke et al. 1997). We will address these and other scientific and technical issues in evaluating the suitability of mesoscale models for climate impacts assessment.

### **D.2.2. Research plan**

#### ***D.2.2.1. Long-Term Regional Climate Simulations***

We will deploy a small group of mesoscale models to perform a set of extended regional climate simulations for a domain that covers most of the continental U.S. It is desirable to use long runs (here 10 years; 1979-1988 which overlaps the AMIP period) in order to include a broader range of climate regimes and to give a more representative sampling with more degrees of freedom. We consider 10 years to be achievable and will attempt 20-year simulations given sufficient computing resources through the NCAR CSL. A second important point is that the longer simulations provide more information on the slow parts of the hydrological cycle. Here, the longer period is important because it is not just the number of hydrological extremes that is important, but their *sequencing* is crucial as well (for example, the 1993 floods were worsened because the soil already was anomalously wet from the previous year's rainfall).

A central question for mesoscale climate modeling is: What added value can it give to global climate-change simulation? The improved resolution should enable more detailed representations both of physical processes (such as the central U.S. nocturnal low-level jet and its effect on the generation of precipitation) and of geographic features (such as land use and topography). Our preliminary results indicate that the regional models give improved representation of the low-level

jet compared with the NCEP/NCAR reanalysis, and that the diurnal variability of warm-season precipitation is better predicted.

The simulations will be done using three distinct regional models to give a preliminary indication of the robustness of regional climate projections. The regional models we will run at the NCAR CSL are the NCAR Regional Climate Model (RegCM2; Giorgi et al. 1993) and the Colorado State University Regional Atmospheric Modeling System (RAMS; Pielke et al. 1992). The third model, HIRHAM4, is being run by Jens Hesselbjerg Christensen at the Danish Meteorological Institute and will not require NCAR resources. Dr. Filippo Giorgi of NCAR, the lead developer of RegCM2, advises us on the use of RegCM2.

#### ***D.2.2.2. Simulation design and computational strategy***

The simulation domain covers the continental U.S. and most of Mexico with a forced outer frame surrounding a free inner region. The horizontal grid uses a nominal resolution of 60 km. The simulations will begin with a “spinup” period that will not be considered as part of the results to be supplied to impacts assessors.

The simulations are being run with three sets of initial and lateral boundary conditions:

- First, climate of the recent past (1979-1988, included within the AMIP period) is being simulated using analyses of observed data to provide the same initial and boundary conditions for all three of the regional models. Driving data are given by the atmospheric reanalyses produced by the National Centers for Environmental Prediction (NCEP) and the National Center for Atmospheric Research (NCAR), using a T62 resolution spectral forecast model and a variational objective analysis (see Kalnay et al. 1996 for a description of the reanalysis methodology). Within caveats regarding analysis accuracy the reanalysis can be viewed as the limiting case of a “perfect” GCM in which to nest the regional model. The regional model results can be compared directly to observed data in order to establish the models’ skill at replicating regional climate and its variability.
- Second, the same three regional models will perform 10-year simulations of regional climate for the same location but with initial and boundary conditions supplied from Hadley Centre GCM simulations of present climate. The second-generation Hadley Centre model is a coupled atmosphere-ocean GCM run on a 2.5° by 3.75° latitude-longitude Arakawa-B grid, which is slightly coarser than the T62 transform grid used in the NCEP/NCAR reanalysis. By comparing these results with the reanalysis driven simulations and with observations we can identify biases that are attributable to driving the regional models with large-scale information from this GCM rather than from observations.
- The third set of simulations will be identical to the second except that initial and boundary data will be taken from Hadley Centre output for future climate perturbed by increased greenhouse gases and by aerosol effects (Johns et al. 1997). Comparison of these results with the preceding two sets allows us to interpret the projected regional climate change in light of the errors induced both by the regional models and by the use of the GCM-supplied initial and boundary conditions.

Codes to develop the regional model boundary conditions from the NCEP/NCAR reanalysis data and the Hadley Centre GCM output have been developed. The NCEP reanalysis output is available at NCAR. We have received and processed 10 years of the Hadley Centre results for both the control climate and CO<sub>2</sub>/aerosol scenario, and soon will receive another 10 years. Additional datasets describing topography, land use, ocean and lake surface temperature have been assembled.

We will use standard statistical measures such as threat scores and pattern correlations in addition to means and higher-order statistics (e.g., Takle and Mearns 1995) to evaluate simulation accuracy for mean behavior, spatial and temporal variability, and extreme events. Additional diagnostics will be computed in order to evaluate whether the regional climate model is appropriately coupled to the observed or GCM-supplied climate (as in Jones et al. 1995). The results will provide an ensemble estimate of simulation accuracy through comparison with observations. This information will help place bounds on types of impacts assessments that can be reasonably attempted using current models. Output analysis may also indicate immediately fruitful avenues for model improvement. Close consultation will be maintained with model developers to correct any anomalous behavior that should arise during the extended simulations.

### **D.2.3. Interactions with impacts assessments**

The physical climate simulated by a mesoscale model provides input into impact assessment models such as agricultural, water resources or human-health models. This is the most direct and immediate use of mesoscale climate simulation. Numerous examples exist of such assessments using global models, which can guide mesoscale-model applications. Output archives for the simulations will be created in ongoing consultation with impacts assessors to ensure that we are archiving the most appropriate data. We will make output available in a timely manner so that impacts assessments for the IPCC Third Assessment Report can start as soon as possible.

The Principal Investigators conducted a workshop, "Interfacing Mesoscale Climate Models with Impact Assessment," held 5–6 December 1996 at NCAR (supported by the Electric Power Research Institute, EPRI). The workshop provided important guidance for developing regional climate simulations and output archives that support impacts assessment. Among the specific recommendations made by the impacts assessors was a consensus that increased computational resources would be better devoted to longer simulations than to improved spatial resolution, so long as the horizontal resolution is not much coarser than about 50 km. Interactions with the impacts community are continuing in part through collaboration with a parallel statistical downscaling project led by T. Wigley and R. Wilby at NCAR. PIRCS will work with WGNE and other oversight groups to establish standards, procedures, and templates for intercomparison experiments and collaborative research programs.

### **D.2.4. Interactions with PIRCS and other modeling programs**

The project proposed here is a part of a larger effort, the "Project for Intercomparison of Regional Climate Simulations" (PIRCS). The mission of PIRCS, as established by the First PIRCS Workshop (Takle 1995), is to advance the status of regional climate modeling by evaluating the strengths and weaknesses of limited-area models and their component procedures through systematic comparative simulations. PIRCS consists of an on-going series of intercomparison

experiments using different regions of the world and different time periods. The first PIRCS experiment is an intercomparison using the same domain as in the present proposal but for shorter periods, specifically 60-day simulations during the 1988 drought and 1993 flood over the central U.S. Preliminary results for the first experiment have been obtained by nine modeling groups from North America, Europe and Australia. The PIRCS web site is located at <http://www.pircs.iastate.edu>.

PIRCS is a project of the International Institute of Theoretical and Applied Physics (IITAP), which is sponsored jointly by UNESCO and Iowa State University. Additional support for PIRCS (and the primary support for the mesoscale climate simulations) is provided under a contract from the Electric Power Research Institute (EPRI). PIRCS is also coordinating its activities with other international programs such as the Comparison of Mesoscale Prediction and Research Experiments (COMPARE), GEWEX (specifically GCIP), AMIP and the Working Group on Numerical Experimentation (WGNE) of the World Climate Research Program. The WGNE has appointed Dr. Andrew Staniforth to serve as liaison between the WGNE and PIRCS and to promote interaction between PIRCS and COMPARE (WGNE 1996). We also are maintaining close contact with related efforts in regional climate modeling and downscaling of climate projections, such as the European Community MERCURE project.

## D.2.5. References

- Christensen, J. H., O. B. Christensen, P. Lopez, E. van Meijaard and M. Botzet, 1996: The HIRHAM4 Regional Atmospheric Climate Model. Scientific Report 96-4, Danish Meteorological Institute, 51 pp.
- Copeland, J.H., R.A. Pielke and T.G.F. Kittel, 1996: Potential climatic impacts of vegetation change: A regional modeling study. *J. Geophys. Res.*, 101 (D3), 7409-7418.
- Giorgi, F., M.R. Marinucci and G.T. Bates, 1993: Development of a second generation regional climate model (RegCM2). Part I. *Mon. Wea. Rev.*, 121, 2794-2813.
- Grell, G. A., J. Dudhia and D. R. Stauffer, 1993: A Description of the Fifth-Generation Penn State/NCAR Mesoscale Model (MM5). NCAR Technical Note NCAR/TN-398+IA, National Center for Atmospheric Research, 116 pp.
- Hirakuchi, H., and F. Giorgi, 1995: Multi-year present day and 2XCO<sub>2</sub> simulations of monsoon-dominated climate over eastern Asia and Japan with a regional climate model nested in a general circulation model. *Journal of Geophysical Research*, 100, 21,105-21,126.
- IPCC, 1996: *Climate Change 1995. Impacts, Adaptations and Mitigation of Climate Change: Scientific-Technical Analyses*. R. Watson, M. Zinyowera and R. Moss, Eds., Cambridge University Press, 879 pp.
- Johns, T.C., R.E. Carnell, J.F. Crossley, J.M.Gregory, J.F.B. Mitchell, C.A. Senior, S.F.B. Tett and R.A. Wood, 1997: The second Hadley Centre coupled ocean-atmosphere GCM: model description, spinup and validation. *Climate Dynamics*, **12**, 103-134.
- Jones, R.G., J.M. Murphy and M. Noguer, 1995: Simulation of climate change over Europe using a nested regional climate model. Part I. Assessment of control climate, including sensitivity to location of lateral boundaries. *Quart. J. Roy. Met. Soc.*, 121, 1413-1449.



Kalnay, E., M. Kanamitsu, R. Kistler, W. Collins, D. Deaven, L. Gandin, M. Iredell, S. Saha, G. White, J. Woollen, Y. Zhu, M. Chelliah, W. Ebisuzaki, W. Higgins, J. Janowiak, K. C. Mo, C. Ropelewski, J. Wang, A. Leetma, R. Reynolds, R. Jenne and D. Joseph, 1996: The NCEP/NCAR 40-year reanalysis project. *Bull. Amer. Meteor. Soc.*, **77**, 437-471.

Pielke, R. A., W. R. Cotton, R. L. Walko, C. J. Tremback, W. A. Lyons, L. D. Grasso, M. E. Nicholls, M. D. Moran, D. A. Wesley, T. J. Lee and J. H. Copeland, 1992: A comprehensive meteorological modeling system - RAMS. *Meteor. Atmos. Phys.*, **49**, 69 - 91.

Pielke, R.A., K. Arpe, K. Keuler, R. Laprise, M.E. Mann, A. Rinke and D.E. Parker, 1997: Summary report: Workshop on High Resolution Climate Modeling, Switzerland, 1996. Submitted to *Eos, Trans. Amer. Geophys. Union*.

PIRCS: Project to Intercompare Regional Climate Simulations. Iowa State University, <http://www.pircs.iastate.edu>

Takle, E. S., and L. O. Mearns, 1995b: Midwest Temperature Means, Extremes, and Variability: Analysis of Results from Climate Models. In G. R. Carmichael, G. E. Folk, and J. L. Schnoor, ed., *Preparing for Global Change: A Midwestern Perspective*. Academic Publishing bv, Amsterdam, p . 135-142.

Takle, E. S., 1995: Project to Intercompare Regional Climate Simulations (PIRCS), Preliminary Workshop, 17-18 November 1994. *Bull. Amer. Meteor. Soc.*, **76**, 1625-1626.

WGNE, 1996: Report of the Eleventh Session of the CAS/JSC Working Group on Numerical Experimentation (Reading, UK, 30 October - 3 November 1995). CAS/JSC WGNE Report No. 11, World Meteorological Organization, 43 pp.

## **E. Computing Requirements**

### ***E.1. Computing resources needed:***

Estimated computing resources are approximately 100 C90 hours per year of simulation for the RegCM2 runs on the domain we intend to use. The 20-year runs would thus require 2,000 C90 hours for a single scenario. The RAMS runs would require computing resources about 50% larger than RegCM2. Combining both models for all three scenarios, plus some allowance for test runs we request 18,000 C90 CPU hours.

### ***E.2. Machine preference***

We request the C90 for this project. The RegCM2 and RAMS models are not parallelized (although experimental parallel versions of RAMS exist). Therefore each of the models will be run on a single C90 processor and will not be run in a parallel-processing configuration. In order to reduce wall-clock time, the models (RegCM2 and RAMS) and scenarios can be run simultaneously on independent processors.

### ***E.3. Memory, output and analysis***

Each simulation requires about 8 MW of memory. Output is about 18 GB per year. Most of the quantitative analysis will be performed at NCAR. The output also will be written on magnetic tapes and transferred to Iowa State University where data can be visualized using tools such as Vis5D.

### ***E.4. Utilization of request***

Utilization will begin immediately upon allocation of the request, as the RegCM2 model has already been implemented on Cray systems and the boundary-condition (reanalysis) data are in hand. RAMS will require setup on the Cray and those runs are projected to begin in June 1998.

### ***E.5. Expected completion date:***

31 December 1998

### ***E.6. Milestones:***

Contractual obligations from the research sponsor (EPRI) require us to have the simulations completed by the end of the contract (December 1998). There is some flexibility in our schedule in that both models have restart capabilities; thus, the use of the C90 allocation need not be continuous, and we can work around other projects that may require all processors on the machine.

## **F. Proposal Manager**

The SCD point of contact will be:

Dr. Zaitao Pan

Tel: (515) 294-4758

Fax: (515) 294-3163

Department of Agronomy

3010 Agronomy Hall, Iowa State University, Ames, IA 50011-1010

email [panz@iastate.edu](mailto:panz@iastate.edu)