


Asymptotics of Lattice Walks

Alexander Kroitor

University of Waterloo

Dec 2023



Generating Functions

Enumerative combinatorics is about studying sequences.

Generating Functions

Enumerative combinatorics is about studying sequences.
Generating functions are a powerful tool.

Generating Functions

Enumerative combinatorics is about studying sequences.
Generating functions are a powerful tool.

Definition (Generating Functions)

The **generating function** of the sequence $\{a_n\}_{n \geq 0}$ is

$$A(z) = \sum_{n \geq 0} a_n z^n.$$

Generating Functions

Enumerative combinatorics is about studying sequences.
Generating functions are a powerful tool.

Definition (Generating Functions)

The **generating function** of the sequence $\{a_n\}_{n \geq 0}$ is

$$A(z) = \sum_{n \geq 0} a_n z^n.$$

We care about *asymptotics* instead of exact values.

Generating Functions

Enumerative combinatorics is about studying sequences.
Generating functions are a powerful tool.

Definition (Generating Functions)

The **generating function** of the sequence $\{a_n\}_{n \geq 0}$ is

$$A(z) = \sum_{n \geq 0} a_n z^n.$$

We care about *asymptotics* instead of exact values.

We can use this framework to count lattice walks!

Definition (Lattice Walk Model)

A d -dimensional **lattice walk model** consists of:

- a **step set** $S \subseteq \mathbb{Z}^d$,
- a **restricting region** $R \subseteq \mathbb{Z}^d$,
- a **starting point** $p \in R$,
- a **terminal set** $T \subseteq R$.

A walk in this lattice walk model consists of a walk that starts at p , stays in R at all times, ends at T , and takes steps using S .

Definition (Lattice Walk Model)

A d -dimensional **lattice walk model** consists of:

- a **step set** $S \subseteq \mathbb{Z}^d$,
- a **restricting region** $R \subseteq \mathbb{Z}^d$,
- a **starting point** $p \in R$,
- a **terminal set** $T \subseteq R$.

A walk in this lattice walk model consists of a walk that starts at p , stays in R at all times, ends at T , and takes steps using S .

We always let $p = 0$ and $T = R$.

Main Question

Given a step set S and a restricting region R , how many lattice walks of length n are there using S ?

Main Question

Given a step set S and a restricting region R , how many lattice walks of length n are there using S ?

Reformulated in generating function language:

Main Question

Given a step set S and a restricting region R , let s_n be the number of lattice walks of length n using S and staying in R .


What is the coefficient of z^n in $S(z) = \sum_{n \geq 0} s_n z^n$?

Unrestricted Walks

If $R = \mathbb{Z}^d$ this is easy.

Unrestricted NSEW steps

Let $S = \{(\pm 1, 0), (0, \pm 1)\}$ and $R = \mathbb{Z}^2$.




Then there are 4 possible steps to take at each time. Thus there are 4^n walks of length n using S .

Unrestricted Walks

If $R = \mathbb{Z}^d$ this is easy.

Unrestricted NSEW steps

Let $S = \{(\pm 1, 0), (0, \pm 1)\}$ and $R = \mathbb{Z}^2$.



Then there are 4 possible steps to take at each time. Thus there are 4^n walks of length n using S .

In general:

Unrestricted Walks

There are $|S|^n$ unrestricted lattice walks of length n using S .

Unrestricted walks are simple. What about other domains?

Halfspace Walks

Unrestricted walks are simple. What about other domains?

Definition (Halfspaces)

$\mathbb{Z}^{d-1} \times \mathbb{N}$ is a **halfspace**.

Unrestricted walks are simple. What about other domains?

Definition (Halfspaces)

$\mathbb{Z}^{d-1} \times \mathbb{N}$ is a **halfspace**.

Let S be any step set and $R = \mathbb{Z}^{d-1} \times \mathbb{N}$ be a halfspace. Let s_n be number of walks.

Unrestricted walks are simple. What about other domains?

Definition (Halfspaces)

$\mathbb{Z}^{d-1} \times \mathbb{N}$ is a **halfspace**.

Let S be any step set and $R = \mathbb{Z}^{d-1} \times \mathbb{N}$ be a halfspace. Let s_n be number of walks.

It turns out $S(z) = \sum_{n \geq 0} s_n z^n$ is **algebraic**.

Unrestricted walks are simple. What about other domains?

Definition (Halfspaces)

$\mathbb{Z}^{d-1} \times \mathbb{N}$ is a **halfspace**.

Let S be any step set and $R = \mathbb{Z}^{d-1} \times \mathbb{N}$ be a halfspace. Let s_n be number of walks.

It turns out $S(z) = \sum_{n \geq 0} s_n z^n$ is **algebraic**.

This means finding asymptotics for s_n is automatic.

Asymptotics for s_n is automatic for halfspaces. What other domains are there?

Asymptotics for s_n is automatic for halfspaces. What other domains are there?

Definition (Orthants)

\mathbb{N}^d is an **orthant**.

This models queues (since negative people does not make sense).

Asymptotics for s_n is automatic for halfspaces. What other domains are there?

Definition (Orthants)

\mathbb{N}^d is an **orthant**.

This models queues (since negative people does not make sense).

This is too hard in general. How do we fix this? Restricting S .

Highly Symmetric Step Sets

We enforce *small steps*.

Let $S \subseteq \{-1, 0, 1\}^d \setminus 0$.

Highly Symmetric Step Sets

We enforce *small steps*.

Let $S \subseteq \{-1, 0, 1\}^d \setminus 0$. Define

$$S(\mathbf{z}) = S(z_1, \dots, z_d) = \sum_{s \in S} \mathbf{z}^s = \sum_{s \in S} z_1^{s_1} \cdots z_d^{s_d}.$$

Highly Symmetric Step Sets

We enforce *small steps*.

Let $S \subseteq \{-1, 0, 1\}^d \setminus \{0\}$. Define

$$S(\mathbf{z}) = S(z_1, \dots, z_d) = \sum_{s \in S} \mathbf{z}^s = \sum_{s \in S} z_1^{s_1} \cdots z_d^{s_d}.$$

Example

If $S = \{(\pm 1, 0), (0, \pm 1), (1, 1)\}$



$$S(z_1, z_2) = z_1 + \frac{1}{z_1} + z_2 + \frac{1}{z_2} + z_1 z_2$$

Highly Symmetric Step Sets

We also enforce some amount of *symmetry*.

Definition (Highly Symmetric Step Sets)

A step set S is **highly symmetric** if

$$S(z_1, \dots, z_j, \dots, z_d) = S\left(z_1, \dots, \frac{1}{z_j}, \dots, z_d\right)$$

for all $1 \leq j \leq d$.

Highly Symmetric Step Sets


Asymptotics for highly symmetric step sets is known.

Theorem (Melczer Mishna 2016 [MM16])

Suppose S is a highly symmetric step set. If s_n is the number of walks of length n using S that remains in \mathbb{N}^d then

$$s_n \sim \left[\frac{|S|^{d/2}}{\pi^{d/2} (a_1 \cdots a_d)^{1/2}} \right] \cdot \frac{|S|^n}{n^{d/2}},$$

where a_j is the number of steps in S that have j -th coordinate 1.



Mostly Symmetric Step Sets

Being highly symmetric is very restrictive.

Mostly Symmetric Step Sets

Being highly symmetric is very restrictive.

Definition (Mostly Symmetric Step Sets)

A step set S is **mostly symmetric** if

$$S(z_1, \dots, z_j, \dots, z_d) = S\left(z_1, \dots, \frac{1}{z_j}, \dots, z_d\right)$$

for all $1 \leq j < d$.

Mostly Symmetric Step Sets

Being highly symmetric is very restrictive.

Definition (Mostly Symmetric Step Sets)

A step set S is **mostly symmetric** if

$$S(z_1, \dots, z_j, \dots, z_d) = S\left(z_1, \dots, \frac{1}{z_j}, \dots, z_d\right)$$

for all $1 \leq j < d$.

This means we can write

$$S(\mathbf{z}) = \frac{1}{z_d} A(z_1, \dots, z_{d-1}) + Q(z_1, \dots, z_{d-1}) + z_d B(z_1, \dots, z_{d-1})$$

where A, Q, B are all highly symmetric.

Asymptotics depend on what the average direction of a step is.

Definition (Drift)

We say that

- S has **positive drift** if $B(\mathbf{1}) - A(\mathbf{1}) > 0$,
- S has **negative drift** if $B(\mathbf{1}) - A(\mathbf{1}) < 0$,
- S has **zero drift** if $B(\mathbf{1}) - A(\mathbf{1}) = 0$.

Asymptotics depend on what the average direction of a step is.

Definition (Drift)

We say that

- S has **positive drift** if $B(\mathbf{1}) - A(\mathbf{1}) > 0$,
- S has **negative drift** if $B(\mathbf{1}) - A(\mathbf{1}) < 0$,
- S has **zero drift** if $B(\mathbf{1}) - A(\mathbf{1}) = 0$.

Asymptotics for the positive and negative case are known.

Mostly Symmetric Step Sets

Asymptotics depend on what the average direction of a step is.

Definition (Drift)

We say that

- S has **positive drift** if $B(\mathbf{1}) - A(\mathbf{1}) > 0$,
- S has **negative drift** if $B(\mathbf{1}) - A(\mathbf{1}) < 0$,
- S has **zero drift** if $B(\mathbf{1}) - A(\mathbf{1}) = 0$.

Asymptotics for the positive and negative case are known.

Asymptotics for the zero drift case is *open*.

Positive Drift Asymptotics

Theorem (Melczer Wilson [MW19])

If S is mostly symmetric with positive drift then

$$s_n \sim \left[\left(1 - \frac{A(1)}{B(1)} \right) \frac{S(1)^{d/2}}{(2\pi)^{d/2}} \cdot \frac{1}{(a_1 \cdots a_d)^{1/2}} \right] \cdot \frac{S(1)^n}{n^{d/2-1/2}}$$

Negative Drift Asymptotics

Theorem (Melczer Wilson [MW19])

If S is mostly symmetric with negative drift and $Q(z) \neq 0$ then

$$s_n \sim C_\rho \cdot \frac{S(1, \rho)^n}{n^{d/2+1}}.$$

If S is mostly symmetric with negative drift and $Q(z) = 0$ then

$$s_n \sim C_\rho \cdot \frac{S(1, \rho)^n}{n^{d/2+1}} + C_{-\rho} \cdot \frac{S(1, -\rho)^n}{n^{d/2+1}}.$$

Where ρ , C_ρ , and $C_{-\rho}$ are explicit constants.

Proof Overview

How is this done?

Through **analytic combinatorics in several variables** (ACSV).

Proof Overview

How is this done?

Through **analytic combinatorics in several variables** (ACSV).

ACSV can find asymptotics of sequences encoded by multivariate rational functions.

Proof Overview

How is this done?

Through **analytic combinatorics in several variables** (ACSV).

ACSV can find asymptotics of sequences encoded by multivariate rational functions.

Proof Outline

Proof Overview

How is this done?

Through **analytic combinatorics in several variables** (ACSV).

ACSV can find asymptotics of sequences encoded by multivariate rational functions.

Proof Outline

- Use **kernel method** to find GF for s_n in terms of higher dimensional rational function.

Proof Overview

How is this done?

Through **analytic combinatorics in several variables** (ACSV).

ACSV can find asymptotics of sequences encoded by multivariate rational functions.

Proof Outline

- Use **kernel method** to find GF for s_n in terms of higher dimensional rational function.
- Characterize singularities of this rational function.

Proof Overview

How is this done?

Through **analytic combinatorics in several variables** (ACSV).

ACSV can find asymptotics of sequences encoded by multivariate rational functions.

Proof Outline

- Use **kernel method** to find GF for s_n in terms of higher dimensional rational function.
- Characterize singularities of this rational function.
- Use ACSV to reduce to asymptotic behaviour of

$$\int A(\mathbf{z}) \exp [-n\phi(\mathbf{z})] d\mathbf{z}.$$

Proof Overview

How is this done?

Through **analytic combinatorics in several variables** (ACSV).

ACSV can find asymptotics of sequences encoded by multivariate rational functions.

Proof Outline

- Use **kernel method** to find GF for s_n in terms of higher dimensional rational function.
- Characterize singularities of this rational function.
- Use ACSV to reduce to asymptotic behaviour of

$$\int A(\mathbf{z}) \exp [-n\phi(\mathbf{z})] d\mathbf{z}.$$

- Approximate integral using saddle-point method.

Why does this approach fail for zero drift?

Why does this approach fail for zero drift?

At the singularity determining asymptotics both the denominator and numerator vanish. Other pathologies give an integral of the form

$$\int \frac{A(\mathbf{z})}{\mathbf{z}} \exp[-n\phi(\mathbf{z})] d\mathbf{z}.$$

We didn't know how to approximate this!

Why does this approach fail for zero drift?

At the singularity determining asymptotics both the denominator and numerator vanish. Other pathologies give an integral of the form

$$\int \frac{A(\mathbf{z})}{\mathbf{z}} \exp[-n\phi(\mathbf{z})] d\mathbf{z}.$$

We didn't know how to approximate this!

Now ~ 4 years later we have a better idea of how to approximate.

Zero Drift Asymptotics

Theorem (K. Melczer [KMon])

Suppose S is a mostly symmetric step set with zero drift. Let s_n be the number of walks of length n using S that remain in \mathbb{N}^d . Then

$$s_n \sim \left[\frac{|S|^{d/2}}{\pi^{d/2} (a_1 \cdots a_d)^{1/2}} \right] \cdot \frac{|S|^n}{n^{d/2}}$$

where a_j is the number of steps in S that have j -th coordinate 1.

Zero Drift Asymptotics

Theorem (K. Melczer [KMon])

Suppose S is a mostly symmetric step set with zero drift. Let s_n be the number of walks of length n using S that remain in \mathbb{N}^d . Then

$$s_n \sim \left[\frac{|S|^{d/2}}{\pi^{d/2} (a_1 \cdots a_d)^{1/2}} \right] \cdot \frac{|S|^n}{n^{d/2}}$$

where a_j is the number of steps in S that have j -th coordinate 1.

This is the same as highly symmetric: two different pathologies (vanishing and “boundary direction asymptotics” cancel).

Zero Drift Asymptotics

Theorem (K. Melczer [KMon])

Suppose S is a mostly symmetric step set with zero drift. Let s_n be the number of walks of length n using S that remain in \mathbb{N}^d . Then

$$s_n \sim \left[\frac{|S|^{d/2}}{\pi^{d/2} (a_1 \cdots a_d)^{1/2}} \right] \cdot \frac{|S|^n}{n^{d/2}}$$

where a_j is the number of steps in S that have j -th coordinate 1.

This is the same as highly symmetric: two different pathologies (vanishing and “boundary direction asymptotics” cancel).

This (sort of) finishes off all walks that can be done with ACSV.

Conclusion

Thank you for listening.

Bibliography



Alex Kroitor and Stephen Melczer.

Asymptotic lattice path enumeration using diagonals.

In preparation.



Stephen Melczer.

An Invitation to Analytic Combinatorics.

Springer Nature Switzerland, Cham, Switzerland.



Stephen Melczer and Marni Mishna.

Asymptotic lattice path enumeration using diagonals.

Algorithmica, 75(4):782–811, 2016.



Stephen Melczer and Mark C. Wilson.

Higher dimensional lattice walks: Connecting combinatorial and analytic behavior.

SIAM J. Discrete Math., 33(4):2140–2174, 2019.