

Asymptotics of Lattice Walks Via ACSV

Alexander Kroitor

University of Waterloo

April 2024




Figure: An Orthant Walk using the *Gessel* step Set

Definition

Let a_n be a complex sequence. The **generating function** corresponding to a_n is

$$A(z) = \sum_{n \geq 0} a_n z^n.$$

Definition

Let a_n be a complex sequence. The **generating function** corresponding to a_n is

$$A(z) = \sum_{n \geq 0} a_n z^n.$$

- GFs are a powerful tool in enumerative combinatorics.

Definition

Let a_n be a complex sequence. The **generating function** corresponding to a_n is

$$A(z) = \sum_{n \geq 0} a_n z^n.$$

- GFs are a powerful tool in enumerative combinatorics.
- Are amenable to (complex) analytic techniques.

Analytic Combinatorics in One Variable

Definition

Let a_n be a complex sequence. The **generating function** corresponding to a_n is

$$A(z) = \sum_{n \geq 0} a_n z^n.$$

- GFs are a powerful tool in enumerative combinatorics.
- Are amenable to (complex) analytic techniques.
- Goal of analytic techniques is almost always asymptotics.

Analytic Combinatorics in One Variable

Definition

Let a_n be a complex sequence. The **generating function** corresponding to a_n is

$$A(z) = \sum_{n \geq 0} a_n z^n.$$

- GFs are a powerful tool in enumerative combinatorics.
- Are amenable to (complex) analytic techniques.
- Goal of analytic techniques is almost always asymptotics.

Principles of AC

Let $A(z)$ be a generating function with underlying sequence a_n .

Analytic Combinatorics in One Variable

Definition

Let a_n be a complex sequence. The **generating function** corresponding to a_n is

$$A(z) = \sum_{n \geq 0} a_n z^n.$$

- GFs are a powerful tool in enumerative combinatorics.
- Are amenable to (complex) analytic techniques.
- Goal of analytic techniques is almost always asymptotics.

Principles of AC

Let $A(z)$ be a generating function with underlying sequence a_n .

- **exponential** behaviour of $a_n \leftrightarrow$ location of singularities

Analytic Combinatorics in One Variable

Definition

Let a_n be a complex sequence. The **generating function** corresponding to a_n is

$$A(z) = \sum_{n \geq 0} a_n z^n.$$

- GFs are a powerful tool in enumerative combinatorics.
- Are amenable to (complex) analytic techniques.
- Goal of analytic techniques is almost always asymptotics.

Principles of AC

Let $A(z)$ be a generating function with underlying sequence a_n .

- **exponential** behaviour of $a_n \leftrightarrow$ location of singularities
- **subexponential** behaviour of $a_n \leftrightarrow$ kind of singularities

Surjections

Let g_n count the number of functions that surject $[n]$ onto a set of the form $[r]$.

Surjections

Let g_n count the number of functions that surject $[n]$ onto a set of the form $[r]$. Then

$$G(z) = \sum_{n \geq 0} \frac{g_n}{n!} z^n = \frac{1}{2 - e^z}.$$

Surjections

Let g_n count the number of functions that surject $[n]$ onto a set of the form $[r]$. Then

$$G(z) = \sum_{n \geq 0} \frac{g_n}{n!} z^n = \frac{1}{2 - e^z}.$$

$G(z)$ has singularities at $\{\log 2 + 2\pi i k\}$. The singularity with smallest modulus is $\log 2$, which has a simple pole.

Surjections

Let g_n count the number of functions that surject $[n]$ onto a set of the form $[r]$. Then

$$G(z) = \sum_{n \geq 0} \frac{g_n}{n!} z^n = \frac{1}{2 - e^z}.$$

$G(z)$ has singularities at $\{\log 2 + 2\pi i k\}$. The singularity with smallest modulus is $\log 2$, which has a simple pole. Then

$$\frac{g_n}{n!} \sim C \left(\frac{1}{\log 2} \right)^n.$$

Here $C = \frac{1}{2 \log 2}$ is easily computed.

Surjections

Let g_n count the number of functions that surject $[n]$ onto a set of the form $[r]$. Then

$$G(z) = \sum_{n \geq 0} \frac{g_n}{n!} z^n = \frac{1}{2 - e^z}.$$

$G(z)$ has singularities at $\{\log 2 + 2\pi i k\}$. The singularity with smallest modulus is $\log 2$, which has a simple pole. Then

$$\frac{g_n}{n!} \sim C \left(\frac{1}{\log 2} \right)^n.$$

Here $C = \frac{1}{2 \log 2}$ is easily computed. If $H(z) = \frac{1}{(2 - e^z)^2}$ then

$$\frac{h_n}{n!} \sim Cn \cdot \left(\frac{1}{\log 2} \right)^n.$$

There are two main techniques in AC:

No. 1: Residue Theorem

Let $G(z) = \sum g_n z^n$ be meromorphic in $D_r(0)$ and analytic at 0.
Let ρ_1, \dots, ρ_k be a list of all the singularities of G in $D_r(0)$. Then

$$g_n = - \sum_{j=1}^k \operatorname{Res}_{z=\rho_j} \left(\frac{G(z)}{z^{n+1}} \right) + \frac{1}{2\pi i} \int_{|z|=r} G(z) \frac{dz}{z^{n+1}}.$$

Here Res is some easily computable operator.

There are two main techniques in AC:

No. 1: Residue Theorem

Let $G(z) = \sum g_n z^n$ be meromorphic in $D_r(0)$ and analytic at 0.
Let ρ_1, \dots, ρ_k be a list of all the singularities of G in $D_r(0)$. Then

$$g_n = - \sum_{j=1}^k \operatorname{Res}_{z=\rho_j} \left(\frac{G(z)}{z^{n+1}} \right) + \frac{1}{2\pi i} \int_{|z|=r} G(z) \frac{dz}{z^{n+1}}.$$

Here Res is some easily computable operator.

If $G(z)$ has no singularities we are out of luck!

No. 2: Saddle-Point Techniques

We describe the general idea.

No. 2: Saddle-Point Techniques

We describe the general idea.

Suppose that $G(z)z^{-n-1}$ has a *critical point* at $z = t$.

No. 2: Saddle-Point Techniques

We describe the general idea.

Suppose that $G(z)z^{-n-1}$ has a *critical point* at $z = t$.

- Write as before $g_n = \frac{1}{2\pi i} \int_{|z|=t} G(z) \frac{dz}{z^{n+1}},$

No. 2: Saddle-Point Techniques

We describe the general idea.

Suppose that $G(z)z^{-n-1}$ has a *critical point* at $z = t$.

- Write as before $g_n = \frac{1}{2\pi i} \int_{|z|=t} G(z) \frac{dz}{z^{n+1}}$,
- (trim) show that $g_n \sim \frac{1}{2\pi i} \int_{(|z|=t) \cap B_\epsilon(t)} G(z) \frac{dz}{z^{n+1}}$,

No. 2: Saddle-Point Techniques

We describe the general idea.

Suppose that $G(z)z^{-n-1}$ has a *critical point* at $z = t$.

- Write as before $g_n = \frac{1}{2\pi i} \int_{|z|=t} G(z) \frac{dz}{z^{n+1}}$,
- (trim) show that $g_n \sim \frac{1}{2\pi i} \int_{(|z|=t) \cap B_\epsilon(t)} G(z) \frac{dz}{z^{n+1}}$,
- (approx.) show that we can approximate

$$\int_{(|z|=t) \cap B_\epsilon(t)} G(z) \frac{dz}{z^{n+1}}$$

by replacing $G(z)z^{-n-1}$ by its leading terms,

No. 2: Saddle-Point Techniques

We describe the general idea.

Suppose that $G(z)z^{-n-1}$ has a *critical point* at $z = t$.

- Write as before $g_n = \frac{1}{2\pi i} \int_{|z|=t} G(z) \frac{dz}{z^{n+1}}$,
- (trim) show that $g_n \sim \frac{1}{2\pi i} \int_{(|z|=t) \cap B_\epsilon(t)} G(z) \frac{dz}{z^{n+1}}$,
- (approx.) show that we can approximate

$$\int_{(|z|=t) \cap B_\epsilon(t)} G(z) \frac{dz}{z^{n+1}}$$

by replacing $G(z)z^{-n-1}$ by its leading terms,

- (compute) compute what's left.

Analytic Combinatorics in Several Variables (ACSV)

This discussion generalizes to several variables.

Analytic Combinatorics in Several Variables (ACSV)

This discussion generalizes to several variables.

Multivariate GFs are natural to consider:

Analytic Combinatorics in Several Variables (ACSV)

This discussion generalizes to several variables.

Multivariate GFs are natural to consider:

- can often store “less nice” single-variable GFs in “nice” multivariate ones,

Analytic Combinatorics in Several Variables (ACSV)

This discussion generalizes to several variables.

Multivariate GFs are natural to consider:

- can often store “less nice” single-variable GFs in “nice” multivariate ones,
- can track multiple characteristics at once,

Analytic Combinatorics in Several Variables (ACSV)

This discussion generalizes to several variables.

Multivariate GFs are natural to consider:

- can often store “less nice” single-variable GFs in “nice” multivariate ones,
- can track multiple characteristics at once,
- strict generalization of single-variable GFs.

Analytic Combinatorics in Several Variables (ACSV)

This discussion generalizes to several variables.

Multivariate GFs are natural to consider:

- can often store “less nice” single-variable GFs in “nice” multivariate ones,
- can track multiple characteristics at once,
- strict generalization of single-variable GFs.

Same two techniques exist; must use both in conjunction.

Analytic Combinatorics in Several Variables (ACSV)

This discussion generalizes to several variables.

Multivariate GFs are natural to consider:

- can often store “less nice” single-variable GFs in “nice” multivariate ones,
- can track multiple characteristics at once,
- strict generalization of single-variable GFs.

Same two techniques exist; must use both in conjunction.

Singular set of GF is no longer discrete, now a complex variety.

Analytic Combinatorics in Several Variables (ACSV)

This discussion generalizes to several variables.

Multivariate GFs are natural to consider:

- can often store “less nice” single-variable GFs in “nice” multivariate ones,
- can track multiple characteristics at once,
- strict generalization of single-variable GFs.

Same two techniques exist; must use both in conjunction.

Singular set of GF is no longer discrete, now a complex variety.

We also need a **direction**; we will only talk about the **main diagonal** of a generating function: given $F = \sum_{j_1, \dots, j_d} f_{j_1, \dots, j_d} \mathbf{z}^{\mathbf{j}}$ we consider $\Delta F = \sum_n f_{n, \dots, n} t^n$.

We have a specialized version of the Cauchy Integral Theorem.

Theorem (CIT For Main Diagonal)

Suppose that

$$F(t) = \sum_{n \geq 0} f_n t^n = \Delta G(z_1, \dots, z_d).$$

We have a specialized version of the Cauchy Integral Theorem.

Theorem (CIT For Main Diagonal)

Suppose that

$$F(t) = \sum_{n \geq 0} f_n t^n = \Delta G(z_1, \dots, z_d).$$

Then

$$f_n = \frac{1}{(2\pi i)^d} \int_{|z_j|=\alpha_j} G(z_1, \dots, z_d) \frac{dz_1 \cdots dz_d}{z_1^{n+1} \cdots z_d^{n+1}}$$

for α_j sufficiently small for all j .

Solution to singular variety: pick certain points on the variety that are amenable to saddle-point analysis.

Solution to singular variety: pick certain points on the variety that are amenable to saddle-point analysis.

Definition (Minimal Critical Points)

Suppose that $F = G/H$. Then \mathbf{z} is a **smooth critical point** of F if $H_{z_1}(\mathbf{z}) \neq 0$ and it is a solution to the system of equations

$$H(\mathbf{z}) = 0, \quad z_1 H_{z_1}(\mathbf{z}) = \cdots = z_d H_{z_d}(\mathbf{z}).$$

Solution to singular variety: pick certain points on the variety that are amenable to saddle-point analysis.

Definition (Minimal Critical Points)

Suppose that $F = G/H$. Then \mathbf{z} is a **smooth critical point** of F if $H_{z_1}(\mathbf{z}) \neq 0$ and it is a solution to the system of equations

$$H(\mathbf{z}) = 0, \quad z_1 H_{z_1}(\mathbf{z}) = \cdots = z_d H_{z_d}(\mathbf{z}).$$

\mathbf{z} is furthermore **minimal** if there is no \mathbf{w} such that $H(\mathbf{w}) = 0$ and $|w_j| < |z_j|$ for all j .

Solution to singular variety: pick certain points on the variety that are amenable to saddle-point analysis.

Definition (Minimal Critical Points)

Suppose that $F = G/H$. Then \mathbf{z} is a **smooth critical point** of F if $H_{z_1}(\mathbf{z}) \neq 0$ and it is a solution to the system of equations

$$H(\mathbf{z}) = 0, \quad z_1 H_{z_1}(\mathbf{z}) = \cdots = z_d H_{z_d}(\mathbf{z}).$$

\mathbf{z} is furthermore **minimal** if there is no \mathbf{w} such that $H(\mathbf{w}) = 0$ and $|w_j| < |z_j|$ for all j .

Our analysis depends on finding minimal critical points.

Consider

$$F(x, y) = \frac{1}{1 - x - y} = \sum_{j, k \geq 0} \binom{j+k}{j} x^j y^k.$$

Central Binomial Coefficients

Consider

$$F(x, y) = \frac{1}{1 - x - y} = \sum_{j, k \geq 0} \binom{j+k}{j} x^j y^k.$$

Then $V = \{x, y \in \mathbb{C} \mid x + y = 1\}$ and the CP equations are

$$\begin{matrix} 1-x-y=0 \\ -x=-y \end{matrix} \implies \text{unique minimal CP is } (1/2, 1/2).$$

Central Binomial Coefficients

Consider

$$F(x, y) = \frac{1}{1 - x - y} = \sum_{j, k \geq 0} \binom{j+k}{j} x^j y^k.$$

Then $V = \{x, y \in \mathbb{C} \mid x + y = 1\}$ and the CP equations are

$$\begin{matrix} 1-x-y=0 \\ -x=-y \end{matrix} \implies \text{unique minimal CP is } (1/2, 1/2).$$

$$\begin{aligned} \binom{2n}{n} &= \frac{1}{(2\pi i)^2} \int_{|x|=1/2} \int_{|y|=1/2-\epsilon} \frac{1}{1-x-y} \frac{dx dy}{x^{n+1} y^{n+1}} \\ &\sim \frac{1}{2\pi i} \int_{\substack{|x|=1/2 \\ \arg(x) \in (-\pi/4, \pi/4)}} \frac{dx}{x^{n+1} (1-x)^{n+1}} \\ &\sim \frac{4^n}{2\pi} \int_{-\pi/4}^{\pi/4} \frac{1}{1 - e^{it}/2} e^{-n(\log(2 - e^{it}) + it)} dt \sim \frac{4^n}{\sqrt{\pi n}} \end{aligned}$$

Lattice Walk Setup

We can use these techniques to find asymptotics of lattice walk models.

Lattice Walk Setup

We can use these techniques to find asymptotics of lattice walk models.

Definition (Lattice Walk Model)

A d -dimensional **weighted short-step lattice walk model** consists of:

- a **step set** $S \subseteq \{-1, 0, +1\}^d \setminus \{(0, \dots, 0)\}$,
- a **restricting region** $R \subseteq \mathbb{Z}^d$,
- a **starting point** $p \in R$,
- a **terminal set** $T \subseteq R$,
- a **weight** $\omega_s > 0$ for each $s \in S$ (if unspecified $\omega_s = 1$).

Lattice Walk Setup

We can use these techniques to find asymptotics of lattice walk models.

Definition (Lattice Walk Model)

A d -dimensional **weighted short-step lattice walk model** consists of:

- a **step set** $S \subseteq \{-1, 0, +1\}^d \setminus \{(0, \dots, 0)\}$,
- a **restricting region** $R \subseteq \mathbb{Z}^d$,
- a **starting point** $p \in R$,
- a **terminal set** $T \subseteq R$,
- a **weight** $\omega_s > 0$ for each $s \in S$ (if unspecified $\omega_s = 1$).

A walk in this lattice walk model consists of a walk that starts at p , stays in R at all times, ends at T , and takes steps using S .

Lattice Walk Setup

We can use these techniques to find asymptotics of lattice walk models.

Definition (Lattice Walk Model)

A d -dimensional **weighted short-step lattice walk model** consists of:

- a **step set** $S \subseteq \{-1, 0, +1\}^d \setminus \{(0, \dots, 0)\}$,
- a **restricting region** $R \subseteq \mathbb{Z}^d$,
- a **starting point** $p \in R$,
- a **terminal set** $T \subseteq R$,
- a **weight** $\omega_s > 0$ for each $s \in S$ (if unspecified $\omega_s = 1$).

A walk in this lattice walk model consists of a walk that starts at p , stays in R at all times, ends at T , and takes steps using S .

We always let $p = 0$ and $T = R$.

We want a convenient way to store S .

Definition (characteristic polynomial)

The **characteristic polynomial** of a step set S is

$$S(\mathbf{z}) = \sum_{s \in S} \omega_s \mathbf{z}^s = \sum_{s \in S} \omega_s z_1^{s_1} \cdots z_d^{s_d}$$

Notation

We write \bar{z} to mean z^{-1} .

We want a convenient way to store S .

Definition (characteristic polynomial)

The **characteristic polynomial** of a step set S is

$$S(\mathbf{z}) = \sum_{s \in S} \omega_s \mathbf{z}^s = \sum_{s \in S} \omega_s z_1^{s_1} \cdots z_d^{s_d}$$

Notation

We write \bar{z} to mean z^{-1} .

Kreweras Step Set

Let $S = \{(-1, 0), (0, -1), (1, 1)\}$. This is the *Kreweras* step set.



Then $S(x, y) = \bar{x} + \bar{y} + xy$.

What are good choices for restricting regions?

What are good choices for restricting regions?

Most obvious is $R = \mathbb{Z}^d$ (so-called **unrestricted walks**). These are too easy!

What are good choices for restricting regions?

Most obvious is $R = \mathbb{Z}^d$ (so-called **unrestricted walks**). These are too easy!

Also natural is $R = \mathbb{Z}^{d-1} \times \mathbb{N}$ (so-called **half-space walks**). The GFs for these walks are *algebraic* and thus asymptotics are automatic.

What are good choices for restricting regions?

Most obvious is $R = \mathbb{Z}^d$ (so-called **unrestricted walks**). These are too easy!

Also natural is $R = \mathbb{Z}^{d-1} \times \mathbb{N}$ (so-called **half-space walks**). The GFs for these walks are *algebraic* and thus asymptotics are automatic.

Another natural restricting region is $R = \mathbb{N}^d$ (**orthant walks**). We consider this region.

What are good choices for restricting regions?

Most obvious is $R = \mathbb{Z}^d$ (so-called **unrestricted walks**). These are too easy!

Also natural is $R = \mathbb{Z}^{d-1} \times \mathbb{N}$ (so-called **half-space walks**). The GFs for these walks are *algebraic* and thus asymptotics are automatic.

Another natural restricting region is $R = \mathbb{N}^d$ (**orthant walks**). We consider this region.

Note that a d -dimensional orthant walk encodes a *multiqueue system* with d queues: each point of \mathbb{N}^d gives the number of people in each queue.

In the pre-seminar we saw how to use the kernel method to get a GF for orthant walks using NSEW steps.

In the pre-seminar we saw how to use the kernel method to get a GF for orthant walks using NSEW steps.

Let $S = \{(\pm 1, 0), (0, \pm 1)\}$.



Let f_n be the number of orthant walks using S of length n .

In the pre-seminar we saw how to use the kernel method to get a GF for orthant walks using NSEW steps.

Let $S = \{(\pm 1, 0), (0, \pm 1)\}$.



Let f_n be the number of orthant walks using S of length n . Then

$$F(t) = \sum_{n \geq 0} f_n t^n = \Delta \left(\frac{(1+x)(1+y)}{1 - txyS(x,y)} \right).$$

In the pre-seminar we saw how to use the kernel method to get a GF for orthant walks using NSEW steps.

Let $S = \{(\pm 1, 0), (0, \pm 1)\}$.



Let f_n be the number of orthant walks using S of length n . Then

$$F(t) = \sum_{n \geq 0} f_n t^n = \Delta \left(\frac{(1+x)(1+y)}{1 - txyS(x,y)} \right).$$

For arbitrary step sets the argument we used does not generalize.

In the pre-seminar we saw how to use the kernel method to get a GF for orthant walks using NSEW steps.

Let $S = \{(\pm 1, 0), (0, \pm 1)\}$.



Let f_n be the number of orthant walks using S of length n . Then

$$F(t) = \sum_{n \geq 0} f_n t^n = \Delta \left(\frac{(1+x)(1+y)}{1 - txyS(x,y)} \right).$$

For arbitrary step sets the argument we used does not generalize.

If S is **mostly symmetric** it does.

Definition (Symmetric Step Sets)

- A step set S is **mostly symmetric** if

$$S(z_1, \dots, z_d) = S(z_1, \dots, \bar{z}_j, \dots, z_d) \quad 1 \leq j \leq d-1,$$

meaning S is unchanged if flipped over any of the first $d-1$ axes.

Definition (Symmetric Step Sets)

- A step set S is **mostly symmetric** if

$$S(z_1, \dots, z_d) = S(z_1, \dots, \bar{z}_j, \dots, z_d) \quad 1 \leq j \leq d-1,$$

meaning S is unchanged if flipped over any of the first $d-1$ axes. If this is the case we can write

$$S(\mathbf{z}) = \bar{z}_d A(\hat{\mathbf{z}}) + Q(\hat{\mathbf{z}}) + z_d B(\hat{\mathbf{z}})$$

where $\hat{\mathbf{z}} = (z_1, \dots, z_{d-1})$ and A, Q, B are highly symmetric.

Definition (Symmetric Step Sets)

- A step set S is **mostly symmetric** if

$$S(z_1, \dots, z_d) = S(z_1, \dots, \bar{z}_j, \dots, z_d) \quad 1 \leq j \leq d-1,$$

meaning S is unchanged if flipped over any of the first $d-1$ axes. If this is the case we can write

$$S(\mathbf{z}) = \bar{z}_d A(\hat{\mathbf{z}}) + Q(\hat{\mathbf{z}}) + z_d B(\hat{\mathbf{z}})$$

where $\hat{\mathbf{z}} = (z_1, \dots, z_{d-1})$ and A, Q, B are highly symmetric.

- A step set S is **highly symmetric** if it is mostly symmetric and also $S(z_1, \dots, z_d) = S(z_1, \dots, \bar{z}_d)$. This is equivalent to the condition $A(\hat{\mathbf{z}}) = B(\hat{\mathbf{z}})$.

Definition (Symmetric Step Sets)

- A step set S is **mostly symmetric** if

$$S(z_1, \dots, z_d) = S(z_1, \dots, \bar{z}_j, \dots, z_d) \quad 1 \leq j \leq d-1,$$

meaning S is unchanged if flipped over any of the first $d-1$ axes. If this is the case we can write

$$S(\mathbf{z}) = \bar{z}_d A(\hat{\mathbf{z}}) + Q(\hat{\mathbf{z}}) + z_d B(\hat{\mathbf{z}})$$

where $\hat{\mathbf{z}} = (z_1, \dots, z_{d-1})$ and A, Q, B are highly symmetric.

- A step set S is **highly symmetric** if it is mostly symmetric and also $S(z_1, \dots, z_d) = S(z_1, \dots, \bar{z}_d)$. This is equivalent to the condition $A(\hat{\mathbf{z}}) = B(\hat{\mathbf{z}})$.

Notation

We also write $\bar{S}(\mathbf{z}) = z_d A(\hat{\mathbf{z}}) + Q(\hat{\mathbf{z}}) + \bar{z}_d B(\hat{\mathbf{z}})$.

Theorem (Melczer Wilson [MW19])

Let S be a mostly symmetric short-step step set. If f_n is the number of orthant walks of length n using S then

$$F(t) = \sum_{n \geq 0} f_n t^n = \Delta \left(\frac{(1 + z_1) \cdots (1 + z_{d-1}) (B(\hat{\mathbf{z}}) - z_d^2 A(\hat{\mathbf{z}}))}{(1 - z_d) B(\hat{\mathbf{z}}) (1 - tz_1 \cdots z_d \bar{S}(\mathbf{z}))} \right).$$

Theorem (Melczer Wilson [MW19])

Let S be a mostly symmetric short-step set. If f_n is the number of orthant walks of length n using S then

$$F(t) = \sum_{n \geq 0} f_n t^n = \Delta \left(\frac{(1 + z_1) \cdots (1 + z_{d-1}) (B(\hat{\mathbf{z}}) - z_d^2 A(\hat{\mathbf{z}}))}{(1 - z_d) B(\hat{\mathbf{z}}) (1 - t z_1 \cdots z_d \bar{S}(\mathbf{z}))} \right).$$

Proof.

Use the kernel method. □

Theorem (Melczer Wilson [MW19])

Let S be a mostly symmetric short-step step set. If f_n is the number of orthant walks of length n using S then

$$F(t) = \sum_{n \geq 0} f_n t^n = \Delta \left(\frac{(1 + z_1) \cdots (1 + z_{d-1}) (B(\hat{\mathbf{z}}) - z_d^2 A(\hat{\mathbf{z}}))}{(1 - z_d) B(\hat{\mathbf{z}}) (1 - tz_1 \cdots z_d \bar{S}(\mathbf{z}))} \right).$$

Proof.

Use the kernel method. □

There are 4 different cases to consider.

Theorem (Melczer Wilson [MW19])

Let S be a mostly symmetric short-step set. If f_n is the number of orthant walks of length n using S then

$$F(t) = \sum_{n \geq 0} f_n t^n = \Delta \left(\frac{(1 + z_1) \cdots (1 + z_{d-1}) (B(\hat{\mathbf{z}}) - z_d^2 A(\hat{\mathbf{z}}))}{(1 - z_d) B(\hat{\mathbf{z}}) (1 - tz_1 \cdots z_d \bar{S}(\mathbf{z}))} \right).$$

Proof.

Use the kernel method. □

There are 4 different cases to consider.
These will be demonstrated by example.

Highly Symmetric Step Set Asymptotics

In the simplest case where S is highly symmetric, this reduces to

$$F(t) = \sum_{n \geq 0} f_n t^n = \Delta \left(\frac{(1 + z_1) \cdots (1 + z_d)}{1 - tz_1 \cdots z_d \overline{S}(\mathbf{z})} \right).$$

Highly Symmetric Step Set Asymptotics

In the simplest case where S is highly symmetric, this reduces to

$$F(t) = \sum_{n \geq 0} f_n t^n = \Delta \left(\frac{(1 + z_1) \cdots (1 + z_d)}{1 - tz_1 \cdots z_d \bar{S}(\mathbf{z})} \right).$$

NSEW Steps

Consider

$$F(x, y, t) = \frac{G(x, y, t)}{H(x, y, t)} = \frac{(1+x)(1+y)}{1 - txyS(x, y)}$$

where $S(x, y) = x + \bar{x} + y + \bar{y}$.

Highly Symmetric Step Set Asymptotics

In the simplest case where S is highly symmetric, this reduces to

$$F(t) = \sum_{n \geq 0} f_n t^n = \Delta \left(\frac{(1+z_1) \cdots (1+z_d)}{1 - tz_1 \cdots z_d \bar{S}(\mathbf{z})} \right).$$

NSEW Steps

Consider

$$F(x, y, t) = \frac{G(x, y, t)}{H(x, y, t)} = \frac{(1+x)(1+y)}{1 - txyS(x, y)}$$

where $S(x, y) = x + \bar{x} + y + \bar{y}$. This has two minimal CPs: $\sigma_- = (-1, -1, -1/4)$ and $\sigma_+ = (1, 1, 1/4)$. Since G vanishes at $(-1, -1)$ then σ_+ determines dominant asymptotics.

NSWE Steps cont.

CIT implies that

$$\begin{aligned}f_n &\sim \frac{1}{(2\pi i)^3} \int_{\substack{|x|=1 \\ |y|=1}} \left(\int_{|t|=1/4-\epsilon} \frac{(1+x)(1+y)}{1-txyS(x,y)} \frac{dt}{t^{n+1}} \right) \frac{dxdy}{x^{n+1}y^{n+1}} \\&\sim \frac{1}{(2\pi i)^2} \int_N \frac{(1+x)(1+y)}{xy} S(x,y)^n dxdy \\&\sim \frac{4^n}{(2\pi)^2} \int_K 4e^{-n(x^2/4+y^2/4)} dxdy \\&\sim \frac{4}{\pi} \cdot \frac{4^n}{n}.\end{aligned}$$

The techniques in this example generalize to arbitrary highly symmetric step sets.

Theorem (Melczer Mishna 2016 [MM16])

Suppose S is a highly symmetric step set. If s_n is the number of walks of length n using S that remains in \mathbb{N}^d then

$$s_n \sim \left[\frac{S(\mathbf{1})^{d/2}}{\pi^{d/2}(a_1 \cdots a_d)^{1/2}} \right] \cdot \frac{S(\mathbf{1})^n}{n^{d/2}},$$

where a_j is the number of steps in S that have j -th coordinate 1.

The techniques in this example generalize to arbitrary highly symmetric step sets.

Theorem (Melczer Mishna 2016 [MM16])

Suppose S is a highly symmetric step set. If s_n is the number of walks of length n using S that remains in \mathbb{N}^d then

$$s_n \sim \left[\frac{S(\mathbf{1})^{d/2}}{\pi^{d/2}(a_1 \cdots a_d)^{1/2}} \right] \cdot \frac{S(\mathbf{1})^n}{n^{d/2}},$$

where a_j is the number of steps in S that have j -th coordinate 1.

Cardinal Directions

If $S = \{\pm e_1, \dots, \pm e_d\}$ then

$$s_n \sim \left[\frac{(2d)^{d/2}}{\pi^{d/2}} \right] \cdot \frac{(2d)^n}{n^{d/2}}$$

Mostly Symmetric Step Set Asymptotics - Setup

In the mostly symmetric case we don't have $B = A$ and so can't cancel.

Mostly Symmetric Step Set Asymptotics - Setup

In the mostly symmetric case we don't have $B = A$ and so can't cancel. Furthermore in the expression

$$\frac{(1 + z_1) \cdots (1 + z_{d-1}) (B(\hat{\mathbf{z}}) - z_d^2 A(\hat{\mathbf{z}}))}{(1 - z_d) B(\hat{\mathbf{z}}) (1 - tz_1 \cdots z_d \overline{S}(\mathbf{z}))}$$

the factor of B in the denominator behaves uncontrollably.

Mostly Symmetric Step Set Asymptotics - Setup

In the mostly symmetric case we don't have $B = A$ and so can't cancel. Furthermore in the expression

$$\frac{(1 + z_1) \cdots (1 + z_{d-1}) (B(\hat{\mathbf{z}}) - z_d^2 A(\hat{\mathbf{z}}))}{(1 - z_d) B(\hat{\mathbf{z}}) (1 - tz_1 \cdots z_d \bar{S}(\mathbf{z}))}$$

the factor of B in the denominator behaves uncontrollably. Instead we write $F(t)$ as the diagonal of

$$\frac{(1 + z_1) \cdots (1 + z_{d-1}) (1 - tz_1 \cdots z_d (Q(\hat{\mathbf{z}}) + 2z_d A(\hat{\mathbf{z}})))}{(1 - z_d) (1 - tz_1 \cdots z_d \bar{S}(\mathbf{z})) (1 - tz_1 \cdots z_d (Q(\hat{\mathbf{z}}) + z_d A(\hat{\mathbf{z}})))}.$$

Mostly Symmetric Step Set Asymptotics - Setup

In the mostly symmetric case we don't have $B = A$ and so can't cancel. Furthermore in the expression

$$\frac{(1 + z_1) \cdots (1 + z_{d-1}) (B(\hat{\mathbf{z}}) - z_d^2 A(\hat{\mathbf{z}}))}{(1 - z_d) B(\hat{\mathbf{z}}) (1 - tz_1 \cdots z_d \bar{S}(\mathbf{z}))}$$

the factor of B in the denominator behaves uncontrollably. Instead we write $F(t)$ as the diagonal of

$$\frac{(1 + z_1) \cdots (1 + z_{d-1}) (1 - tz_1 \cdots z_d (Q(\hat{\mathbf{z}}) + 2z_d A(\hat{\mathbf{z}})))}{(1 - z_d) (1 - tz_1 \cdots z_d \bar{S}(\mathbf{z})) (1 - tz_1 \cdots z_d (Q(\hat{\mathbf{z}}) + z_d A(\hat{\mathbf{z}})))}.$$

This looks awful, but it's easier to deal with the denominator.

We have one last important distinction to make.

We have one last important distinction to make.

Definition

Let S be a mostly symmetric (and not highly symmetric) step set.

We have one last important distinction to make.

Definition

Let S be a mostly symmetric (and not highly symmetric) step set.

- S has **negative drift** if $A(\mathbf{1}) > B(\mathbf{1})$,

We have one last important distinction to make.

Definition

Let S be a mostly symmetric (and not highly symmetric) step set.

- S has **negative drift** if $A(\mathbf{1}) > B(\mathbf{1})$,
- S has **positive drift** if $A(\mathbf{1}) < B(\mathbf{1})$,

We have one last important distinction to make.

Definition

Let S be a mostly symmetric (and not highly symmetric) step set.

- S has **negative drift** if $A(1) > B(1)$,
- S has **positive drift** if $A(1) < B(1)$,
- S has **zero drift** if $A(1) = B(1)$.

We have one last important distinction to make.

Definition

Let S be a mostly symmetric (and not highly symmetric) step set.

- S has **negative drift** if $A(\mathbf{1}) > B(\mathbf{1})$,
- S has **positive drift** if $A(\mathbf{1}) < B(\mathbf{1})$,
- S has **zero drift** if $A(\mathbf{1}) = B(\mathbf{1})$.

We can view this as a condition on the vector sum $\sum \omega_s s \in \mathbb{Z}$. If this sum is positive S has positive drift etc.

We have one last important distinction to make.

Definition

Let S be a mostly symmetric (and not highly symmetric) step set.

- S has **negative drift** if $A(\mathbf{1}) > B(\mathbf{1})$,
- S has **positive drift** if $A(\mathbf{1}) < B(\mathbf{1})$,
- S has **zero drift** if $A(\mathbf{1}) = B(\mathbf{1})$.

We can view this as a condition on the vector sum $\sum \omega_s s \in \mathbb{Z}$. If this sum is positive S has positive drift etc.

In each of the different cases the singular variety is differently positioned, and so the asymptotics change.

Mostly Symmetric Step Set Asymptotics - Negative Drift

The second simplest case is negative drift mostly symmetric.

Negative Drift

Let $S = \{(-1, -1), (1, -1), (0, 1)\}$. Then the GF is the diagonal of

$$\frac{(1+x)(1-2t(x^2y^2+1))}{(1-y)(1-t(x^2y^2+y^2+x))(1-t(x^2y^2+1))}.$$

Mostly Symmetric Step Set Asymptotics - Negative Drift

The second simplest case is negative drift mostly symmetric.

Negative Drift

Let $S = \{(-1, -1), (1, -1), (0, 1)\}$. Then the GF is the diagonal of

$$\frac{(1+x)(1-2t(x^2y^2+1))}{(1-y)(1-t(x^2y^2+y^2+x))(1-t(x^2y^2+1))}.$$

There are now four minimal critical points

$$(1, 1/\sqrt{2}, 1/2), (1, -1/\sqrt{2}, 1/2), (-1, \pm i/\sqrt{2}, -1/2).$$

The numerator vanishes at the last two CPs, implying that the first two give dominant asymptotics. Only one factor of the denominator vanishes (the “smooth” case).

Mostly Symmetric Step Set Asymptotics - Negative Drift

The second simplest case is negative drift mostly symmetric.

Negative Drift

Let $S = \{(-1, -1), (1, -1), (0, 1)\}$. Then the GF is the diagonal of

$$\frac{(1+x)(1-2t(x^2y^2+1))}{(1-y)(1-t(x^2y^2+y^2+x))(1-t(x^2y^2+1))}.$$

There are now four minimal critical points

$$(1, 1/\sqrt{2}, 1/2), (1, -1/\sqrt{2}, 1/2), (-1, \pm i/\sqrt{2}, -1/2).$$

The numerator vanishes at the last two CPs, implying that the first two give dominant asymptotics. Only one factor of the denominator vanishes (the “smooth” case). We have

$$s_n \sim \frac{16 + 12\sqrt{2}}{\pi} \cdot \frac{(2\sqrt{2})^n}{n^2} + \frac{-16 + 12\sqrt{2}}{\pi} \cdot \frac{(-2\sqrt{2})^n}{n^2}$$

In general we have the following.

Theorem (Melczer Wilson [MW19])

If S is mostly symmetric with negative drift and $Q(z) \neq 0$ then

$$s_n \sim C_\rho \cdot \frac{S(1, \rho)^n}{n^{d/2+1}}.$$

If S is mostly symmetric with negative drift and $Q(z) = 0$ then

$$s_n \sim C_\rho \cdot \frac{S(1, \rho)^n}{n^{d/2+1}} + C_{-\rho} \cdot \frac{S(1, -\rho)^n}{n^{d/2+1}}.$$

Where ρ , C_ρ , and $C_{-\rho}$ are explicit constants.

Mostly Symmetric Step Set Asymptotics - Positive Drift

In the positive drift case we encounter CPs that lie on the intersection of two factors of the denominator.

Positive Drift

Let $S = \{(-1, 1), (1, 1), (0, -1)\}$. Then the GF is the diagonal of

$$F(x, y, t) = \frac{(1+x)(1-2txy^2)}{(1-y)(1-t(xy^2+x^2+1))(1-txy^2)}.$$

Mostly Symmetric Step Set Asymptotics - Positive Drift

In the positive drift case we encounter CPs that lie on the intersection of two factors of the denominator.

Positive Drift

Let $S = \{(-1, 1), (1, 1), (0, -1)\}$. Then the GF is the diagonal of

$$F(x, y, t) = \frac{(1+x)(1-2txy^2)}{(1-y)(1-t(xy^2+x^2+1))(1-txy^2)}.$$

The only minimal CP is $(1, 1, 1/3)$. Two terms in the denominator vanish at this point! This means we are no longer in the “smooth” case. Since the main diagonal is “generic” we are able to take two residues.

Positive Drift cont.

$$\begin{aligned} s_n &\sim \frac{1}{(2\pi i)^3} \int_{|x|=1} \left(\int_{\substack{|y|=1-\epsilon \\ |t|=1/3-\epsilon}} F(x, y, t) \frac{dy dt}{y^{n+1} t^{n+1}} \right) \frac{dx}{x^{n+1}} \\ &\sim \frac{1}{2\pi i} \int_N \frac{(x^2 - x + 1)(1 + x)}{x(x^2 + 1)} (x + 1 + 1/x)^n dx \\ &\sim \frac{\sqrt{3}}{2\sqrt{\pi}} \cdot \frac{3^n}{\sqrt{n}}. \end{aligned}$$

Theorem (Melczer Wilson [MW19])

If S is mostly symmetric with positive drift then

$$s_n \sim \left[\left(1 - \frac{A(1)}{B(1)} \right) \frac{S(1)^{d/2}}{(2\pi)^{d/2}} \cdot \frac{1}{(a_1 \cdots a_d)^{1/2}} \right] \cdot \frac{S(1)^n}{n^{d/2-1/2}}$$

Mostly Symmetric Step Set Asymptotics - Zero Drift

In the zero drift case $A(\mathbf{1}) = B(\mathbf{1})$; we expect the numerator to vanish when $\mathbf{z} = \mathbf{1}$.

Zero Drift

Let $S = \{(-1, -1), (1, -1), 2 \cdot (0, 1)\}$. Then the GF is the diagonal of

$$F(x, y, t) = \frac{(1+x)(1-2ty^2(x^2+1))}{(1-y)(1-txy(2/y+xy+y/x))(1-ty^2(x^2+1))}.$$

Mostly Symmetric Step Set Asymptotics - Zero Drift

In the zero drift case $A(\mathbf{1}) = B(\mathbf{1})$; we expect the numerator to vanish when $\mathbf{z} = \mathbf{1}$.

Zero Drift

Let $S = \{(-1, -1), (1, -1), 2 \cdot (0, 1)\}$. Then the GF is the diagonal of

$$F(x, y, t) = \frac{(1+x)(1-2ty^2(x^2+1))}{(1-y)(1-txy(2/y+xy+y/x))(1-ty^2(x^2+1))}.$$

There are four minimal critical points

$$(1, 1, 1/4), (1, -1, -1/4), (-1, \pm i, \mp i).$$

Mostly Symmetric Step Set Asymptotics - Zero Drift

In the zero drift case $A(\mathbf{1}) = B(\mathbf{1})$; we expect the numerator to vanish when $\mathbf{z} = \mathbf{1}$.

Zero Drift

Let $S = \{(-1, -1), (1, -1), 2 \cdot (0, 1)\}$. Then the GF is the diagonal of

$$F(x, y, t) = \frac{(1+x)(1-2ty^2(x^2+1))}{(1-y)(1-txy(2/y+xy+y/x))(1-ty^2(x^2+1))}.$$

There are four minimal critical points

$$(1, 1, 1/4), (1, -1, -1/4), (-1, \pm i, \mp i).$$

It turns out the contribution from the last three CPs is negligible, and only the first matters. Two terms in the denominator vanish, but this time the main diagonal is “non-generic”, and so we are only able to take one residue.

Zero Drift cont.

$$\begin{aligned}
 s_n &\sim \frac{1}{(2\pi i)^3} \int_{|x|=1} \left(\int_{\substack{|y|=1-\epsilon \\ |t|=1/4-\epsilon}} F(x, y, t) \frac{dy dt}{y^{n+1} t^{n+1}} \right) \frac{dx}{x^{n+1}} \\
 &\sim \frac{1}{(2\pi i)^2} \int_{N(1)} \int_{N(1-\epsilon)} \frac{1+x}{2x^2 y} \cdot \frac{2x - y^2(1+x^2)}{1-y} \bar{S}(x, y)^n dx dy.
 \end{aligned}$$

$$\begin{aligned}
s_n &\sim \frac{1}{(2\pi i)^3} \int_{|x|=1} \left(\int_{\substack{|y|=1-\epsilon \\ |t|=1/4-\epsilon}} F(x, y, t) \frac{dy dt}{y^{n+1} t^{n+1}} \right) \frac{dx}{x^{n+1}} \\
&\sim \frac{1}{(2\pi i)^2} \int_{N(1)} \int_{N(1-\epsilon)} \frac{1+x}{2x^2 y} \cdot \frac{2x - y^2(1+x^2)}{1-y} \bar{S}(x, y)^n dx dy.
\end{aligned}$$

We cannot take any more residues, and must use saddle-point techniques. Note that at $x = y = 1$ both the numerator and denominator of $P(x, y) = \frac{2x - y^2(1+x^2)}{1-y}$ vanish.

$$\begin{aligned}
s_n &\sim \frac{1}{(2\pi i)^3} \int_{|x|=1} \left(\int_{\substack{|y|=1-\epsilon \\ |t|=1/4-\epsilon}} F(x, y, t) \frac{dy dt}{y^{n+1} t^{n+1}} \right) \frac{dx}{x^{n+1}} \\
&\sim \frac{1}{(2\pi i)^2} \int_{N(1)} \int_{N(1-\epsilon)} \frac{1+x}{2x^2 y} \cdot \frac{2x - y^2(1+x^2)}{1-y} \bar{S}(x, y)^n dx dy.
\end{aligned}$$

We cannot take any more residues, and must use saddle-point techniques. Note that at $x = y = 1$ both the numerator and denominator of $P(x, y) = \frac{2x - y^2(1+x^2)}{1-y}$ vanish. One can carefully show that $P(x, y)$ has leading term equal to 4, with other terms contributing a negligible amount to asymptotics.

$$\begin{aligned}
 s_n &\sim \frac{1}{(2\pi i)^3} \int_{|x|=1} \left(\int_{\substack{|y|=1-\epsilon \\ |t|=1/4-\epsilon}} F(x, y, t) \frac{dy dt}{y^{n+1} t^{n+1}} \right) \frac{dx}{x^{n+1}} \\
 &\sim \frac{1}{(2\pi i)^2} \int_{N(1)} \int_{N(1-\epsilon)} \frac{1+x}{2x^2 y} \cdot \frac{2x - y^2(1+x^2)}{1-y} \bar{S}(x, y)^n dx dy.
 \end{aligned}$$

We cannot take any more residues, and must use saddle-point techniques. Note that at $x = y = 1$ both the numerator and denominator of $P(x, y) = \frac{2x - y^2(1+x^2)}{1-y}$ vanish. One can carefully show that $P(x, y)$ has leading term equal to 4, with other terms contributing a negligible amount to asymptotics. Thus

$$s_n \sim \frac{4}{(2\pi i)^2} \int_N \frac{1+x}{2x^2 y} \bar{S}(x, y)^n dx dy \sim \frac{4^n}{n} \cdot \frac{2\sqrt{2}}{\pi}.$$

Dealing with this tricky term gives us the following result.

Dealing with this tricky term gives us the following result.

Theorem (K. Melczer [KM24])

Suppose S is a mostly symmetric step set with zero drift. Let s_n be the number of walks of length n using S that remain in \mathbb{N}^d . Then

$$s_n \sim \left[\frac{|S|^{d/2}}{\pi^{d/2} (a_1 \cdots a_d)^{1/2}} \right] \cdot \frac{|S|^n}{n^{d/2}}$$

where a_j is the number of steps in S that have j -th coordinate 1.

Higher Order Terms

We can find the dominant term for zero drift walks. What about higher order terms?

Higher Order Terms

We can find the dominant term for zero drift walks. What about higher order terms?

Need to get more terms in the expansion of $P(x, y)$.

Higher Order Terms

We can find the dominant term for zero drift walks. What about higher order terms?

Need to get more terms in the expansion of $P(x, y)$.

One can show

$$\begin{aligned} & \frac{1}{(2\pi i)^2} \int_{N(1)} \int_{N(1-\epsilon)} \frac{1+x}{2x^2y} \cdot \frac{2x - y^2(1+x^2)}{1-y} \bar{S}(x, y)^n dx dy \\ & \sim \frac{1}{(2\pi)^2} \int_{K+i(0,\epsilon)} \left(4 + \frac{is^2}{t} \right) e^{-n(s^2/4 + t^2/2)} ds dt \\ & \sim \frac{4^n}{n} \cdot \frac{2\sqrt{2}}{\pi} + \frac{4^n}{n^{3/2}} \cdot \frac{1}{\sqrt{\pi}}. \end{aligned}$$

Conclusion

To find the higher order asymptotic terms of f_n we need to be able to evaluate integrals of the form

$$\int_{K+i\epsilon} \frac{A(\mathbf{z})}{\mathbf{z}^{\mathbf{k}}} \exp[-n\phi(\mathbf{z})] d\mathbf{z}.$$

Conclusion

To find the higher order asymptotic terms of f_n we need to be able to evaluate integrals of the form

$$\int_{K+i\epsilon} \frac{A(\mathbf{z})}{\mathbf{z}^{\mathbf{k}}} \exp[-n\phi(\mathbf{z})] d\mathbf{z}.$$

This is a whole other can of worms.

Conclusion

To find the higher order asymptotic terms of f_n we need to be able to evaluate integrals of the form

$$\int_{K+i\epsilon} \frac{A(z)}{z^k} \exp[-n\phi(z)] dz.$$

This is a whole other can of worms.

Fin.



Alexander Kroitor and Stephen Melczer.

Completing the asymptotic classification of mostly symmetric short step walks in an orthant, 2024.



Stephen Melczer.

An Invitation to Analytic Combinatorics.

Springer Nature Switzerland, Cham, Switzerland.



Stephen Melczer and Marni Mishna.

Asymptotic lattice path enumeration using diagonals.

Algorithmica, 75(4):782–811, 2016.



Stephen Melczer and Mark C. Wilson.

Higher dimensional lattice walks: Connecting combinatorial and analytic behavior.

SIAM J. Discrete Math., 33(4):2140–2174, 2019.