

Quasar-Lyman α forest cross-correlation from BOSS DR11: Baryon Acoustic Oscillations

This content has been downloaded from IOPscience. Please scroll down to see the full text.

JCAP05(2014)027

(<http://iopscience.iop.org/1475-7516/2014/05/027>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 136.152.142.26

This content was downloaded on 07/12/2015 at 19:37

Please note that [terms and conditions apply](#).

Quasar-Lyman α forest cross-correlation from BOSS DR11: Baryon Acoustic Oscillations

Andreu Font-Ribera,^{a,b} David Kirkby,^c Nicolas Busca,^d
Jordi Miralda-Escudé,^{e,f} Nicholas P. Ross,^{b,g} Anže Slosar,^h
James Rich,^l Éric Aubourg,^d Stephen Bailey,^b Vaishali Bhardwaj,^{i,b}
Julian Bautista,^d Florian Beutler,^b Dmitry Bizyaev,ⁱ
Michael Blomqvist,^c Howard Brewington,^j Jon Brinkmann,^j
Joel R. Brownstein,^k Bill Carithers,^b Kyle S. Dawson,^k
Timothée Delubac,^l Garrett Ebelke,^j Daniel J. Eisenstein,^m
Jian Ge,ⁿ Karen Kinemuchi,ⁱ Khee-Gan Lee,^o
Viktor Malanushenko,^j Elena Malanushenko,^j Moses Marchante,ⁱ
Daniel Margala,^c Demitri Muna,^p Adam D. Myers,^q
Pasquier Noterdaeme,^r Daniel Oravetz,ⁱ
Nathalie Palanque-Delabrouille,^l Isabelle Pâris,^s Patrick Petitjean,^r
Matthew M. Pieri,^t Graziano Rossi,^l Donald P. Schneider,^{u,v}
Audrey Simmons,ⁱ Matteo Viel,^{w,x} Christophe Yèche^l and
Donald G. York^y

^aInstitute of Theoretical Physics, University of Zurich,
Winterthurerstrasse 190, 8057 Zurich, Switzerland

^bLawrence Berkeley National Laboratory,
1 Cyclotron Road, Berkeley, CA, U.S.A.

^cDepartment of Physics and Astronomy, University of California,
4129 Frederick Reines Hall, Irvine, CA, 92697, U.S.A.

^dAPC, Université Paris Diderot-Paris 7, CNRS/IN2P3, CEA, Observatoire de Paris,
10, rue A. Domon & L. Duquet, Paris, France

^eInstitut de Ciències del Cosmos (IEEC/UB),
Martí i Franquès 1, Barcelona, 08028 Catalonia

^fInstitució Catalana de Recerca i Estudis Avançats,
Passeig Lluís Companys 23, Barcelona, 08010 Catalonia

^gDepartment of Physics, Drexel University,
3141 Chestnut Street, Philadelphia, PA, 19104 U.S.A.

^hBrookhaven National Laboratory,
Blgd 510, Upton, NY, 11375 U.S.A.

ⁱDepartment of Astronomy, University of Washington,
Box 351580, Seattle, WA, 98195 U.S.A.

^jApache Point Observatory and New Mexico State University,
P.O. Box 59, Sunspot, NM, 88349-0059 U.S.A.

^kDepartment of Physics and Astronomy, University of Utah,
115 S 1400 E, Salt Lake City, UT, 84112 U.S.A.

^lCEA, Centre de Saclay, IRFU,
Gif-sur-Yvette, 91191 France

^mHarvard-Smithsonian Center for Astrophysics,
60 Garden Street, Cambridge, MA, 02138 U.S.A.

ⁿAstronomy Department, University of Florida,
211 Bryant Space Science Center, Gainesville, FL, 32611-2055 U.S.A.

^oMax-Planck-Institut für Astronomie,
Königstuhl 17, Heidelberg, D-69117 Germany

^pDepartment of Astronomy, Ohio State University,
Columbus, OH, 43210 U.S.A.

^qDepartment of Physics and Astronomy 3905, University of Wyoming,
1000 East University, Laramie, WY, 82071 U.S.A.

^rInstitut d'Astrophysique de Paris, CNRS-UPMC, UMR7095,
98bis bd Arago, Paris, 75014 France

^sDepartamento de Astronomía, Universidad de Chile,
Casilla 36-D, Santiago, Chile

^tInstitute of Cosmology and Gravitation, Dennis Sciama Building, University of Portsmouth,
Portsmouth, PO1 3FX U.K.

^uDepartment of Astronomy and Astrophysics, The Pennsylvania State University,
University Park, PA, 16802 U.S.A.

^vInstitute for Gravitation and the Cosmos, The Pennsylvania State University,
University Park, PA, 16802 U.S.A.

^wINAF, Osservatorio Astronomico di Trieste,
Via G.B. Tiepolo 11, Trieste, 34131 Italy

^xINFN/National Institute for Nuclear Physics,
Via Valerio 2, Trieste, I-34127 Italy

^yDepartment of Astronomy and Astrophysics and The Fermi Institute,
The University of Chicago, 5640 South Ellis Avenue, Chicago, IL, 60615 U.S.A.

E-mail: font@physik.uzh.ch

Received November 5, 2013

Revised January 24, 2014

Accepted April 22, 2014

Published May 20, 2014

Abstract. We measure the large-scale cross-correlation of quasars with the Ly α forest absorption, using over 164,000 quasars from Data Release 11 of the SDSS-III Baryon Oscillation Spectroscopic Survey. We extend the previous study of roughly 60,000 quasars from Data Release 9 to larger separations, allowing a measurement of the Baryonic Acoustic Oscillation

(BAO) scale along the line of sight $c/(H(z = 2.36) r_s) = 9.0 \pm 0.3$ and across the line of sight $D_A(z = 2.36) / r_s = 10.8 \pm 0.4$, consistent with CMB and other BAO data. Using the best fit value of the sound horizon from Planck data ($r_s = 147.49 \text{ Mpc}$), we can translate these results to a measurement of the Hubble parameter of $H(z = 2.36) = 226 \pm 8 \text{ km s}^{-1} \text{ Mpc}^{-1}$ and of the angular diameter distance of $D_A(z = 2.36) = 1590 \pm 60 \text{ Mpc}$. The measured cross-correlation function and an update of the code to fit the BAO scale (baofit) are made publicly available.

Keywords: Lyman alpha forest, baryon acoustic oscillations, redshift surveys, dark energy experiments

ArXiv ePrint: [1311.1767](https://arxiv.org/abs/1311.1767)

Contents

1	Introduction	1
2	Data sample	3
2.1	Quasar sample	3
2.2	$\text{Ly}\alpha$	3
2.3	Independent sub-samples	4
3	Cross-correlation	4
3.1	Continuum fitting	4
3.2	From flux to δ_F	5
3.3	Estimator and covariance matrix	6
3.4	Measured cross-correlation	7
4	Fitting the BAO scale	8
4.1	BAO model	8
4.1.1	Theoretical model for the cross-correlation	9
4.1.2	Quasar redshift errors	9
4.1.3	Broadband distortion	10
4.2	BAO fits	10
4.3	Systematic tests	11
4.4	Test of the covariance matrix	11
4.5	Alternative uncertainty estimates of the BAO scales	13
4.6	Visualizing the BAO peak	13
5	Discussion & conclusions	15
5.1	$\text{Ly}\alpha$ auto-correlation vs. quasar- $\text{Ly}\alpha$ cross-correlation	16
A	Public access to data and code	18
B	Fisher matrix forecasts	18
B.1	Auto-correlation	18
B.2	Cross-correlation	19
B.3	Forecast for a BOSS-like survey	19

1 Introduction

Fifteen years ago, two independent studies of the luminosity distance of type Ia supernovae ([1, 2]) showed that the Universe was undergoing an accelerated expansion. In order to explain such an unintuitive result, different authors have suggested the need for a cosmological constant in Einstein’s equations of general relativity, more profound modifications of the gravitational theory, or the presence of a new energy component usually referred to as *dark energy*.

Following this discovery, different cosmological probes have provided a wealth of new data, allowing us to constrain the cosmological parameters of the model at a few-percent

level. The simplest possible solution, a flat universe with a cosmological constant, is able to explain all current data [3], and ongoing and future cosmological surveys will continue to reduce the errorbars of these measurements and place even more stringent constraints on the models.

There are different observational probes that can measure the history of the accelerated expansion (see [4] for a review). One is the measurement of the Baryon Acoustic Oscillation (BAO) scale on the clustering of any tracer of the density field, which can be used as a cosmic ruler to study the geometry of the Universe [5]. This technique has gained considerable attention during the last decade, and the list of BAO measurements is rapidly increasing.

In theory, any tracer of the large-scale matter distribution can be used to measure BAO. Even though the first measurements came from the clustering of low redshift galaxies (Sloan Digital Sky Survey [6] at $0.2 < z < 0.4$, Two-degree-Field Galaxy Redshift Survey [7] at $0.1 < z < 0.2$), and the tightest constraints are obtained from intermediate redshift galaxy surveys (WiggleZ Dark Energy Survey [8] at $0.4 < z < 0.8$, Baryon Oscillation Spectroscopic Survey [9] at $0.4 < z < 0.7$), there are also a variety of undergoing or planned surveys that aim to measure BAO at higher redshift from the clustering of x-ray sources (eROSITA [10]), 21cm emission (Canadian Hydrogen Intensity Mapping Experiment,¹ Baryon Acoustic Oscillation Broadband and Broad-beam Array [11]), Ly α emitting galaxies (Hobby-Eberly Telescope Dark Energy Experiment [12], Dark Energy Spectroscopic Instrument [13]) and quasars (Extended Baryon Oscillation Spectroscopic Survey [14]).

The spatial distribution of neutral hydrogen, as traced by the Lyman- α forest (Ly α forest) can also be used to measure BAO. The first measurement of the three-dimensional large-scale structure of Ly α absorption was presented in [15], using over 14,000 spectra from the first year of BOSS. This study was extended using approximately 50,000 quasar spectra from the ninth data release of SDSS (DR9, [16]), and the first detections of the BAO at $z = 2.4$ were presented in [17, 18] and [19].

Using the same set of spectra, [20] presented an analysis of the large scale cross-correlation of quasars and the Ly α absorption. In this analysis the cross-correlation was clearly detected up to separations of $r \sim 70 h^{-1}$ Mpc, and was accurately described by a linear bias and redshift-space distortion theory for comoving separations $r > 15 h^{-1}$ Mpc. Their measurement of the quasar bias of $b_q = 3.64 \pm 0.14$ was fully consistent with measurements of the quasar auto-correlation function at the same redshift, e.g., [21].

In this paper, we use over 164,000 quasars from the eleventh data release of SDSS (DR11, which will be publicly released at the end of 2014 together with DR12) to extend the measurement of the cross-correlation to larger separations, and we present an accurate determination of the BAO scale in cross-correlation at high-redshift ($z \sim 2.4$).

Throughout, we use the fiducial cosmology ($\Omega_m = 0.27$, $\omega_b = 0.0227$, $h = 0.7$, $\sigma_8 = 0.8$, $n_s = 0.97$) that was used in the previous Ly α BAO measurements ([17–19]). We use the publicly available code CAMB [22] to compute the comoving distance to the sound horizon at the redshift at which baryon-drag optical depth equals unity, $z_{\text{drag}} = 1059.97$, and obtain a value of $r_s = 149.72$ Mpc. Some previous BAO studies have used the equations from [23] to compute the value of r_s , resulting in a few percent difference with respect to the value computed with CAMB, as discussed in [3].

We start by introducing our data sample in section 2. In section 3 we present our measurement of the quasar-Ly α cross-correlation and summarize our analysis method. In

¹<http://chime.phas.ubc.ca/>.

section 4 we describe our fits of the BAO scale, present our main results, and test possible systematic effects. In section 5 we compare our results with previous BAO measurements at similar redshifts.

2 Data sample

In this section we describe the data set used in this study, and present a series of references for further details.

The eleventh Data Release (DR11) of the SDSS-III Collaboration ([24]) contains all spectra obtained during the first four years of the Baryon Oscillation Spectroscopic Survey (BOSS, [30]), including spectra of 238,978 visually confirmed quasars. The quasar target selection used in BOSS is summarized in [31], and combines different targeting methods described in [32, 33], and [34].

In this study we measure the cross-correlation of two tracers of the underlying density field: the number density of quasars and the $\text{Ly}\alpha$ absorption along a set of lines of sight. We will use the term “quasar sample” to refer to the quasars used as tracers of the density field, and the term “ $\text{Ly}\alpha$ sample” to refer to those quasar lines of sight where the $\text{Ly}\alpha$ absorption is measured.

2.1 Quasar sample

We use a preliminary version of the DR11Q quasar catalog, an updated version of the DR9Q catalog presented in [35]. This catalog contains a total of 238,978 visually confirmed quasars, distributed in an area of 8976 square degrees in two disconnected parts of the sky: the South Galactic Cap (SGC) and the North Galactic Cap (NGC). To avoid repeated observations of the same object, we only use quasars in the catalog that have the SPECPRIMARY flag [30].

The performance of the BOSS spectrograph rapidly deteriorates at wavelengths bluer than $\lambda \sim 3650 \text{ \AA}$, corresponding to a $\text{Ly}\alpha$ absorption of $z = 2$; this wavelength sets the lower limit in the redshift range used in this study. Since the number of identified quasars drops rapidly at high redshift, we restrict our quasar sample to those with redshifts in the range $2.0 \leq z_q \leq 3.5$. This redshift constraint reduces the number of quasars used as tracers of the density field to 164,017. In this paper we use the Z_{VI} redshift estimate from [35].

2.2 $\text{Ly}\alpha$ sample

Not all spectra from the quasar sample are included in our $\text{Ly}\alpha$ sample. We first drop the spectra from quasars with a redshift lower than $z_q = 2.15$, since only a small part of their $\text{Ly}\alpha$ forest can be observed with the BOSS spectrograph. This choice reduces the number of spectra to 153,496.

During the visual inspection, Broad Absorption Line quasars (BAL) are identified [35]. We discard any spectra from BAL quasars, reducing the number of spectra to 136,431. We finally exclude spectra with less than 150 pixels covering the $\text{Ly}\alpha$ forest, further reducing the number of spectra used in the $\text{Ly}\alpha$ sample to 130,825. We use the same definition of the $\text{Ly}\alpha$ forest as in [36], which contains all pixels in the rest-frame wavelength range $1040 \text{ \AA} \leq \lambda_{\text{rest}} \leq 1200 \text{ \AA}$.

In the left panel of figure 1 we show the redshift distribution of objects in the quasar sample (red in the NGC, blue in the SGC) and the distribution of these quasars that have spectra included in the $\text{Ly}\alpha$ sample (green in the NGC, purple in the SGC).




Figure 1. Left panel: redshift distribution of the 164,017 quasars used as density tracers (red in NGC, blue in SGC), and of the 130,825 quasars with Ly α spectra (green in NGC, purple in SGC). Right panel: DR11 footprint in J2000 equatorial coordinates with the 66 sub-samples indicated in different colors.

We use an updated (DR11) version of the Damped Lyman α system (DLA) catalogue from [37], based on DLA profile recognition as described in [38], to mask the central part of DLA in the spectra (up to a transmitted flux fraction of $F < 0.8$), and correct the rest of the spectra using the inferred Voigt profile.

We correct the noise estimate from the pipeline using the method described in [17]. Following the same reference, we rebin our spectra by averaging the flux over three adjacent pipeline pixels. These new pixels have a width of 210 km s^{-1} , $\sim 2 \text{ } h^{-1} \text{ Mpc}$ at the redshift of interest, much smaller than the minimum separation in which we are interested ($r > 40 \text{ } h^{-1} \text{ Mpc}$), and smaller than the width of the BAO peak ($\Delta r \sim 25 \text{ } h^{-1} \text{ Mpc}$). We will use the term “pixel” to refer to these rebinned pixels.

2.3 Independent sub-samples

In section 3 we explain the method to estimate the covariance matrix of our measurement. To reduce the required computing time, the survey is split into 66 sub-samples with a similar number of quasars and combine the measurement of each sub-sample assuming that they are independent. The right panel of figure 1 shows the different sub-samples, 51 of them in the north galactic cap (NGC) and 15 in the south (SGC).

These sub-samples are also used in section 4 to compute bootstrap errors on the best fit parameters. Since we are interested in scales smaller than $150 \text{ } h^{-1} \text{ Mpc}$ and the typical area of these sub-samples is roughly $(800 \text{ } h^{-1} \text{ Mpc})^2$ (140 deg^2 at $z = 2.4$), the assumption that the sub-samples are independent is justified.

3 Cross-correlation

In this section we briefly describe the method used to measure the cross-correlation and its covariance matrix, referring the reader to previous publications for a detailed explanation ([20, 39]), and present the measured cross-correlation.

3.1 Continuum fitting

The first step necessary to estimate the Ly α transmitted flux fraction $F(\lambda) = e^{-\tau(\lambda)}$ from a set of pixels with flux $f(\lambda)$ is estimating the quasar continuum, $C_q(\lambda)$,

$$F(\lambda) = \frac{f(\lambda)}{C_q(\lambda)}. \quad (3.1)$$

	β_F	b_q	a_{peak}	α	α_{\parallel}	α_{\perp}	χ^2 (d.o.f)
FID	1.09 ± 0.29	3.02 ± 0.22	-	-	1.042 ± 0.034	0.930 ± 0.036	426.4 (420)
ISO	1.09 ± 0.30	3.00 ± 0.22	-	0.988 ± 0.022	-	-	429.5 (421)
NW	1.13 ± 0.34	2.71 ± 0.21	-	-	-	-	448.5 (422)
AMP	1.14 ± 0.32	3.02 ± 0.22	1.15 ± 0.24	-	1.041 ± 0.033	0.933 ± 0.035	426.0 (419)
PCA	1.54 ± 0.42	3.14 ± 0.22	-	-	1.052 ± 0.035	0.922 ± 0.044	474.7 (420)

Table 1. Best fit parameters for different analyses: fiducial BAO fit (FID), isotropic BAO fit (ISO), non-BAO fit (NW), fit with free amplitude (AMP) and using a continuum fitting method based on a Principal Component Analysis (PCA, [40]).

Among various approaches for the determination of the continuum available in the literature, we use *Method 2* in [17], which is also being used for the analysis of the BAO Ly α autocorrelation for DR11 [36]. This method assumes that all quasars have the same continuum $\bar{C}(\lambda_{\text{rest}})$, except for a linear multiplicative function that varies for each quasar:

$$C_q(\lambda) = (a_q + b_q \lambda) \bar{C}(\lambda_{\text{rest}}), \quad (3.2)$$

where a_q and b_q are fitted to match an assumed probability distribution function (PDF), as explained in [17].

The construction of the continuum is a critical step for those Ly α studies focused on the line of sight power spectrum or in the flux PDF, where errors in the continuum fitting can systematically bias the results. In three dimensional clustering measurements of Ly α absorption, one would expect that the continuum fitting errors in different lines of sight are uncorrelated, getting rid of any potential bias in the measurement. However, as noted by [15], if the continuum of each quasar is rescaled in order to match an external mean transmission or flux PDF, the errors in the continuum fitting will be correlated with large scale density fluctuations. We discuss this issue further in section 3.4.

In section 4 (table 1) we present an alternative analysis using a different method to fit the continua, based on the Principal Component Analysis (PCA) presented in [40], and show that the measurement of the BAO scale is in very good agreement with our fiducial analysis.

3.2 From flux to δ_F

Using the continua described above, we now measure the mean transmitted flux fraction $\bar{F}(z)$, also known as the “mean transmission”. The redshift of absorption z is related to the observed wavelength λ by the Ly α transition line $\lambda_{\alpha} = 1216 \text{ \AA} = \lambda/(1+z)$. We measure the mean transmission as a function of redshift, in $N_z = 300$ bins between $1.9 \leq z \leq 3.4$,

$$\bar{F}(z) = \left\langle \frac{f(\lambda)}{C_q(\lambda)} \right\rangle, \quad (3.3)$$

where the average is computed over all pixels in each redshift bin. The Ly α absorption fluctuation is then defined as

$$\delta_F = \frac{f}{C_q \bar{F}} - 1. \quad (3.4)$$

As noted in [17], there are sharp features in the measurement of the mean transmission due to imperfections in the calibration vector of the BOSS data reduction pipeline. We do not expect this error to bias our results on the quasar-Ly α cross-correlation because it should

be corrected when the quasar spectra are divided by the measured mean transmission, and any residual errors are not expected to correlate with the quasar detection efficiency that varies across the BOSS survey area.

3.3 Estimator and covariance matrix

We estimate the cross-correlation ξ_A between quasars and Ly α absorption, in a bin r_A , employing the same method that was used in previous analyses of cross-correlations in BOSS ([20, 39]):

$$\hat{\xi}_A = \frac{\sum_{i \in A} w_i \delta_{Fi}}{\sum_{i \in A} w_i}, \quad (3.5)$$

where the sum is over all pixels i that are at a separation r_i in bin A from a quasar, and where the weights w_i are computed independently at each pixel from the pipeline noise variance and assuming a model for the intrinsic Ly α absorption variance (equation (3.10) in [39]).

The covariance matrix of the correlation measurements in two bins A and B , \tilde{C}_{AB} , is too large to be computed using resampling techniques or from synthetic data sets. Instead we use an analytical estimate similar to the method used in Ly α auto-correlation analyses ([15, 36]), which was first applied to cross-correlations studies in [39]:

$$\tilde{C}_{AB} \equiv \langle \hat{\xi}_A \hat{\xi}_B \rangle - \langle \hat{\xi}_A \rangle \langle \hat{\xi}_B \rangle = \frac{\sum_{i \in A} \sum_{j \in B} w_i w_j C_{ij}^F}{\sum_{i \in A} w_i \sum_{j \in B} w_j}, \quad (3.6)$$

where $C_{ij}^F = \langle \delta_{Fi} \delta_{Fj} \rangle$ is the correlation of the Lyman alpha fluctuations δ_F measured in pixels i and j , separated in redshift space by \mathbf{r}_{ij} , and includes both the cosmological signal and the contribution from instrumental noise when $i = j$. In this study, we further simplify the calculation by ignoring the correlation among Ly α pixels in different lines of sight.

This method assumes Gaussian errors and ignores the contribution from cosmic variance, a reasonable approximation given the large volume of the survey and its sparse sampling. We also assume that the variance in the quasar density field is dominated by shot noise, which is justified on the same grounds. It is useful to examine these assumptions in Fourier space (see appendix B for an extended discussion on this). The variance in the measurement of the cross-power spectrum $P_{qF}(\mathbf{k})$ for a single Fourier mode \mathbf{k} can be approximated by:

$$\text{var}(P_{qF}(\mathbf{k})) = P_{qF}(\mathbf{k})^2 + (P_{qq}(\mathbf{k}) + n_q^{-1}) (P_{FF}(\mathbf{k}) + P^{1D}(k\mu) n_{\text{eff}}^{-1}), \quad (3.7)$$

where n_q is the quasar density, P_{qq} the quasar auto-power spectrum, P_{FF} the Ly α auto-power spectrum, P^{1D} the line of sight Ly α power spectrum and n_{eff} the effective density of Ly α lines of sight as defined in [46]. Ignoring cosmic variance is equivalent to removing the term P_{qF}^2 , while assuming that quasars are shot-noise dominated is equivalent to removing the P_{qq} term. These approximations are supported by the analysis of the various terms in this equation presented in B.

We use 16 bins of constant width $10 h^{-1} \text{Mpc}$ in transverse separation r_{\perp} , up to a maximum separation of $r_{\perp} < 160 h^{-1} \text{Mpc}$. Since r_{\parallel} can be positive (pixel behind the quasar) or negative (pixel in front of the quasar) we use 32 bins in r_{\parallel} with the limits $-160 h^{-1} \text{Mpc} < r_{\parallel} < 160 h^{-1} \text{Mpc}$. We use a single bin in redshift, ranging from $2.0 < z < 3.4$. Other BAO studies that measure the correlation in multipoles, or grids defined in the $(r = \sqrt{r_{\parallel}^2 + r_{\perp}^2}, \mu = r_{\parallel}/r)$ plane, use narrower bins in order to better resolve the BAO peak. Coarser bins




Figure 2. Left panel: observed quasar-Ly α cross-correlation, $r^2\xi(r_{\parallel}, r_{\perp})$, as a function of line of sight (r_{\parallel}) and transverse (r_{\perp}) separations. Mid panel: cosmological contribution to the best fit model (see next section). Right panel: contribution of the broadband distortion to the best fit model. The fit to the observed cross-correlation is the sum of the middle and right panels. As discussed in the text, the asymmetric nature of the broadband distortion can be explained by the continuum fitting method.

can be used in studies where the correlation is measured in the $(r_{\parallel}, r_{\perp})$ plane, since each point corresponds to a different value of r . For instance, we cover 48 different values of r in the range $90 h^{-1} \text{ Mpc} < r < 120 h^{-1} \text{ Mpc}$.

As discussed in 2.3, the cross-correlations and their covariance matrices are measured in 66 sub-samples (shown in figure 1), ξ_{α} and \tilde{C}_{α} . Assuming that these are independent, the optimal way to combine them is:

$$\tilde{C}^{-1} = \sum_{\alpha} \tilde{C}_{\alpha}^{-1}, \quad \hat{\xi} = \tilde{C} \sum_{\alpha} \tilde{C}_{\alpha}^{-1} \hat{\xi}_{\alpha}. \quad (3.8)$$

When measuring the correlation in one of the sub-samples we only use Ly α pixels from spectra in that given part of the sky. However, we cross-correlate the absorption in these pixels not only with quasars from the sub-sample, but also with quasars in the neighboring sub-samples. We are therefore not losing any interesting quasar-pixel pairs, at the expense of adding a small correlation between the different measurements.

3.4 Measured cross-correlation

The quasar-Ly α cross-correlation that is obtained with the method just described is plotted in the left panel of figure 2. The model that we use to fit its functional form has two components: first, the theoretical cross-correlation function in the absence of systematics, and second, a broadband function that models systematic distortions that are introduced into the measurement. This will be described in detail in section 4, but it is useful to discuss now the general reason to fit a model with these two terms.

The middle panel in figure 2 shows the cosmological component of the best fit model, and the right panel is the broadband distortion part. The fit to the observed cross-correlation is the sum of the functions in the middle and right panels. The shape of the broadband distortion and its asymmetric nature can be explained by our method to determine the quasar

continuum, which involves fitting a multiplicative function $a_q + b_q\lambda$ to each spectrum to match an assumed PDF of the transmission F . This effectively removes large-scale power in the observed Ly α forest: roughly speaking, the mean value and the gradient of the large-scale density fluctuations over the line of sight of each Ly α spectrum are removed by the continuum fitting operation.

The distortion effect that this introduces on three-dimensional correlation measurements was first discussed in the context of the Ly α auto-correlation in appendix A of [15]. The corresponding distortion on the cross-correlation was considered in [39], where it was modeled and computed in terms of the quasar redshift distribution and the interval of the observed Ly α forest spectra. This expected distortion, plotted in figure 17 of [39], is a strong function of r_\perp and a weaker function of r_\parallel , and is asymmetric under a sign change of r_\parallel because the average quasar redshift is higher than the average Ly α pixel redshift.

An analytical prescription to correct for this distortion in the fitted theoretical model was presented in [39], valid for the simpler continuum fitting method that was used there. This was crucial for that work and in [20], where the goal was to accurately measure the full shape of the cross-correlation to obtain the bias and redshift distortion parameters.

In this paper, our goal is to measure the position of the BAO peak without any dependence on possible systematics in our modeling of the broadband shape of the cross-correlation. We therefore decide not to apply any correction to the theoretical model. Instead, a broadband term is added to the cross-correlation with enough free parameters to absorb a generic smooth distortion, as explained in section 4. This approach, also used in the recent BAO measurements from the Ly α auto-correlation ([17–19]), relies on the narrowness of the BAO peak, which decouples its position from the broadband shape. Unfortunately, this degrades our ability to measure the bias and redshift distortion parameters because they are affected by the broadband model.

4 Fitting the BAO scale

In this section we describe the method used to measure the scale of the Baryon Acoustic Oscillations (BAO) from the measured cross-correlations, and present our main results. We conclude with a detailed analysis of possible sources of systematic errors.

4.1 BAO model

We adapted the publicly available fitting code `baofit` [19] to work with cross-correlations. The code can be downloaded from the URL in footnote,² together with the measured cross-correlation and its covariance matrix, as described in appendix A.

A detailed description of the fitting code can be found in [19]; here we only summarize the main points and highlight the differences between fitting the Ly α auto-correlation and the quasar-Ly α cross-correlation.

We model the measured cross-correlation $\xi(r_\parallel, r_\perp)$ as a sum of the *cosmological* correlation ξ_{cosmo} and a broadband distortion term ξ_{bb} due primarily to continuum fitting (as discussed in section 3.4)

$$\xi(r_\parallel, r_\perp) = \xi_{\text{cosmo}}(r_\parallel, r_\perp, \alpha_\parallel, \alpha_\perp) + \xi_{\text{bb}}(r_\parallel, r_\perp) . \quad (4.1)$$

²<http://darkmatter.ps.uci.edu/baofit/>.

The quantity ξ_{cosmo} is described as a sum of two terms

$$\xi_{\text{cosmo}}(r_{\parallel}, r_{\perp}) = \xi_{\text{smooth}}(r_{\parallel}, r_{\perp}) + a_{\text{peak}} \cdot \xi_{\text{peak}}(r_{\parallel} \alpha_{\parallel}, r_{\perp} \alpha_{\perp}), \quad (4.2)$$

where a_{peak} controls the amplitude of the BAO peak. The correct ξ_{cosmo} in the Cold Dark Matter standard model is obtained only for $a_{\text{peak}} = 1$, which we use in all of our analyses except when we want to test the consistency of our results with the prediction for the peak amplitude (row labeled AMP in table 1).

The main goal of this study is to measure the scale of BAO relative to the fiducial cosmological model, along the line of sight α_{\parallel} and across the line of sight α_{\perp} :

$$\alpha_{\parallel} = \frac{[D_H(\bar{z})/r_s]}{[D_H(\bar{z})/r_s]_{\text{fid}}}, \quad \alpha_{\perp} = \frac{[D_A(\bar{z})/r_s]}{[D_A(\bar{z})/r_s]_{\text{fid}}}, \quad (4.3)$$

where r_s is the sound horizon, $D_H(z) = c/H(z)$ is the Hubble distance, and $D_A(z)$ the comoving angular diameter distance. The mean redshift of our measurement is $\bar{z} = 2.36$.

The scale factors $(\alpha_{\parallel}, \alpha_{\perp})$ only appear in the *peak* part of the correlation, to ensure that no information comes from the broadband shape. A detailed description of the decomposition of the cosmological signal into a *peak* and a *smooth* component can be found in [19].

4.1.1 Theoretical model for the cross-correlation

We model the *cosmological* correlation $\xi_{\text{cosmo}}(\mathbf{r})$ as the 3D Fourier transform of the cross-power spectrum $P_{qF}(\mathbf{k})$:

$$P_{qF}(\mathbf{k}) = b_q [1 + \beta_q \mu_k^2] b_F [1 + \beta_F \mu_k^2] P(k), \quad (4.4)$$

where b_q is the linear bias of quasars, b_F the linear bias of Ly α forest, and β_q and β_F the redshift space distortion parameters for quasars and Ly α forest. The matter power spectrum is $P(k)$, which includes the non-linear broadening of the peak [19] and μ_k is the cosine of the angle between the Fourier mode vector \mathbf{k} and the line of sight.

Following [20] we leave two of the four bias parameters free (b_q and β_F) and derive the other two from them, using the well-constrained combination $b_F(1 + \beta_F) = -0.336 \pm 0.03$ [15] and the Kaiser relation $b_q \beta_q = f(\Omega_m)$ [41], where $f(\Omega_m)$ is the logarithmic growth rate of structure. Note that the same relation does not apply to the Ly α forest (e.g., [15]). These values of the bias parameters are defined at $z = 2.25$, and we translate them to our mean redshift $\bar{z} = 2.36$ assuming that only b_F evolves with redshift, following $(b_F(z)g(z))^2 \propto (1+z)^{3.8}$, where $g(z)$ is the linear growth factor (as discussed in [15]).

4.1.2 Quasar redshift errors

Determining precise quasar redshifts is a difficult task. As noted in [20], quasar redshift errors have two main effects on the cross-correlation: a) the r.m.s. in the quasar redshift estimates ($\sigma_z \sim 500 \text{ km s}^{-1}$) smooths the cross-correlation along the line of sight (with an equivalent effect on the quasar auto-correlation, [21]); b) a systematic offset in the BOSS redshift estimates shifts the cross-correlation along the line of sight by a non-negligible amount $\Delta_z \sim -180 \text{ km s}^{-1}$ (see [20]).

Since we restrict our analysis to large separations ($r > 40 h^{-1} \text{ Mpc}$), we do not expect quasar redshift errors to have a significant impact on our fits. We leave Δ_z as a free parameter in all our fits, presenting our results after marginalizing over it. We do not include an explicit σ_z parameter since it would be highly degenerate with the non-linear broadening model.

4.1.3 Broadband distortion

All BAO analyses to date have used a broadband model that parameterizes each multipole as a function of r , or a parameterization as a function of (r, μ) as in [18]. However, the shape of the distortion discussed in section 3.4 that is introduced into the cross-correlation by the continuum fitting operation is better separated in terms of the $(r_{\parallel}, r_{\perp})$ coordinates, as inferred from the analysis that was presented in [39] (see their figure 17). In this figure, one can see that the distortion decreases rapidly with r_{\perp} , and it has a non-trivial r_{\parallel} dependence. Therefore, we use the following parameterization for the broadband distortion model:

$$\xi_{bb}(r_{\parallel}, r_{\perp}) = \sum_{i=i_{\min}}^{i_{\max}} \sum_{j=j_{\min}}^{j_{\max}} b_{ij} r_{\parallel}^i r_{\perp}^j, \quad (4.5)$$

where the sums are understood to be over consecutive integers, and they go from $i_{\min} = 0$ to $i_{\max} = 2$, and from $j_{\min} = -3$ to $j_{\max} = 1$ in our fiducial analysis. The dependence of our results on the broadband distortion model is discussed in section 4.3.

4.2 BAO fits

Our fiducial BAO fit is performed over the separation range $40 h^{-1} \text{ Mpc} < r < 180 h^{-1} \text{ Mpc}$ using a broadband model with $(i_{\min} = 0, i_{\max} = 2, j_{\min} = -3, j_{\max} = 1)$. The total number of bins included is 440, and the number of free parameters is 20: $\alpha_{\parallel}, \alpha_{\perp}, \beta_F, b_q, \Delta_z$ and the 15 parameters b_{ij} in our broadband distortion model. In table 1 we present the best fit values for our fiducial analysis, and for a series of illustrative alternative analyses: an isotropic BAO analysis (ISO) imposing $\alpha \equiv \alpha_{\parallel} = \alpha_{\perp}$; a no-wiggles fit (NW) with $a_{\text{peak}} = 0$; a fit allowing the amplitude of the peak a_{peak} to vary (AMP); and a fit using a different method to fit the continua based on a Principal Component Analysis (PCA, [40]).

The BAO peak position is significantly measured to an accuracy better than $\sim 4\%$ both along and across the line of sight directions. The measured amplitude of the BAO peak is consistent with the expected in our fiducial model.

In figure 3 we present the main result of this paper: the value of $\Delta\chi^2$ as a function of $(\alpha_{\parallel}, \alpha_{\perp})$ for our fiducial BAO analysis, fully marginalized over the other 18 free parameters. The solid contours correspond to $\Delta\chi^2 = 2.27, 5.99$ and 11.62 , equivalent to likelihood contours of 68%, 95% and 99.7% for a Gaussian likelihood. The fiducial model is consistent at the $\sim 1.5\sigma$ level.

We can translate our measurement of $(\alpha_{\parallel}, \alpha_{\perp})$ to a measurement of the Hubble parameter and the angular diameter distance at our mean redshift $z = 2.36$, up to a factor r_s :

$$c/(H(z = 2.36) r_s) = 9.0 \pm 0.3, \quad D_A(z = 2.36) / r_s = 10.8 \pm 0.4. \quad (4.6)$$

Using the best fit value of the sound horizon from the Planck collaboration ($r_s = 147.49 \text{ Mpc}$) [3],³ we can present the results as:

$$H(z = 2.36) = 226 \pm 8 \text{ km s}^{-1} \text{ Mpc}^{-1} \quad D_A(z = 2.36) = 1590 \pm 60 \text{ Mpc}. \quad (4.7)$$

³Table 2, column with 68% limits for Planck+WP.




Figure 3. $\Delta\chi^2$ as a function of $\alpha_{\parallel}, \alpha_{\perp}$ (defined in equation (4.3)) in our fiducial analysis, after marginalizing over the remaining 18 parameters. The solid contours correspond to $\Delta\chi^2 = 2.27, 5.99$ and 11.62 , equivalent to Gaussian probabilities of 68%, 95% and 99.7%. The fiducial model is consistent at the $\sim 1.5\sigma$ level.

4.3 Systematic tests

Table 2 presents the dependence of our results on the broadband model for our fiducial analysis. We start by presenting the results in the absence of any broadband distortion term (NO BB row), and we increasingly add more free parameters to our model in an attempt to remove the distortion caused by the continuum fitting method (see figure 17 of [39]). Adding a single constant (BB_0) does not improve the fit, but adding a r_{\perp}^{-1} term reduces the best fit χ^2 by 30. The goodness of fit keeps improving while adding new free parameters, until it saturates close to our fiducial model (BB_7), after which adding new parameters does not improve the fit much. The BAO results are very insensitive to the chosen broadband function form for all models with more than 6 free parameters (BB_4 - BB_14), and even the model with only 2 free parameters gives very similar results.

Table 3 presents the dependence on the separation range over which the cross-correlation is fitted, when the maximum separation is modified from the fiducial value of $180 h^{-1}$ Mpc to $170 h^{-1}$ Mpc (RMAX_170) or to $190 h^{-1}$ Mpc (RMAX_190), and the minimum separation from $40 h^{-1}$ Mpc to $30 h^{-1}$ Mpc (RMIN_30) or to $50 h^{-1}$ Mpc (RMIN_50). The last three rows show the results of restricting the range of the angle cosine $\mu = r_{\parallel}/r$ to $\|\mu\| < 0.8$ (MU_08), $\|\mu\| < 0.9$ (MU_09), or $\|\mu\| < 0.95$ (MU_095). The results in this table show that the BAO measurement in general has little dependence on the fitting range. The broadband distortion is most important for separations near the line of sight (i.e., $\|\mu\|$ near one), but the removal of this most contaminated part does not significantly alter the BAO peak position that is obtained, except in the MU_095 case where the position that is obtained shifts to a value closer to the expected one in our fiducial model by nearly 1σ .

4.4 Test of the covariance matrix

In table 1 we can see that the χ^2 value in our fiducial fit is good, in the sense that it is compatible with being drawn from a χ^2 distribution with mean equal to the degrees of

Model	N_{bb}	β_F	b_q	α_{\parallel}	α_{\perp}	χ^2 Prob
NO BB (-)	0	2.85 ± 0.62	2.75 ± 0.16	1.049 ± 0.039	0.902 ± 0.048	0.056
BB_0 (0, 0, 0, 0)	1	2.87 ± 0.62	2.75 ± 0.16	1.050 ± 0.039	0.902 ± 0.048	0.054
BB_1 (0, 0, -1, 0)	2	1.55 ± 0.30	2.73 ± 0.16	1.038 ± 0.039	0.921 ± 0.041	0.255
BB_2 (0, 1, -1, 0)	4	1.51 ± 0.29	2.75 ± 0.16	1.039 ± 0.038	0.921 ± 0.040	0.254
BB_3 (0, 2, -1, 0)	6	1.22 ± 0.25	2.94 ± 0.18	1.047 ± 0.036	0.916 ± 0.040	0.318
BB_4 (0, 2, -2, 0)	9	0.91 ± 0.22	3.11 ± 0.20	1.044 ± 0.035	0.927 ± 0.037	0.350
BB_5 (0, 2, -2, 1)	12	0.95 ± 0.23	3.14 ± 0.21	1.043 ± 0.034	0.933 ± 0.036	0.397
BB_6 (0, 2, -3, 0)	12	0.94 ± 0.24	3.05 ± 0.22	1.047 ± 0.035	0.924 ± 0.037	0.381
BB_7 (0, 2, -3, 1)	15	1.09 ± 0.29	3.02 ± 0.22	1.042 ± 0.034	0.930 ± 0.036	0.404
BB_8 (0, 3, -2, 1)	16	0.88 ± 0.22	3.14 ± 0.21	1.043 ± 0.033	0.932 ± 0.036	0.467
BB_9 (0, 3, -3, 1)	20	1.01 ± 0.27	3.01 ± 0.22	1.042 ± 0.034	0.929 ± 0.036	0.463
BB_10 (-1, 3, -2, 1)	20	0.89 ± 0.22	3.14 ± 0.21	1.045 ± 0.033	0.931 ± 0.036	0.443
BB_11 (0, 4, -2, 1)	20	0.74 ± 0.18	3.23 ± 0.26	1.052 ± 0.034	0.930 ± 0.035	0.525
BB_12 (0, 5, -2, 1)	24	0.80 ± 0.23	3.17 ± 0.27	1.051 ± 0.034	0.932 ± 0.035	0.517
BB_13 (0, 4, -2, 2)	25	0.75 ± 0.21	3.15 ± 0.28	1.052 ± 0.034	0.923 ± 0.035	0.514
BB_14 (0, 4, -3, 1)	25	0.90 ± 0.27	3.03 ± 0.28	1.047 ± 0.035	0.929 ± 0.036	0.513

Table 2. Best fit parameters for different broadband distortion models, defined by their values of $(i_{\min}, i_{\max}, j_{\min}, j_{\max})$ in equation (4.5). N_{bb} is the total of free parameters of the broadband distortion model. The last column shows the probability inferred from the χ^2 value in the fit and the number of degrees of freedom (different in each row). The first row (NO BB) shows the results for a fit without distortion, and row BB_7 corresponds to our fiducial model.

	β_F	b_q	α_{\parallel}	α_{\perp}	χ^2 (d.o.f)
FID	1.09 ± 0.29	3.02 ± 0.22	1.042 ± 0.034	0.930 ± 0.036	426.4 (420)
RMAX_170	1.07 ± 0.28	3.07 ± 0.22	1.046 ± 0.034	0.926 ± 0.036	397.4 (394)
RMAX_190	1.05 ± 0.27	3.04 ± 0.22	1.042 ± 0.034	0.929 ± 0.036	438.3 (436)
RMIN_30	1.03 ± 0.21	2.85 ± 0.14	1.043 ± 0.035	0.930 ± 0.038	443.5 (430)
RMIN_50	1.65 ± 0.71	3.01 ± 0.34	1.044 ± 0.034	0.923 ± 0.038	407 (406)
MU_08	2.26 ± 2.0	2.61 ± 0.38	1.025 ± 0.088	0.937 ± 0.055	257.1 (244)
MU_09	1.39 ± 0.89	2.73 ± 0.41	1.023 ± 0.054	0.938 ± 0.041	306.7 (302)
MU_095	0.88 ± 0.32	3.13 ± 0.27	1.009 ± 0.041	0.949 ± 0.037	353.6 (344)

Table 3. Best fit parameters for the fiducial analysis (FID), and different fitting ranges (in h^{-1} Mpc). The last rows show the results when using only bins that are far from the line of sight, with $|\mu| < 0.8$ (MU_08), $|\mu| < 0.9$ (MU_09) and $|\mu| < 0.95$ (MU_095).

freedom in the problem, i.e., the number of bins used in the fit (440) minus the number of free parameters (20).

In order to test our estimate of the covariance matrix, we examine the distribution of χ^2 for its different eigenmodes. The results of this test are compared to a zero-mean Gaussian with variance $(440 - 20)/440$ in figure 4. The agreement supports the validity of our covariance matrix.




Figure 4. Histogram of χ^2 values for the different eigenmodes of the covariance matrix, compared to a zero-mean Gaussian with variance $(440 - 20)/440$, where 440 is the number of bins in the fit, and 20 is the number of free parameters. The agreement between the distributions supports the validity of our covariance matrix.

4.5 Alternative uncertainty estimates of the BAO scales

The error on the fitted parameters reported so far have been computed from the second derivatives of the log-likelihood function at its maximum, assuming this likelihood function to be Gaussian at 1σ . The BAO scale uncertainties in the fiducial analysis obtained in this way are 0.034 for α_{\parallel} and 0.036 for α_{\perp} . An alternative error estimate can be computed from the full likelihood surface in figure 3, without assuming a Gaussian likelihood. The uncertainties obtained in the fiducial analysis are then 0.032 for α_{\parallel} and 0.036 for α_{\perp} , in good agreement with the previous ones.

Both these estimates rely on the accuracy of the covariance matrix that we have computed as described in section 3. We test this by computing an alternative bootstrap error on the BAO scale parameters, that does not rely on our covariance matrix. We generate 1,000 bootstrap realizations of the survey [42], combining the measurements from the 66 different sub-samples. The fitting analysis is done for each realization, and the uncertainties on α_{\parallel} and α_{\perp} are computed from their distribution of best fit values. The resulting uncertainties on the BAO scales are 0.031 on α_{\parallel} and 0.036 on α_{\perp} , in excellent agreement with the previous estimates.

4.6 Visualizing the BAO peak

Even though we do not use multipoles anywhere in our analysis, we present here a fit of the multipoles from the measured cross-correlation in order to better *see* the BAO peak in the data. We start by constructing a multipole expansion of our measured cross-correlation, $\xi(r_{\parallel}, r_{\perp})$, using a linear least-squares fit to:

$$\xi(r, \mu) = \sum_l L_l(\mu) \xi_l(r), \quad (4.8)$$

where $r = \sqrt{r_{\parallel}^2 + r_{\perp}^2}$ and $\mu = r_{\parallel}/r$, $L_l(x)$ is the Legendre polynomial of order l and $\xi_l(r)$ are the multipoles we wish to measure.

In figure 2 we show the measured cross-correlation, as a function of line of sight (r_{\parallel}) and transverse (r_{\perp}) separation, together with our best fit model. From the right panel of the figure, one can see that the best fit model of the broadband distortion is asymmetric with




Figure 5. Transverse (left) and parallel (right) correlations, defined as $\xi(r, \mu = 0) = \xi_0(r) - \xi_2(r)/2$ and $\xi(r, \mu = 1) = \xi_0(r) + \xi_2(r)$, after projecting out the mode responsible for most of the correlation between separations. The best fit theory is shown in a solid black curve, its BAO-only part in a red dashed curve and the distortion in a dotted green curve. The orange dot-dashed curve shows the cosmological signal for our fiducial cosmology ($\alpha = 1$), using a quasar bias of $b_q = 3.64$ and a Ly α redshift-space distortion parameter $\beta_F = 1.1$, as measured in [20]. All datapoints and lines are weighted by r^2 and are plotted after the projection (see text for details).

respect to $r_{\parallel} = 0$. Therefore we expect a net non-zero contribution from odd multipoles. For the purpose of visualization, however, we only fit the monopole ($l = 0$) and the quadrupole ($l = 2$), since these two multipoles contain most of the cosmological information. We use 36 equidistant interpolation points separated by $4 h^{-1}$ Mpc and ranging from $40 h^{-1}$ Mpc to $180 h^{-1}$ Mpc.

Our estimates of the multipoles at different separations are highly correlated. In order to improve the visualization of the BAO peak, we apply a correction to the multipoles based on the analysis presented in [19]. We start by examining the eigenmodes of the covariance matrix and identify a particular mode being essentially a DC offset of the monopole, and therefore responsible for much of the correlations between separations. We then project out the mode from the data and its covariance matrix, and refit for the distortion while keeping all other parameters fixed from the baseline best fit.

Figure 5 shows the resulting monopole and quadrupole of the quasar-Ly α cross-correlation, expressed as the transverse correlation, $\xi(r, \mu = 0) = \xi_0(r) - \xi_2(r)/2$ (left panel), and the parallel correlation $\xi(r, \mu = 1) = \xi_0(r) + \xi_2(r)$. We superimpose a fit with all parameters fixed from the 2D BAO fit except for the distortion. The solid black curve shows the best fit, the red dashed curve is the BAO-only part (with parameters fixed from the 2D fit), and the green dotted curve shows the distortion, which is parabolic after r^2 weighting. Since the Ly α fluctuation is defined in equation (3.4) as a transmission fluctuation, positive values of δ_F reflect negative density fluctuations, implying a negative value for the bias factor b_F . This explains why the BAO feature in the quasar-Ly α cross-correlation appears as a dip instead of a peak, as seen in figure 5.

The orange curve in figure 5 shows the predicted cross-correlation for our fiducial cosmological model with $\alpha_{\parallel} = \alpha_{\perp} = 1$, and an amplitude determined by a quasar bias factor

Analysis	Probe	Data Release	α	α_{\parallel}	α_{\perp}
Busca_2013	Auto	DR9	1.01 ± 0.03	-	-
Slosar_2013	Auto	DR9	0.98 ± 0.020	0.99 ± 0.035	0.98 ± 0.070
This work	Cross	DR10	1.00 ± 0.027	1.06 ± 0.038	0.91 ± 0.041
This work	Cross	DR11	0.99 ± 0.022	1.04 ± 0.034	0.93 ± 0.036
Busca_2013	Auto	to DR11	± 0.019	-	-
Slosar_2013	Auto	to DR11	± 0.013	± 0.022	± 0.046

Table 4. Comparison of different BAO analysis at $z \sim 2.4$ from BOSS, from the auto-correlation in DR9 (Busca_2013 [17], Slosar_2013 [18]), and from this work. We show our results when using only DR10 data and when including DR11 data. In the last two rows we extrapolate the uncertainties of previous work to DR11, assuming that these scale with the inverse of the square root of the survey area.

$b_q = 3.64$ and a Ly α redshift distortion parameter $\beta_F = 1.1$, as measured in [20]. The fact that this model is consistent with the best fit that is obtained here to the DR11 data proves that our result is consistent with that obtained in [20] using the DR9 data, and that the different values that are obtained in our fit for b_q and β_F are caused by our addition of an arbitrary broadband function, with parameters that are degenerate with b_q and β_F . The amplitude of the BAO dip, as visualized in figure 5, is consistent with our expectation. This is seen also in the model AMP in table 1, where the parameter a_{peak} has a best fit value that is consistent with unity. A model with a suppressed BAO peak (model NW in table 1) has a χ^2 that is worse than our fiducial model by 20, although we warn that this is not to be directly interpreted as a statistical significance of a BAO detection because our likelihood function is not necessarily Gaussian. In any case, our interest here lies in the statistical constraint obtained on the BAO scale, rather than the significance of the BAO detection in the quasar-Ly α cross-correlation only.

5 Discussion & conclusions

We have presented a measurement of the quasar-Ly α cross-correlation using approximately 164,000 quasars from the eleventh Data Release (DR11) of SDSS. We are able to measure the BAO scale along and across the line of sight (α_{\parallel} , α_{\perp}) with an uncertainty of 3.4% and 3.6% respectively. The measurement is in agreement with our fiducial cosmology well within the 95% confidence level.

We have checked the robustness of our measurement under changes of broadband models, separation range used, and different error estimates. As discussed in section 3.4, we are not particularly careful in our treatment of the non-BAO part of the cross-correlation. However, the best fit values for the bias parameters of both quasars and Ly α forest are roughly consistent with previous analyses, with rather large uncertainties since we only use large separations to measure the BAO scale.

In table 4 we compare the results with other BAO measurements at the same redshift from the Ly α auto-correlation measured with DR9 ([17, 18]). We also present our results when using only data from DR10. Assuming that the uncertainties in these BAO measurements scale with the inverse of the square root of the survey area, we can extrapolate them from DR9 to DR11, using $\sqrt{A_{\text{DR11}}/A_{\text{DR9}}} = 1.60$. We show these extrapolations in the last two rows of table 4.

The errors on the BAO scale ($\alpha_{\parallel}, \alpha_{\perp}$) from our DR10 analysis are considerably smaller than those reported in [43]. An extensive comparison of the two analyses within the BOSS Ly α working group concluded that the discrepancy can be explained by the differences in the analysis. While [43] uses only the monopole and the quadrupole to fit the BAO scale, in this analysis we use the full 2D contours of the cross-correlation function.

In the absence of any broadband distortion of $\xi(r_{\parallel}, r_{\perp})$ (or with a distortion that is a priori known), we find that essentially all of the BAO signal is contained within the monopole $\xi_0(r)$ and quadrupole $\xi_2(r)$. However, when broadband distortion is present, as in our analysis, it contributes significantly to multipoles other than the monopole and quadrupole, and leads to correlated uncertainties between distortion and BAO parameters and corresponding parameter degeneracies. As a result, we find that the unknown broadband distortion parameters can be determined more precisely with a fit to the full $\xi(r_{\parallel}, r_{\perp})$ (or, equivalently, a larger set of multipoles) instead of a fit to only the monopole and quadrupole. Similarly, we find that a fit to $\xi(r_{\parallel}, r_{\perp})$ yields a more precise determination of the BAO parameters by helping to break the degeneracy between distortion and BAO parameters. The actual improvement we find is a factor of 1.2 in α_{\parallel} and a factor of 1.3 in α_{\perp} . [36] found that a similar improvement is also seen when measuring BAO from the Ly α auto-correlation function, although a detailed study on mock data sets revealed a large scatter in the gain from realization to realization.

5.1 Ly α auto-correlation vs. quasar-Ly α cross-correlation

In appendix B we present a Fisher matrix projection comparing the relative strength of measuring BAO with the Ly α auto-correlation and with the quasar-Ly α cross-correlation. In a BOSS-like survey, both probes should measure the transverse BAO scale with similar uncertainties, while the Ly α auto-correlation should be able to measure the line of sight scale $\sim 40\%$ better than the cross-correlation with quasars.

The measurement of BAO from the Ly α auto-correlation in DR9 was presented in [17] and [18]. Most of the difference between the uncertainties in these results can be explained by the looser data cuts used in [18], that included lines with DLAs and that defined their Ly α forest with a wider wavelength range. In this analysis we used data cuts similar to those in [18], and therefore we compare here our uncertainties with those from [18] extrapolated to DR11 (see table 4).

Our measurement of α_{\parallel} is $\sim 55\%$ worse than the results from the Ly α auto-correlation of [18] extrapolated to DR11, in good agreement with the prediction of $\sim 40\%$ computed in the appendix. The Fisher forecast formalism predicted similar uncertainties in α_{\perp} , and we find that our measurement is $\sim 20\%$ better than the extrapolated results from the auto-correlation.

In the same appendix we also show that on the scales of interest for BAO measurements ($k > 0.05 h \text{ Mpc}^{-1}$) cosmic variance is not the dominant contribution to our error budget. Assuming that the shot noise in the quasar density field is uncorrelated with the small scale fluctuations in the Ly α absorption and with the instrumental noise, we can then combine both BAO measurements as if they were independent.

In figure 6 we compare the contours on $(\alpha_{\parallel}, \alpha_{\perp})$ from the Ly α auto-correlation function from DR9 ([18] in blue, generated from the files in <http://darkmatter.ps.uci.edu/baofit/>), and compare it to our measurement from the cross-correlation function from DR11 (red) and the sum of their χ^2 surfaces (in black), assuming they are independent. We compare these constraints with the 68% and 95% confidence limits obtained from the Planck results [3] in




Figure 6. Contours of $\Delta\chi^2 = 2.27$ and 5.99 , corresponding to Gaussian confidence levels of 68% and 95%, from the $\text{Ly}\alpha$ auto-correlation analysis from DR9 ([18], in blue), from the cross-correlation from DR11 (this work, in red) and from the joint analysis (in black). The green contours show the 68% and 95% contours for the regions of this parameter space allowed by the Planck results [3] in an open ΛCDM cosmology.

an open ΛCDM cosmology, shown in green.⁴ Note that by allowing for space curvature, the Planck constraints on the distance and expansion rate at our mean redshift $z = 2.36$ are much less restrictive compared to a flat model.

We have shown that adding the cross-correlation of $\text{Ly}\alpha$ and quasars to the auto-correlation of $\text{Ly}\alpha$ can certainly improve the constraints on BAO scales at high redshift. A detailed analysis of the cosmological implications of the measurements of the $\text{Ly}\alpha$ auto-correlation and the quasar- $\text{Ly}\alpha$ cross-correlation will be presented in a future publication, which will include the DR11 results from the $\text{Ly}\alpha$ auto-correlation, together with a more complete examination of potential correlations between the two measurements.

Acknowledgments

We would like to thank Ross O’Connell for detailed comparisons with his analysis, and Shirley Ho for very useful comments.

This research used resources of the National Energy Research Scientific Computing Center (NERSC), which is supported by the Office of Science of the U.S. Department of Energy under Contract No. DE-AC02-05CH11231. DK would like to thank CEA Saclay for their hospitality and productive environment during his sabbatical. JM is supported in part by Spanish grant AYA2012-33938.

Funding for SDSS-III has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, and the U.S. Department of Energy Office of Science. The SDSS-III web site is <http://www.sdss3.org/>.

SDSS-III is managed by the Astrophysical Research Consortium for the Participating Institutions of the SDSS-III Collaboration including the University of Arizona, the Brazilian

⁴We use the Planck + ACT/SPT + WP public chains available under the name `base_omegak_planck_lowL_lowLike_highL`.

Participation Group, Brookhaven National Laboratory, University of Cambridge, Carnegie Mellon University, University of Florida, the French Participation Group, the German Participation Group, Harvard University, the Instituto de Astrofísica de Canarias, the Michigan State/Notre Dame/JINA Participation Group, Johns Hopkins University, Lawrence Berkeley National Laboratory, Max Planck Institute for Astrophysics, Max Planck Institute for Extraterrestrial Physics, New Mexico State University, New York University, Ohio State University, Pennsylvania State University, University of Portsmouth, Princeton University, the Spanish Participation Group, University of Tokyo, University of Utah, Vanderbilt University, University of Virginia, University of Washington, and Yale University.

A Public access to data and code

The `baofit` software used in this paper is publicly available at <https://github.com/deepzot/baofit/>. The measured cross-correlation function and its covariance matrix, and the instructions to reproduce the BAO constraints presented in this paper, can be downloaded from <http://darkmatter.ps.uci.edu/baofit/>, together with the likelihood surface used to generate figure 3. The software is written in C++ and uses MINUIT [44] for likelihood minimization.

B Fisher matrix forecasts

The goal of this appendix is to compare the expected accuracy with which one should be able to measure the BAO scale from a BOSS-like survey using the $\text{Ly}\alpha$ auto-correlation function and the quasar- $\text{Ly}\alpha$ cross-correlation.

Instead of undertaking a full forecast of the uncertainty on the BAO scale, we will do a simpler comparison and evaluate the signal to noise ratio (S/N) that one should obtain for a certain Fourier mode \mathbf{k} . This is a fair comparison, since the uncertainty on the BAO scale should be proportional to the uncertainty on the power spectrum over the relevant BAO scales.

B.1 Auto-correlation

We start by computing the expected signal to noise ratio in the auto-correlation of $\text{Ly}\alpha$ and in the auto-correlation of quasars. On large scales, the signal in the auto-correlation can be described with a simple linear bias model with the Kaiser model to account for redshift space distortions:

$$P_A(\mathbf{k}) = b_A^2 (1 + \beta_A \mu^2)^2 P(k), \quad (\text{B.1})$$

where b_A and β_A are the linear bias parameter of the tracer A and its redshift space distortion parameter, $P(k)$ is the matter power spectrum, and μ is the cosine of the angle between the Fourier mode \mathbf{k} and the line of sight.

The accuracy with which one can measure the quasar power spectrum $P_A(\mathbf{k})$ in a given bin centered at (k, μ) can be quantified by the signal to noise ratio (S/N),

$$\left(\frac{S}{N}\right)_A^2 = N_k \frac{P_A^2(\mathbf{k})}{\text{var}[P_A(\mathbf{k})]}, \quad (\text{B.2})$$

where N_k is the number of modes in the bin. Since we only care about relative performance in this appendix, we will drop any N_k and will plot signal to noise ratio per mode.

For a sample of point-like sources (for instance quasars), the variance of its measured power spectrum can be approximated by

$$\text{var} [P_A(\mathbf{k})] = 2 \left(P_A(\mathbf{k}) + n_A^{-1} \right)^2, \quad (\text{B.3})$$

with n_A the number density of systems.

Since the Ly α forest is not a discrete point sampling of the underlying matter density field, but rather a non-linear transformation of a continuous sampling along discrete lines of sight, we need to use a slightly different approach. [45] computed the expected S/N in the measurement of $P_{FF}(\mathbf{k})$ in a spectroscopic survey, and highlighted the importance of the “aliasing term” due to the sparse sampling of the universe. Here we use the formalism from [46] that combines both the noise term and the aliasing term defining a noise-weighted density of lines of sight per unit area n_{eff} ,

$$\text{var}[P_{FF}(\mathbf{k})] = 2 \left(P_{FF}(\mathbf{k}) + P^{1D}(k\mu) n_{\text{eff}}^{-1} \right)^2, \quad (\text{B.4})$$

where $P^{1D}(k\mu)$ is the one-dimensional flux power spectrum.

B.2 Cross-correlation

The cross correlation between the Ly α absorption and the quasar density field can be defined as

$$\langle \delta_F(\mathbf{k}) \delta_q(\mathbf{k}') \rangle = (2\pi)^3 \delta^D(\mathbf{k} + \mathbf{k}') P_{qF}(\mathbf{k}). \quad (\text{B.5})$$

Again, in the linear regime we can relate the cross-correlation power spectrum with the linear power spectrum $P(k)$ using the linear bias parameters defined above,

$$P_{qF}(\mathbf{k}) = b_q (1 + \beta_q \mu^2) b_F (1 + \beta_F \mu^2) P(k). \quad (\text{B.6})$$

[46] showed that the variance in the measurement of the cross-correlation can be approximated by

$$\text{var} (P_{qF}(\mathbf{k})) = P_{qF}(\mathbf{k})^2 + (P_{qq}(\mathbf{k}) + n_q^{-1}) (P_{FF}(\mathbf{k}) + P^{1D}(k\mu) n_{\text{eff}}^{-1}). \quad (\text{B.7})$$

In this approximation, the expected S/N in a bin of (k, μ_k) can be approximated by

$$\left(\frac{S}{N} \right)_{Fg}^2 = N_k \frac{P_{qF}^2(\mathbf{k})}{P_{qF}(\mathbf{k})^2 + (P_{qq}(\mathbf{k}) + n_q^{-1}) (P_{FF}(\mathbf{k}) + P^{1D}(k\mu) n_{\text{eff}}^{-1})}. \quad (\text{B.8})$$

B.3 Forecast for a BOSS-like survey

Here we quantify the previous results for the case of a spectroscopic survey with properties similar to the BOSS survey. The BOSS survey has an area of $A = 10^4 \text{ deg}^2$, and if we restrict the analysis to the redshift range $2 < z < 3$, the total volume of the survey is roughly $V = 40(h^{-1} \text{ Gpc})^3$. The quasar density in the BOSS survey is roughly $n_q = 160000/V \sim 4 \times 10^{-6} (h^{-1} \text{ Mpc})^{-3}$, and we assume a quasar bias of $b_q = 3.6$ ([20, 21]). The effective density of lines of sight for BOSS is estimated in [46] to be $n_{\text{eff}} \approx 10^{-3} (h^{-1} \text{ Mpc})^{-2}$, and we assume the values for the Ly α biases of $b_F = -0.15$ and $\beta_F = 1.2$, both compatible with the 1D measurement of [47] and the 3D clustering from [15]. We compute the power spectra at our fiducial redshift of $z_c = 2.36$.




Figure 7. Study of the signal to noise ratio in different analyses: auto-correlation of Ly α (top left), auto-correlation of quasars (top right) and their cross-correlation (bottom left). The red solid lines show the signal, the green dashed lines the “shot noise” level and the dotted blue lines their sum, for different values of μ (increasing from lower to upper lines). The bottom-right panel shows the expected $(S/N)^2$ per mode for the Ly α auto-correlation (dashed green), quasar auto-correlation (solid red) and cross-correlation (dotted blue).

In figure 7 we compare the signal and the different noise contributions for the different analyses: Ly α auto-correlation (top left), quasar auto-correlation (top right) and quasar-Ly α cross-correlation (bottom left). In the bottom-right panel we compare the expected signal to noise ratio (squared) per mode for the three different analyses, and for different values of μ_k . We can see that the S/N of the quasar-Ly α cross-correlation is much higher than the quasar auto-correlation, and that for transverse modes (lower lines) is as high as the Ly α auto-correlation. It is also clear from the figure that on scales relevant for BAO ($k > 0.05 h \text{ Mpc}^{-1}$), we are in the noise-dominated regime, and therefore cosmic variance is at best a secondary contribution to the error budget.

References

- [1] SUPERNOVA SEARCH TEAM collaboration, A.G. Riess et al., *Observational evidence from supernovae for an accelerating universe and a cosmological constant*, *Astron. J.* **116** (1998) 1009 [[astro-ph/9805201](#)] [[INSPIRE](#)].
- [2] SUPERNOVA COSMOLOGY PROJECT collaboration, S. Perlmutter et al., *Measurements of Ω and Λ from 42 high redshift supernovae*, *Astrophys. J.* **517** (1999) 565 [[astro-ph/9812133](#)] [[INSPIRE](#)].

- [3] PLANCK collaboration, P.A.R. Ade et al., *Planck 2013 results. XVI. Cosmological parameters*, [arXiv:1303.5076](#) [INSPIRE].
- [4] D.H. Weinberg, M.J. Mortonson, D.J. Eisenstein, C. Hirata, A.G. Riess and E. Rozo, *Observational Probes of Cosmic Acceleration*, *Phys. Rept.* **530** (2013) 87 [[arXiv:1201.2434](#)] [INSPIRE].
- [5] H.-J. Seo and D.J. Eisenstein, *Probing dark energy with baryonic acoustic oscillations from future large galaxy redshift surveys*, *Astrophys. J.* **598** (2003) 720 [[astro-ph/0307460](#)] [INSPIRE].
- [6] SDSS collaboration, D.J. Eisenstein et al., *Detection of the baryon acoustic peak in the large-scale correlation function of SDSS luminous red galaxies*, *Astrophys. J.* **633** (2005) 560 [[astro-ph/0501171](#)] [INSPIRE].
- [7] 2dFGRS collaboration, S. Cole et al., *The 2dF Galaxy Redshift Survey: Power-spectrum analysis of the final dataset and cosmological implications*, *Mon. Not. Roy. Astron. Soc.* **362** (2005) 505 [[astro-ph/0501174](#)] [INSPIRE].
- [8] C. Blake et al., *The WiggleZ Dark Energy Survey: mapping the distance-redshift relation with baryon acoustic oscillations*, *Mon. Not. Roy. Astron. Soc.* **418** (2011) 1707 [[arXiv:1108.2635](#)] [INSPIRE].
- [9] L. Anderson et al., *The clustering of galaxies in the SDSS-III Baryon Oscillation Spectroscopic Survey: Baryon Acoustic Oscillations in the Data Release 9 Spectroscopic Galaxy Sample*, *Mon. Not. Roy. Astron. Soc.* **427** (2012) 3435 [[arXiv:1203.6594](#)] [INSPIRE].
- [10] A. Merloni et al., *eROSITA Science Book: Mapping the Structure of the Energetic Universe*, [arXiv:1209.3114](#) [INSPIRE].
- [11] J.C. Pober et al., *The Baryon Acoustic Oscillation Broadband and Broad-beam Array: Design Overview and Sensitivity Forecasts*, *Astron. J.* **145** (2013) 65 [[arXiv:1210.2413](#)] [INSPIRE].
- [12] G.J. Hill et al., *The Hobby-Eberly Telescope Dark Energy Experiment (HETDEX): Description and Early Pilot Survey Results*, T. Kodama, T. Yamada and K. Aoki eds., *Astr. Soc. P.* **399** (2008) 115 [[arXiv:0806.0183](#)] [INSPIRE].
- [13] D.J. Schlegel et al., *BigBOSS: The Ground-Based Stage IV Dark Energy Experiment*, [arXiv:0904.0468](#) [INSPIRE].
- [14] G. Zhao et al., *The Extended BOSS Survey (eBOSS)*, in preparation (2013).
- [15] A. Slosar et al., *The Lyman- α forest in three dimensions: measurements of large scale flux correlations from BOSS 1st-year data*, *JCAP* **09** (2011) 001 [[arXiv:1104.5244](#)] [INSPIRE].
- [16] SDSS collaboration, C.P. Ahn et al., *The Ninth Data Release of the Sloan Digital Sky Survey: First Spectroscopic Data from the SDSS-III Baryon Oscillation Spectroscopic Survey*, *Astrophys. J. Suppl.* **203** (2012) 21 [[arXiv:1207.7137](#)] [INSPIRE].
- [17] N.G. Busca et al., *Baryon Acoustic Oscillations in the Ly- α forest of BOSS quasars*, *Astron. Astrophys.* **552** (2013) A96 [[arXiv:1211.2616](#)] [INSPIRE].
- [18] A. Slosar et al., *Measurement of Baryon Acoustic Oscillations in the Lyman- α Forest Fluctuations in BOSS Data Release 9*, *JCAP* **04** (2013) 026 [[arXiv:1301.3459](#)] [INSPIRE].
- [19] D. Kirkby et al., *Fitting Methods for Baryon Acoustic Oscillations in the Lyman- α Forest Fluctuations in BOSS Data Release 9*, *JCAP* **03** (2013) 024 [[arXiv:1301.3456](#)] [INSPIRE].
- [20] A. Font-Ribera et al., *The large-scale Quasar-Lyman α Forest Cross-Correlation from BOSS*, *JCAP* **05** (2013) 018 [[arXiv:1303.1937](#)] [INSPIRE].
- [21] M. White et al., *The clustering of intermediate redshift quasars as measured by the Baryon Oscillation Spectroscopic Survey*, *Mon. Not. Roy. Astron. Soc.* **424** (2012) 933 [[arXiv:1203.5306](#)] [INSPIRE].

- [22] A. Lewis, A. Challinor and A. Lasenby, *Efficient computation of CMB anisotropies in closed FRW models*, *Astrophys. J.* **538** (2000) 473 [[astro-ph/9911177](#)] [[INSPIRE](#)].
- [23] D.J. Eisenstein and W. Hu, *Baryonic features in the matter transfer function*, *Astrophys. J.* **496** (1998) 605 [[astro-ph/9709112](#)] [[INSPIRE](#)].
- [24] SDSS collaboration, D.J. Eisenstein et al., *SDSS-III: Massive Spectroscopic Surveys of the Distant Universe, the Milky Way Galaxy and Extra-Solar Planetary Systems*, *Astron. J.* **142** (2011) 72 [[arXiv:1101.1529](#)] [[INSPIRE](#)].
- [25] CUTLER group, LP collaboration, A.S. Bolton et al., *Spectral Classification and Redshift Measurement for the SDSS-III Baryon Oscillation Spectroscopic Survey*, *Astron. J.* **144** (2012) 144 [[arXiv:1207.7326](#)] [[INSPIRE](#)].
- [26] SDSS collaboration, J.E. Gunn et al., *The Sloan digital sky survey photometric camera*, *Astron. J.* **116** (1998) 3040 [[astro-ph/9809085](#)] [[INSPIRE](#)].
- [27] SDSS collaboration, J.E. Gunn et al., *The 2.5 m Telescope of the Sloan Digital Sky Survey*, *Astron. J.* **131** (2006) 2332 [[astro-ph/0602326](#)] [[INSPIRE](#)].
- [28] S.A. Smee et al., *The Multi-Object, Fiber-Fed Spectrographs for SDSS and the Baryon Oscillation Spectroscopic Survey*, *Astron. J.* **146** (2013) 32 [[arXiv:1208.2233](#)] [[INSPIRE](#)].
- [29] SDSS collaboration, D.G. York et al., *The Sloan Digital Sky Survey: Technical Summary*, *Astron. J.* **120** (2000) 1579 [[astro-ph/0006396](#)] [[INSPIRE](#)].
- [30] BOSS collaboration, K.S. Dawson et al., *The Baryon Oscillation Spectroscopic Survey of SDSS-III*, *Astron. J.* **145** (2013) 10 [[arXiv:1208.0022](#)] [[INSPIRE](#)].
- [31] N.P. Ross et al., *The SDSS-III Baryon Oscillation Spectroscopic Survey: Quasar Target Selection for Data Release Nine*, *Astrophys. J. Suppl.* **199** (2012) 3 [[arXiv:1105.0606](#)] [[INSPIRE](#)].
- [32] C.Yèche et al., *Artificial neural networks for quasar selection and photometric redshift determination*, *Astron. Astrophys.* **523** (2010) A14.
- [33] J.A. Kirkpatrick et al., *A Simple Likelihood Method for Quasar Target Selection*, *Astrophys. J.* **743** (2011) 125 [[arXiv:1104.4995](#)] [[INSPIRE](#)].
- [34] J. Bovy et al., *Think Outside the Color-Box: Probabilistic Target Selection and the SDSS-XDQSO Quasar Targeting Catalog*, *Astrophys. J.* **729** (2011) 141 [[arXiv:1011.6392](#)] [[INSPIRE](#)].
- [35] I. Pâris et al., *The Sloan Digital Sky Survey quasar catalog: ninth data release*, *Astron. Astrophys.* **548** (2012) A66 [[arXiv:1210.5166](#)] [[INSPIRE](#)].
- [36] T. Delubac et al., *Lyman alpha BAO constraints from SDSS DR11*, in preparation (2013).
- [37] P. Noterdaeme et al., *Column density distribution and cosmological mass density of neutral gas: Sloan Digital Sky Survey-III Data Release 9*, *Astron. Astrophys.* **547** (2012) L1 [[arXiv:1210.1213](#)] [[INSPIRE](#)].
- [38] P. Noterdaeme, P. Petitjean, C. Ledoux and R. Srianand, *Evolution of the cosmological mass density of neutral gas from Sloan Digital Sky Survey II - Data Release 7*, *Astron. Astrophys.* **505** (2009) 1087 [[arXiv:0908.1574](#)] [[INSPIRE](#)].
- [39] A. Font-Ribera et al., *The large-scale cross-correlation of Damped Lyman Alpha Systems with the Lyman Alpha Forest: First Measurements from BOSS*, *JCAP* **11** (2012) 059 [[arXiv:1209.4596](#)] [[INSPIRE](#)].
- [40] K.-G. Lee, N. Suzuki and D.N. Spergel, *Mean-flux Regulated PCA Continuum Fitting of SDSS Lyman- α Forest Spectra*, *Astron. J.* **143** (2012) 51 [[arXiv:1108.6080](#)] [[INSPIRE](#)].
- [41] N. Kaiser, *Clustering in real space and in redshift space*, *Mon. Not. Roy. Astron. Soc.* **227** (1987) 1 [[INSPIRE](#)].

- [42] B. Efron and G. Gong, *A Leisurely Look at the Bootstrap, the Jackknife, and Cross-Validation*, *Am. Stat.* **37** (1983) 36.
- [43] R. O’Connell et al., *BAO in QSO-LyaF Cross-correlation*, in preparation (2013).
- [44] F. James and M. Roos, *Minuit: A System for Function Minimization and Analysis of the Parameter Errors and Correlations*, *Comput. Phys. Commun.* **10** (1975) 343 [INSPIRE].
- [45] P. McDonald and D.J. Eisenstein, *Dark energy and curvature from a future baryonic acoustic oscillation survey using the Lyman- α forest*, *Phys. Rev. D* **76** (2007) 063009 [[astro-ph/0607122](#)] [INSPIRE].
- [46] M. McQuinn and M. White, *On Estimating Lyman- α Forest Correlations between Multiple Sightlines*, *Mon. Not. Roy. Astron. Soc.* **415** (2011) 2257 [[arXiv:1102.1752](#)] [INSPIRE].
- [47] SDSS collaboration, P. McDonald et al., *The Lyman- α forest power spectrum from the Sloan Digital Sky Survey*, *Astrophys. J. Suppl.* **163** (2006) 80 [[astro-ph/0405013](#)] [INSPIRE].