

Homework 2

Due September 26 at 11 pm

Unless stated otherwise, justify any answers you give. You can work in groups, but each student must write their own solution based on their own understanding of the problem.

When uploading your homework to Gradescope you will have to select the relevant pages for each question. Please submit each problem on a separate page (i.e., 1a and 1b can be on the same page but 1 and 2 must be on different pages). We understand that this may be cumbersome but this is the best way for the grading team to grade your homework assignments and provide feedback in a timely manner. Failure to adhere to these guidelines may result in a loss of points. Note that it may take some time to select the pages for your submission. Please plan accordingly. We suggest uploading your assignment at least 30 minutes before the deadline so you will have ample time to select the correct pages for your submission. If you are using L^AT_EX, consider using the `minted` or `listings` packages for typesetting code.

1. (Geometric random variable) Let \tilde{a} be a geometric random variable with parameter α . What is the probability that \tilde{a} equals a for $a = 1, 2, 3, \dots$ if we condition on the event $\tilde{a} > 5$? Justify your answer mathematically, and also explain why it makes sense intuitively (for example by referring to the coin flip example that we used to derive the geometric pmf).
2. (Chess games) Garry and Anish decide to play 10 chess games. Garry wins 4, they draw 4, and Anish wins 2. We decide to model the games probabilistically, assuming that they are independent and in each game Garry has a probability θ of winning and Anish has a probability α of winning.
 - (a) Plot the log-likelihood function of the parametric model.
 - (b) What is the maximum likelihood estimate of θ and α ?
 - (c) Model the data as realizations from a discrete random variable and compute its empirical pmf. Compare this nonparametric model to the parametric model from the previous questions.
3. (Darts) In a game of darts, a player needs to hit a certain number k times. Assume that all attempts are mutually independent, and that the probability of success in each attempt is θ . Derive the pmf of a random variable representing the number of required attempts.
4. (Call center) Complete the code in the iPython notebook:

https://github.com/cfgranda/prob_stats_for_data_science/blob/main/modeling_discrete_data/call_center_EXERCISE.ipynb

Verify that you reproduce the figures in the notes (no need to submit them), and submit your code for the functions *empirical_pmf* and *poisson_model*.