

Introduction to probability

1. Classical and geometric probability

Wojciech Kotłowski


Institute of Computing Science, PUT

<http://www.cs.put.poznan.pl/wkotlowski/>


2.03.2021

Historical overview

- France, XVII century: gambling is popular and is becoming more complicated
- 1654: known gambler Chevalier de Méré consults **Blaise Pascal** about the chance of winning in a certain game of dice
- Pascal starts to correspond with **Pierre de Fermat** and they jointly formulate the mathematical basis of probability
- The ideas of Pascal and Fermat are developed in the following centuries (e.g., de Moivre, Bernoulli)



Blaise Pascal (1623-1662)



Pierre de Fermat (1601-1665)

Historical overview

- 1814: Pierre Laplace in his book *Théorie analytique des probabilités* formulates the mathematical theory of probability
- Laplace's theory is now known as classical probability
- Based on a principle of indifference:
„Having n mutually-exclusive and collectively exhausting possible outcomes, in the absence of any prior information, assign equal probability to every outcome.”




Pierre Simon de Laplace
(1749-1827)

Sample space

- Every possible result of a **random experiment** (**trial**) is called an **outcome** or an **elementary event** and is denoted with ω

Sample space

- Every possible result of a **random experiment** (**trial**) is called an **outcome** or an **elementary event** and is denoted with ω
 - ▶ Example: rolling a die



Sample space

- Every possible result of a **random experiment** (**trial**) is called an **outcome** or an **elementary event** and is denoted with ω
 - ▶ Example: rolling a die



ω_1



ω_3



ω_5

ω_2



ω_4



ω_6



Sample space

- Every possible result of a **random experiment** (**trial**) is called an **outcome** or an **elementary event** and is denoted with ω


► Example: rolling a die



- The set of all possible outcomes is called a **sample space** Ω

Sample space


- Every possible result of a **random experiment** (**trial**) is called an **outcome** or an **elementary event** and is denoted with ω



- The set of all possible outcomes is called a **sample space** Ω
 - ▶ Example: $\Omega = \{\omega_1, \omega_2, \omega_3, \omega_4, \omega_5, \omega_6\}$.

Sample space


- Every possible result of a **random experiment** (**trial**) is called an **outcome** or an **elementary event** and is denoted with ω



- The set of all possible outcomes is called a **sample space** Ω
 - ▶ Example: $\Omega = \{\omega_1, \omega_2, \omega_3, \omega_4, \omega_5, \omega_6\}$.
- An **event** is a subset of the sample space Ω .
We say that **event A occurred**, if the actual outcome $\omega \in A$

Sample space


- Every possible result of a **random experiment** (trial) is called an **outcome** or an **elementary event** and is denoted with ω



- The set of all possible outcomes is called a **sample space** Ω
 - ▶ Example: $\Omega = \{\omega_1, \omega_2, \omega_3, \omega_4, \omega_5, \omega_6\}$.
- An **event** is a subset of the sample space Ω .
We say that **event A occurred**, if the actual outcome $\omega \in A$
 - ▶ Example: event „roll 1”: $A = \{\omega_1\}$

Sample space


- Every possible result of a **random experiment** (trial) is called an **outcome** or an **elementary event** and is denoted with ω



- The set of all possible outcomes is called a **sample space** Ω
 - ▶ Example: $\Omega = \{\omega_1, \omega_2, \omega_3, \omega_4, \omega_5, \omega_6\}$.
- An **event** is a subset of the sample space Ω .
We say that **event** A **occurred**, if the actual outcome $\omega \in A$
 - ▶ Example: event „roll 1”: $A = \{\omega_1\}$
 - ▶ Example: event „roll at most 2”: $B = \{\omega_1, \omega_2\}$

Sample space

- Every possible result of a **random experiment** (trial) is called an **outcome** or an **elementary event** and is denoted with ω



- The set of all possible outcomes is called a **sample space** Ω
 - ▶ Example: $\Omega = \{\omega_1, \omega_2, \omega_3, \omega_4, \omega_5, \omega_6\}$.
- An **event** is a subset of the sample space Ω .
We say that **event** A **occurred**, if the actual outcome $\omega \in A$
 - ▶ Example: event „roll 1”: $A = \{\omega_1\}$
 - ▶ Example: event „roll at most 2”: $B = \{\omega_1, \omega_2\}$
 - ▶ Example: event „roll even number”: $C = \{\omega_2, \omega_4, \omega_6\}$

Classical probability

- Sample space Ω
- Events $A \subseteq \Omega$ are subsets of the sample space
- The probability of event A :

$$P(A) = \frac{|A|}{|\Omega|}$$

Classical probability

- Sample space Ω
- Events $A \subseteq \Omega$ are subsets of the sample space
- The probability of event A :

$$P(A) = \frac{|A|}{|\Omega|}$$

number of outcomes
in A

total number
of outcomes

Classical probability

- Sample space Ω
- Events $A \subseteq \Omega$ are subsets of the sample space
- The probability of event A :

$$P(A) = \frac{|A|}{|\Omega|}$$

number of outcomes
in A

total number
of outcomes

- Principle of indifference: „every outcome is equally likely”

Classical probability

- Sample space Ω
- Events $A \subseteq \Omega$ are subsets of the sample space
- The probability of event A :

$$P(A) = \frac{|A|}{|\Omega|}$$

number of outcomes
in A

total number
of outcomes


- Principle of indifference: „every outcome is equally likely”
- It holds:

$$P(\emptyset) = 0$$

$$P(\Omega) = 1$$


$$P(A) \in [0, 1]$$

Example – rolling a die




- Sample space $\Omega = \{\omega_1, \omega_2, \omega_3, \omega_4, \omega_5, \omega_6\}$, $|\Omega| = 6$

Example – rolling a die




- Sample space $\Omega = \{\omega_1, \omega_2, \omega_3, \omega_4, \omega_5, \omega_6\}$, $|\Omega| = 6$
- Event „roll 1“:
 $A = \{\omega_1\}$, $|A| = 1$, $P(A) = \frac{1}{6}$

Example – rolling a die



- Sample space $\Omega = \{\omega_1, \omega_2, \omega_3, \omega_4, \omega_5, \omega_6\}$, $|\Omega| = 6$
- Event „roll 1“:
 $A = \{\omega_1\}$, $|A| = 1$, $P(A) = \frac{1}{6}$
- Event „roll at most 2“:
 $B = \{\omega_1, \omega_2\}$, $|B| = 2$, $P(B) = \frac{2}{6} = \frac{1}{3}$

Example – rolling a die



- Sample space $\Omega = \{\omega_1, \omega_2, \omega_3, \omega_4, \omega_5, \omega_6\}$, $|\Omega| = 6$
- Event „roll 1”:
 $A = \{\omega_1\}$, $|A| = 1$, $P(A) = \frac{1}{6}$
- Event „roll at most 2”:
 $B = \{\omega_1, \omega_2\}$, $|B| = 2$, $P(B) = \frac{2}{6} = \frac{1}{3}$
- Event „roll even number”:
 $C = \{\omega_2, \omega_4, \omega_6\}$, $|C| = 3$, $P(A) = \frac{3}{6} = \frac{1}{2}$

Example – tossing three coins



Head



Tail

- Sample space:
- Event „getting three heads”:
- Event „getting (exactly) two heads”:

Example – tossing three coins



Head



Tail

- Sample space:

$$\Omega = \{HHH, HHT, HTH, THH, HTT, THT, TTH, TTT\}$$

$$|\Omega| = 8$$

- Event „getting three heads”:
- Event „getting (exactly) two heads”:

Example – tossing three coins



Head



Tail

- Sample space:

$$\Omega = \{HHH, HHT, HTH, THH, HTT, THT, TTH, TTT\}$$

$$|\Omega| = 8$$

- Event „getting three heads”:

$$A = \{HHH\}, \quad |A| = 1, \quad P(A) = \frac{1}{8}$$

- Event „getting (exactly) two heads”:

Example – tossing three coins



Head



Tail

- Sample space:

$$\Omega = \{HHH, HHT, HTH, THH, HTT, THT, TTH, TTT\}$$

$$|\Omega| = 8$$

- Event „getting three heads”:

$$A = \{HHH\}, \quad |A| = 1, \quad P(A) = \frac{1}{8}$$

- Event „getting (exactly) two heads”:

$$B = \{HHT, HTH, THH\}, \quad |B| = 3, \quad P(B) = \frac{3}{8}$$

Example – rolling two dice

- Sample space:

Example – rolling two dice

- Sample space ($|\Omega| = 36$):

$$\Omega = \{(1, 1), (1, 2), \dots, (1, 6), (2, 1), (2, 2), \dots, (6, 5), (6, 6)\}$$

Example – rolling two dice

- Sample space ($|\Omega| = 36$):

$$\Omega = \{(1, 1), (1, 2), \dots, (1, 6), (2, 1), (2, 2), \dots, (6, 5), (6, 6)\}$$

- Event: „the sum of both dice is S ”

S	Event	Probability
2		
3		
4		
5		
6		
7		
8		
9		
10		
11		
12		

Example – rolling two dice

- Sample space ($|\Omega| = 36$):

$$\Omega = \{(1, 1), (1, 2), \dots, (1, 6), (2, 1), (2, 2), \dots, (6, 5), (6, 6)\}$$

- Event: „the sum of both dice is S ”

S	Event	Probability
2	$A_2 = \{(1, 1)\}$	
3	$A_3 = \{(1, 2), (2, 1)\}$	
4	$A_4 = \{(1, 3), (2, 2), (3, 1)\}$	
5	$A_5 = \{(1, 4), (2, 3), (3, 2), (4, 1)\}$	
6	$A_6 = \{(1, 5), (2, 4), (3, 3), (4, 2), (5, 1)\}$	
7	$A_7 = \{(1, 6), (2, 5), (3, 4), (4, 3), (5, 2), (6, 1)\}$	
8	$A_8 = \{(2, 6), (3, 5), (4, 4), (5, 3), (6, 2)\}$	
9	$A_9 = \{(3, 6), (4, 5), (5, 4), (6, 3)\}$	
10	$A_{10} = \{(4, 6), (5, 5), (6, 4)\}$	
11	$A_{11} = \{(5, 6), (6, 5)\}$	
12	$A_{12} = \{(6, 6)\}$	

Example – rolling two dice

- Sample space ($|\Omega| = 36$):


$$\Omega = \{(1, 1), (1, 2), \dots, (1, 6), (2, 1), (2, 2), \dots, (6, 5), (6, 6)\}$$

- Event: „the sum of both dice is S ”


S	Event	Probability
2	$A_2 = \{(1, 1)\}$	$P(A_2) = 1/36$
3	$A_3 = \{(1, 2), (2, 1)\}$	$P(A_3) = 2/36$
4	$A_4 = \{(1, 3), (2, 2), (3, 1)\}$	$P(A_4) = 3/36$
5	$A_5 = \{(1, 4), (2, 3), (3, 2), (4, 1)\}$	$P(A_5) = 4/36$
6	$A_6 = \{(1, 5), (2, 4), (3, 3), (4, 2), (5, 1)\}$	$P(A_6) = 5/36$
7	$A_7 = \{(1, 6), (2, 5), (3, 4), (4, 3), (5, 2), (6, 1)\}$	$P(A_7) = 6/36$
8	$A_8 = \{(2, 6), (3, 5), (4, 4), (5, 3), (6, 2)\}$	$P(A_8) = 5/36$
9	$A_9 = \{(3, 6), (4, 5), (5, 4), (6, 3)\}$	$P(A_9) = 4/36$
10	$A_{10} = \{(4, 6), (5, 5), (6, 4)\}$	$P(A_{10}) = 3/36$
11	$A_{11} = \{(5, 6), (6, 5)\}$	$P(A_{11}) = 2/36$
12	$A_{12} = \{(6, 6)\}$	$P(A_{12}) = 1/36$

Operations with events


Events are sets!




sum $A \cup B$
„A or B occurred”



intersection $A \cap B$
„A and B occurred”




difference $A \setminus B$
„A occurred, but not B”




complement A'
„A did not occur”


Operations with events




$$A \cup B$$



$$A \cap B$$



$$A \setminus B$$




$$A'$$

$$|A \cup B| = |A| + |B| - |A \cap B|$$


$$|A \setminus B| = |A| - |A \cap B|$$

$$|A'| = |\Omega| - |A|$$


Operations with events




$$A \cup B$$



$$A \cap B$$



$$A \setminus B$$




$$A'$$

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$P(A \setminus B) = P(A) - P(A \cap B)$$


$$P(A') = 1 - P(A)$$

Mutually exclusive (disjoint) events



- If $A \cap B = \emptyset$ then $P(A \cup B) = P(A) + P(B)$

Mutually exclusive (disjoint) events



- If $A \cap B = \emptyset$ then $P(A \cup B) = P(A) + P(B)$
- More generally: if A_1, \dots, A_n are mutually exclusive (disjoint), $A_i \cap A_j = \emptyset$ for $i \neq j$, then:

$$P(A_1 \cup A_2 \cup \dots \cup A_n) = P(A_1) + P(A_2) + \dots + P(A_n)$$

Counting: variation with repetition

If an experiment consists of k independent trials, and there are n possible outcomes in each trial, then the total number of outcomes is

$$\underbrace{n \cdot n \cdot \dots \cdot n}_k = n^k.$$

This is the number of ways one can choose k elements (with repetitions allowed) from a set of n elements (the order of elements matters)

- Number of possible outcomes from rolling 4 dice? $6 \cdot 6 \cdot 6 \cdot 6 = 6^4$
- Number of possible outcomes from tossing 10 coins? $2^{10} = 1024$
- Number of binary sequences of length n ? $2 \cdot 2 \cdot \dots \cdot 2 = 2^n$
- Number of 5-letter words formed from a 26-letter Latin alphabet? 26^5

Counting: variation without repetition

The number of ways one can choose k **distinct** (**without repetition**) elements from a set of n elements (the order of elements **matters**):

$$n \cdot (n-1) \cdot \dots \cdot (n-k+1) = \frac{n!}{(n-k)!}.$$

- Number of ways one can draw 5 cards (without replacement) from a deck of 52 cards (the order of cards matters)? $52 \cdot 51 \cdot 50 \cdot 49 \cdot 48$
- Number of possible outcomes from rolling 3 dice when each die gives a different number? $6 \cdot 5 \cdot 4$
- Number of 5-letter words formed from a 26-letter alphabet in which every letter occurs at most once? $26 \cdot 25 \cdot 24 \cdot 23 \cdot 22$

Counting: permutations

The number of ways one can order a set of n elements:

$$n! = 1 \cdot 2 \cdot \dots \cdot n$$

- Number of ways 5 people can be arranged in a queue? $5! = 120$
- Number of all possible results of shuffling a deck of 52 cards? $52!$

Counting: combination

The number of ways one can choose a k -element subset (the order **does not matter**) from a set of n elements:

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}$$

- The number of ways one can form a group of size 3 from a set of 5 people? $\binom{5}{3} = \frac{5!}{3!2!} = 10$
- 10 teams play with each other. How many matches are going to be played? $\binom{10}{2} = \frac{10!}{8!2!} = 45$
- The number of binary sequences of length 8 with exactly 3 ones? $\binom{8}{3} = \frac{8!}{5!3!} = 56$ (hint: with each binary sequence associate a subset of an n -element set)

Exercise

We draw 5 cards from a deck. What is the probability of getting “four of a kind” (4 cards of the same rank)

Exercise

We draw 5 cards from a deck. What is the probability of getting “four of a kind” (4 cards of the same rank)

Ω – set of all 5-element subsets of 52-element card deck.

$$|\Omega| = \binom{52}{5}$$

Exercise

We draw 5 cards from a deck. What is the probability of getting “four of a kind” (4 cards of the same rank)

Ω – set of all 5-element subsets of 52-element card deck.

$$|\Omega| = \binom{52}{5}$$

Event A – „four of a kind”

Exercise

We draw 5 cards from a deck. What is the probability of getting “four of a kind” (4 cards of the same rank)

Ω – set of all 5-element subsets of 52-element card deck.

$$|\Omega| = \binom{52}{5}$$

Event A – „four of a kind”

$$|A| = 13 \times 48$$

Exercise

We draw 5 cards from a deck. What is the probability of getting “four of a kind” (4 cards of the same rank)

Ω – set of all 5-element subsets of 52-element card deck.

$$|\Omega| = \binom{52}{5}$$

Event A – „four of a kind”

$$|A| = 13 \times 48$$

$$P(A) = \frac{13 \cdot 48}{\binom{52}{5}} \simeq 0.00024$$

Exercise

What is the probability that we get 10 heads in 20 tosses of a fair coin?

Exercise

What is the probability that we get 10 heads in 20 tosses of a fair coin?

Ω – the set of all outcomes of 20 coin tosses

$$|\Omega| = 2^{20}$$

Exercise

What is the probability that we get 10 heads in 20 tosses of a fair coin?

Ω – the set of all outcomes of 20 coin tosses

$$|\Omega| = 2^{20}$$

Event A – „we got exactly 10 heads”

Exercise

What is the probability that we get 10 heads in 20 tosses of a fair coin?

Ω – the set of all outcomes of 20 coin tosses

$$|\Omega| = 2^{20}$$

Event A – „we got exactly 10 heads”

$$|A| = \binom{20}{10}$$

Exercise

What is the probability that we get 10 heads in 20 tosses of a fair coin?

Ω – the set of all outcomes of 20 coin tosses

$$|\Omega| = 2^{20}$$

Event A – „we got exactly 10 heads”

$$|A| = \binom{20}{10}$$

$$P(A) = \frac{\binom{20}{10}}{2^{20}} \simeq 0.176$$

The problem of Chevalier de Méré

What is more likely?

1. Getting at least one 1 in four rolls of a die
2. Getting at least one “double 1” in 24 rolls of a pair of dice?

The problem of Chevalier de Méré

What is more likely?

1. Getting at least one 1 in four rolls of a die
2. Getting at least one “double 1” in 24 rolls of a pair of dice?

According to de Méré:

- The probability of getting a “double 1” ($1/36$) is **six times** less than that of a single 1 ($1/6$)
- To compensate this, one needs to roll a pair of dice **six times more** than one would roll a single die
- Conclusion: both events are **equally likely**

The problem of Chevalier de Méré

What is more likely?

1. Getting at least one 1 in four rolls of a die
2. Getting at least one “double 1” in 24 rolls of a pair of dice?

According to de Méré:

- The probability of getting a “double 1” ($1/36$) is **six times** less than that of a single 1 ($1/6$)
- To compensate this, one needs to roll a pair of dice **six times more** than one would roll a single die
- Conclusion: both events are **equally likely**

This reasoning is actually **wrong!!!**

The problem of Chevalier de Méré

What is more likely?

1. Getting at least one 1 in four rolls of a die
2. Getting at least one “double 1” in 24 rolls of a pair of dice?

Experiment 1:

The problem of Chevalier de Méré

What is more likely?

1. Getting at least one 1 in four rolls of a die
2. Getting at least one “double 1” in 24 rolls of a pair of dice?

Experiment 1:

- Ω : all possible outcomes of rolling 4 dice:
 $|\Omega| = 6 \cdot 6 \cdot 6 \cdot 6 = 6^4$

The problem of Chevalier de Méré

What is more likely?

1. Getting at least one 1 in four rolls of a die
2. Getting at least one “double 1” in 24 rolls of a pair of dice?

Experiment 1:

- Ω : all possible outcomes of rolling 4 dice:
 $|\Omega| = 6 \cdot 6 \cdot 6 \cdot 6 = 6^4$
- A : “at least one 1”

The problem of Chevalier de Méré

What is more likely?

1. Getting at least one 1 in four rolls of a die
2. Getting at least one “double 1” in 24 rolls of a pair of dice?

Experiment 1:

- Ω : all possible outcomes of rolling 4 dice:
 $|\Omega| = 6 \cdot 6 \cdot 6 \cdot 6 = 6^4$
- A : “at least one 1”
- A' : “all numbers different than 1”

The problem of Chevalier de Méré

What is more likely?

1. Getting at least one 1 in four rolls of a die
2. Getting at least one “double 1” in 24 rolls of a pair of dice?

Experiment 1:

- Ω : all possible outcomes of rolling 4 dice:

$$|\Omega| = 6 \cdot 6 \cdot 6 \cdot 6 = 6^4$$

- A : “at least one 1”
- A' : “all numbers different than 1”

$$|A'| = 5 \cdot 5 \cdot 5 \cdot 5 = 5^4, \quad P(A') = \frac{5^4}{6^4}$$

The problem of Chevalier de Méré

What is more likely?

1. Getting at least one 1 in four rolls of a die
2. Getting at least one “double 1” in 24 rolls of a pair of dice?

Experiment 1:

- Ω : all possible outcomes of rolling 4 dice:

$$|\Omega| = 6 \cdot 6 \cdot 6 \cdot 6 = 6^4$$

- A : “at least one 1”

- A' : “all numbers different than 1”

$$|A'| = 5 \cdot 5 \cdot 5 \cdot 5 = 5^4, \quad P(A') = \frac{5^4}{6^4}$$

- Therefore:

$$P(A) = 1 - P(A') = 1 - \frac{5^4}{6^4} \simeq 0.5177$$

The problem of Chevalier de Méré

What is more likely?

1. Getting at least one 1 in four rolls of a die
2. Getting at least one “double 1” in 24 rolls of a pair of dice?

Experiment 2:

- Ω : all possible outcomes of rolling a pair of dice 24 times:
 $|\Omega| = 36^{24}$
- A : “at least one double 1”
- A' : “no double 1 rolled”
 $|A'| = 35^{24}$, $P(A') = \frac{35^{24}}{36^{24}}$
- Therefore:

$$P(A) = 1 - P(A') = 1 - \frac{35^{24}}{36^{24}} \simeq 0.4914$$

Lotto

What is the probability of winning in the Lotto game (we pick 6 out of 49 numbers, the machine draws 6 out of 49 numbers and all the number must match).

Ω – set of 6-element subsets of a set of size 49

Lotto

What is the probability of winning in the Lotto game (we pick 6 out of 49 numbers, the machine draws 6 out of 49 numbers and all the number must match).

Ω – set of 6-element subsets of a set of size 49

$$|\Omega| = \binom{49}{6}$$

Lotto

What is the probability of winning in the Lotto game (we pick 6 out of 49 numbers, the machine draws 6 out of 49 numbers and all the number must match).

Ω – set of 6-element subsets of a set of size 49

$$|\Omega| = \binom{49}{6}$$

Event A – „all six numbers matched”

Lotto

What is the probability of winning in the Lotto game (we pick 6 out of 49 numbers, the machine draws 6 out of 49 numbers and all the number must match).

Ω – set of 6-element subsets of a set of size 49

$$|\Omega| = \binom{49}{6}$$

Event A – „all six numbers matched”

$$|A| = 1 \quad \implies \quad P(A) = \frac{1}{\binom{49}{6}} = \frac{6!43!}{49!} = \frac{1}{13\,983\,816}$$

Not much

(thus, playing Lotto is sometimes called a “tax on dreams”)

Birthday paradox

What is the probability that in a set of 23 people some pair of them will have the same birthday? (for simplicity, assume 365 days in a year)

Birthday paradox

What is the probability that in a set of 23 people some pair of them will have the same birthday? (for simplicity, assume 365 days in a year)

- Number of possible outcomes: $|\Omega| =$

Birthday paradox

What is the probability that in a set of 23 people some pair of them will have the same birthday? (for simplicity, assume 365 days in a year)

- Number of possible outcomes: $|\Omega| = 365^{23}$
- Event A – „at least one pair with the same birthday”

Birthday paradox

What is the probability that in a set of 23 people some pair of them will have the same birthday? (for simplicity, assume 365 days in a year)

- Number of possible outcomes: $|\Omega| = 365^{23}$
- Event A – „at least one pair with the same birthday”
- Event A' – „every person has a different birthday”

Birthday paradox

What is the probability that in a set of 23 people some pair of them will have the same birthday? (for simplicity, assume 365 days in a year)

- Number of possible outcomes: $|\Omega| = 365^{23}$
- Event A – „at least one pair with the same birthday”
- Event A' – „every person has a different birthday”

$$|A'| = 365 \cdot 364 \cdot \dots \cdot (365 - 22)$$

Birthday paradox

What is the probability that in a set of 23 people some pair of them will have the same birthday? (for simplicity, assume 365 days in a year)

- Number of possible outcomes: $|\Omega| = 365^{23}$
- Event A – „at least one pair with the same birthday”
- Event A' – „every person has a different birthday”

$$|A'| = 365 \cdot 364 \cdot \dots \cdot (365 - 22)$$

- Therefore:

$$P(A') = \frac{365 \cdot 364 \cdot \dots \cdot 343}{365^{23}} \simeq 0.493 \quad P(A) = 1 - P(A') \simeq 0.507$$

- The probability is surprisingly large (above 50%)!
- Direct applications to calculating the probability of collision for hash functions

Geometric probability

- Application of the principle of indifference to sample space \mathbb{R}^n
- In complete analogy to classical probability, but:
 - ▶ The outcomes of an experiment are points in \mathbb{R}^n ,
 - ▶ The events are sets in \mathbb{R}^n
 - ▶ The “size” of a set is its n -dimensional **measure**: length ($n = 1$), area ($n = 2$), volume ($n = 3$), etc.
- The concept of geometric probability already considered by Newton in 1666-1668

Geometric probability

- For each subset $A \subset \mathbb{R}^n$, let $|A|$ denote its n -dimensional measure (length for $n = 1$, area for $n = 2$, volume for $n = 3$, etc.)

Geometric probability

- For each subset $A \subset \mathbb{R}^n$, let $|A|$ denote its n -dimensional measure (length for $n = 1$, area for $n = 2$, volume for $n = 3$, etc.)
- Sample space $\Omega \subset \mathbb{R}^n$, where $|\Omega| < \infty$
- Events $A \subset \Omega$ are subsets of Ω

Geometric probability

- For each subset $A \subset \mathbb{R}^n$, let $|A|$ denote its n -dimensional measure (length for $n = 1$, area for $n = 2$, volume for $n = 3$, etc.)
- Sample space $\Omega \subset \mathbb{R}^n$, where $|\Omega| < \infty$
- Events $A \subset \Omega$ are subsets of Ω
- The probability of A :

$$P(A) = \frac{|A|}{|\Omega|}$$

Geometric probability

- For each subset $A \subset \mathbb{R}^n$, let $|A|$ denote its n -dimensional measure (length for $n = 1$, area for $n = 2$, volume for $n = 3$, etc.)
- Sample space $\Omega \subset \mathbb{R}^n$, where $|\Omega| < \infty$
- Events $A \subset \Omega$ are subsets of Ω
- The probability of A :

$$P(A) = \frac{|A|}{|\Omega|}$$

- Interpretation:
 - ▶ Classical probability: „each element in Ω equally likely”
 - ▶ Geometric probability: „each point in Ω equally likely”

Geometric probability

- For each subset $A \subset \mathbb{R}^n$, let $|A|$ denote its n -dimensional measure (length for $n = 1$, area for $n = 2$, volume for $n = 3$, etc.)
- Sample space $\Omega \subset \mathbb{R}^n$, where $|\Omega| < \infty$
- Events $A \subset \Omega$ are subsets of Ω
- The probability of A :

$$P(A) = \frac{|A|}{|\Omega|}$$

- Interpretation:
 - ▶ Classical probability: „each element in Ω equally likely”
 - ▶ Geometric probability: „each point in Ω equally likely”
- Geometric probability has identical properties as its classical counterpart, e.g. $P(\emptyset) = 0$, $P(A') = 1 - P(A)$,
 $P(A \cup B) = P(A) + P(B) - P(A \cap B)$, itp.


Geometric probability: example

We random draw a point from interval $[-1, 1]$. What is the probability that the point is contained within the interval $[0.1, 0.5]$?



Geometric probability: example

We random draw a point from interval $[-1, 1]$. What is the probability that the point is contained within the interval $[0.1, 0.5]$?




Answer:

$$\Omega = [-1, 1], A = [0.1, 0.5]$$

$$P(A) = \frac{|A|}{|\Omega|} = \frac{0.4}{2} = 0.2$$

Geometric probability: example

We random draw a point from interval $[-1, 1]$. What is the probability that the point is contained within the interval $[0.1, 0.5]$?




Answer:

$$\Omega = [-1, 1], A = [0.1, 0.5]$$

$$P(A) = \frac{|A|}{|\Omega|} = \frac{0.4}{2} = 0.2$$

What is the probability that we draw a point exactly equal to zero?



$$A = \{0\}, P(A) = \frac{0}{2} = 0$$


Geometric probability: example

We draw two points x_1 and x_2 from a unit interval $[0, 1]$. Calculate the probability that $x_1 > x_2$?

Geometric probability: example

We draw two points x_1 and x_2 from a unit interval $[0, 1]$. Calculate the probability that $x_1 > x_2$?

Sample space: set of pairs (x_1, x_2)



$$\Omega = [0, 1] \times [0, 1] = [0, 1]^2$$

Geometric probability: example


We draw two points x_1 and x_2 from a unit interval $[0, 1]$. Calculate the probability that $x_1 > x_2$?

Sample space: set of pairs (x_1, x_2)

$$\Omega = [0, 1] \times [0, 1] = [0, 1]^2$$

The event we are interested in:

$$A = \{(x_1, x_2) \in \Omega : x_1 > x_2\}$$



Geometric probability: example


We draw two points x_1 and x_2 from a unit interval $[0, 1]$. Calculate the probability that $x_1 > x_2$?

Sample space: set of pairs (x_1, x_2)

$$\Omega = [0, 1] \times [0, 1] = [0, 1]^2$$

The event we are interested in:

$$A = \{(x_1, x_2) \in \Omega : x_1 > x_2\}$$




$$P(A) = \frac{|A|}{|\Omega|} = \frac{1/2}{1} = \frac{1}{2}$$

Geometric probability: example

Alice and Bob go on a date. The first person to show up will only wait up to 20 minutes for the other. What is the probability that they meet if they arrive independently at a random moment between 10:00 and 11:00?

Geometric probability: example

Alice and Bob go on a date. The first person to show up will only wait up to 20 minutes for the other. What is the probability that they meet if they arrive independently at a random moment between 10:00 and 11:00?




Let $x, y \in [0, 60]$ denote the times of arrival (in minutes) of Alice and Bob counting from 10:00.

$$\Omega = [0, 60] \times [0, 60]$$

Geometric probability: example

Alice and Bob go on a date. The first person to show up will only wait up to 20 minutes for the other. What is the probability that they meet if they arrive independently at a random moment between 10:00 and 11:00?




Let $x, y \in [0, 60]$ denote the times of arrival (in minutes) of Alice and Bob counting from 10:00.

$$\Omega = [0, 60] \times [0, 60]$$

$$A = \{(x, y) \in \Omega : |x - y| \leq 20\} \text{ (red area)}$$

Geometric probability: example

Alice and Bob go on a date. The first person to show up will only wait up to 20 minutes for the other. What is the probability that they meet if they arrive independently at a random moment between 10:00 and 11:00?



Let $x, y \in [0, 60]$ denote the times of arrival (in minutes) of Alice and Bob counting from 10:00.

$$\Omega = [0, 60] \times [0, 60]$$

$$A = \{(x, y) \in \Omega : |x - y| \leq 20\} \text{ (red area)}$$


$$P(A) = 1 - P(A') = 1 - \frac{|A'|}{|\Omega|} = 1 - \frac{40 \times 40}{60 \times 60} = \frac{5}{9}.$$

Geometric probability: Buffon's needle

We drop a needle of length ℓ onto the floor made of parallel strips of wood of the same width $a \geq \ell$. What is the probability that the needle will lie across a line between two strips?

Geometric probability: Buffon's needle

We drop a needle of length ℓ onto the floor made of parallel strips of wood of the same width $a \geq \ell$. What is the probability that the needle will lie across a line between two strips?




Outcomes are pairs (x, α) :

x – distance from the center of the needle to the closest line, $x \in [0, a/2]$

α – acute angle between the needle and one of the lines, $\alpha \in [0, \frac{\pi}{2}]$

Geometric probability: Buffon's needle

We drop a needle of length ℓ onto the floor made of parallel strips of wood of the same width $a \geq \ell$. What is the probability that the needle will lie across a line between two strips?



Outcomes are pairs (x, α) :


x – distance from the center of the needle to the closest line, $x \in [0, a/2]$

α – acute angle between the needle and one of the lines, $\alpha \in [0, \frac{\pi}{2}]$

$$\Omega = [0, a/2] \times [0, \frac{\pi}{2}], |\Omega| = \frac{a\pi}{4}$$

Geometric probability: Buffon's needle

We drop a needle of length ℓ onto the floor made of parallel strips of wood of the same width $a \geq \ell$. What is the probability that the needle will lie across a line between two strips?



Outcomes are pairs (x, α) :

x – distance from the center of the needle to the closest line, $x \in [0, a/2]$


α – acute angle between the needle and one of the lines, $\alpha \in [0, \frac{\pi}{2}]$

$$\Omega = [0, a/2] \times [0, \frac{\pi}{2}], |\Omega| = \frac{a\pi}{4}$$

The needle will lie across a line if $x \leq d = \frac{\ell}{2} \sin \alpha$ (event A)

Geometric probability: Buffon's needle

We drop a needle of length ℓ onto the floor made of parallel strips of wood of the same width $a \geq \ell$. What is the probability that the needle will lie across a line between two strips?




Outcomes are pairs (x, α) :

x – distance from the center of the needle to the closest line, $x \in [0, a/2]$

α – acute angle between the needle and one of the lines, $\alpha \in [0, \frac{\pi}{2}]$


$$\Omega = [0, a/2] \times [0, \frac{\pi}{2}], |\Omega| = \frac{a\pi}{4}$$

The needle will lie across a line if $x \leq d = \frac{\ell}{2} \sin \alpha$ (event A)



Geometric probability: Buffon's needle

We drop a needle of length ℓ onto the floor made of parallel strips of wood of the same width $a \geq \ell$. What is the probability that the needle will lie across a line between two strips?




Outcomes are **pairs** (x, α) :

x – distance from the center of the needle to the closest line, $x \in [0, a/2]$

α – acute angle between the needle and one of the lines, $\alpha \in [0, \frac{\pi}{2}]$

$$\Omega = [0, a/2] \times [0, \frac{\pi}{2}], |\Omega| = \frac{a\pi}{4}$$


The needle will lie across a line if $x \leq d = \frac{\ell}{2} \sin \alpha$ (event A)



$$P(A) = \frac{|A|}{|\Omega|} = \frac{4}{a\pi} \frac{\ell}{2} \underbrace{\int_0^{\pi/2} \sin \alpha \, d\alpha}_{=1} = \frac{2\ell}{a\pi}$$

Geometric probability: Buffon's needle

We drop a needle of length ℓ onto the floor made of parallel strips of wood of the same width $a \geq \ell$. What is the probability that the needle will lie across a line between two strips?




Outcomes are pairs (x, α) :

x – distance from the center of the needle to the closest line, $x \in [0, a/2]$

α – acute angle between the needle and one of the lines, $\alpha \in [0, \frac{\pi}{2}]$

$$\Omega = [0, a/2] \times [0, \frac{\pi}{2}], |\Omega| = \frac{a\pi}{4}$$

The needle will lie across a line if $x \leq d = \frac{\ell}{2} \sin \alpha$ (event A)



$$P(A) = \frac{|\Omega_A|}{|\Omega|} = \frac{4}{a\pi} \frac{\ell}{2} \underbrace{\int_0^{\pi/2} \sin \alpha \, d\alpha}_{=1} = \frac{2\ell}{a\pi}$$

If $\ell = \frac{a}{2}$, $P(A) = \frac{1}{\pi}$. Can be used to experimentally estimate the number π !