

Linear Least-Squares Problem (LSP)

QR Method

Rafikul Alam
Department of Mathematics
Indian Institute of Technology Guwahati
Guwahati - 781039, INDIA

Outline

- QR method for LSP

Special matrices

Complex matrix	Real matrix
Hermitian: $A^* = A$	Symmetric: $A^T = A$
Unitary: $AA^* = A^*A = I$	Orthogonal: $AA^T = A^T A = I$
Isometry: $A^*A = I$	Isometry: $A^T A = I$

Special matrices

Complex matrix	Real matrix
Hermitian: $A^* = A$	Symmetric: $A^T = A$
Unitary: $AA^* = A^*A = I$	Orthogonal: $AA^T = A^T A = I$
Isometry: $A^*A = I$	Isometry: $A^T A = I$

Fact: An $n \times n$ matrix A is unitary (resp., orthogonal) if and only if columns of A are orthonormal.

Special matrices

Complex matrix	Real matrix
Hermitian: $A^* = A$	Symmetric: $A^T = A$
Unitary: $AA^* = A^*A = I$	Orthogonal: $AA^T = A^T A = I$
Isometry: $A^*A = I$	Isometry: $A^T A = I$

Fact: An $n \times n$ matrix A is unitary (resp., orthogonal) if and only if columns of A are orthonormal. An $m \times n$ matrix is isometry if and only if columns of A are orthonormal.

Special matrices

Complex matrix	Real matrix
Hermitian: $A^* = A$	Symmetric: $A^\top = A$
Unitary: $AA^* = A^*A = I$	Orthogonal: $AA^\top = A^\top A = I$
Isometry: $A^*A = I$	Isometry: $A^\top A = I$

Fact: An $n \times n$ matrix A is unitary (resp., orthogonal) if and only if columns of A are orthonormal. An $m \times n$ matrix is isometry if and only if columns of A are orthonormal.

Example: The matrix $U := \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{i}{\sqrt{2}} \\ i & \frac{1}{\sqrt{2}} \end{bmatrix}$ is unitary and $P := \begin{bmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix}$ is orthogonal.

Special matrices

Complex matrix	Real matrix
Hermitian: $A^* = A$	Symmetric: $A^T = A$
Unitary: $AA^* = A^*A = I$	Orthogonal: $AA^T = A^T A = I$
Isometry: $A^*A = I$	Isometry: $A^T A = I$

Fact: An $n \times n$ matrix A is unitary (resp., orthogonal) if and only if columns of A are orthonormal. An $m \times n$ matrix is isometry if and only if columns of A are orthonormal.

Example: The matrix $U := \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{i}{\sqrt{2}} \\ i & \frac{1}{\sqrt{2}} \end{bmatrix}$ is unitary and $P := \begin{bmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix}$ is orthogonal. The

matrix $Q := \begin{bmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{3}} \\ 0 & \frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{3}} \end{bmatrix}$ is an isometry.

Special matrices

Complex matrix	Real matrix
Hermitian: $A^* = A$	Symmetric: $A^T = A$
Unitary: $AA^* = A^*A = I$	Orthogonal: $AA^T = A^T A = I$
Isometry: $A^*A = I$	Isometry: $A^T A = I$

Fact: An $n \times n$ matrix A is unitary (resp., orthogonal) if and only if columns of A are orthonormal. An $m \times n$ matrix is isometry if and only if columns of A are orthonormal.

Example: The matrix $U := \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{i}{\sqrt{2}} \\ \frac{i}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix}$ is unitary and $P := \begin{bmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix}$ is orthogonal. The matrix $Q := \begin{bmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{3}} \\ 0 & \frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{3}} \end{bmatrix}$ is an isometry.

Exercise: An $n \times n$ matrix U is unitary (resp., orthogonal) if and only if $\langle Ux, Uy \rangle = \langle x, y \rangle$ for all x and y . An $m \times n$ matrix Q is an isometry if and only if $\langle Qx, Qy \rangle = \langle x, y \rangle$ for all x and y .

QR factorization

Theorem: Let $A \in \mathbb{C}^{m \times n}$. Then there is a unitary matrix $Q \in \mathbb{C}^{m \times m}$ and an upper triangular matrix $\mathcal{R} \in \mathbb{C}^{m \times n}$ such that $A = Q\mathcal{R}$.

QR factorization

Theorem: Let $A \in \mathbb{C}^{m \times n}$. Then there is a unitary matrix $Q \in \mathbb{C}^{m \times m}$ and an upper triangular matrix $\mathcal{R} \in \mathbb{C}^{m \times n}$ such that $A = Q\mathcal{R}$. If $\text{rank}(A) = n$ then \mathcal{R} is of the form $\mathcal{R} = \begin{bmatrix} R \\ 0 \end{bmatrix}$ for some nonsingular upper triangular matrix $R \in \mathbb{C}^{n \times n}$.

QR factorization

Theorem: Let $A \in \mathbb{C}^{m \times n}$. Then there is a unitary matrix $Q \in \mathbb{C}^{m \times m}$ and an upper triangular matrix $\mathcal{R} \in \mathbb{C}^{m \times n}$ such that $A = Q\mathcal{R}$. If $\text{rank}(A) = n$ then \mathcal{R} is of the form $\mathcal{R} = \begin{bmatrix} R \\ 0 \end{bmatrix}$ for some nonsingular upper triangular matrix $R \in \mathbb{C}^{n \times n}$. Let $Q = \begin{bmatrix} Q_n & Q_{m-n} \end{bmatrix}$ with $Q_n \in \mathbb{C}^{m \times n}$. Then Q_n is an isometry and

$$A = Q\mathcal{R} = \begin{bmatrix} Q_n & Q_{m-n} \end{bmatrix} \begin{bmatrix} R \\ 0 \end{bmatrix} = Q_n R. \quad \blacksquare$$

QR factorization

Theorem: Let $A \in \mathbb{C}^{m \times n}$. Then there is a unitary matrix $Q \in \mathbb{C}^{m \times m}$ and an upper triangular matrix $\mathcal{R} \in \mathbb{C}^{m \times n}$ such that $A = Q\mathcal{R}$. If $\text{rank}(A) = n$ then \mathcal{R} is of the form $\mathcal{R} = \begin{bmatrix} R \\ 0 \end{bmatrix}$ for some nonsingular upper triangular matrix $R \in \mathbb{C}^{n \times n}$. Let $Q = \begin{bmatrix} Q_n & Q_{m-n} \end{bmatrix}$ with $Q_n \in \mathbb{C}^{m \times n}$. Then Q_n is an isometry and

$$A = Q\mathcal{R} = \begin{bmatrix} Q_n & Q_{m-n} \end{bmatrix} \begin{bmatrix} R \\ 0 \end{bmatrix} = Q_n R. \blacksquare$$

Remark: The factorization $A = Q\mathcal{R}$ is called a full QR factorization and $A = Q_n R$ is called a compact (or economy size) QR factorization of A .

QR factorization

Theorem: Let $A \in \mathbb{C}^{m \times n}$. Then there is a unitary matrix $Q \in \mathbb{C}^{m \times m}$ and an upper triangular matrix $\mathcal{R} \in \mathbb{C}^{m \times n}$ such that $A = Q\mathcal{R}$. If $\text{rank}(A) = n$ then \mathcal{R} is of the form $\mathcal{R} = \begin{bmatrix} R \\ 0 \end{bmatrix}$ for some nonsingular upper triangular matrix $R \in \mathbb{C}^{n \times n}$. Let $Q = [Q_n \quad Q_{m-n}]$ with $Q_n \in \mathbb{C}^{m \times n}$. Then Q_n is an isometry and

$$A = Q\mathcal{R} = [Q_n \quad Q_{m-n}] \begin{bmatrix} R \\ 0 \end{bmatrix} = Q_n R. \quad \blacksquare$$

Remark: The factorization $A = Q\mathcal{R}$ is called a **full QR factorization** and $A = Q_n R$ is called a **compact (or economy size) QR factorization** of A . MATLAB commands: $[Q, R] = \text{qr}(A)$ and $[Q, R] = \text{qr}(A, 0)$, respectively, compute full and compact QR factorization of A

QR factorization

Theorem: Let $A \in \mathbb{C}^{m \times n}$. Then there is a unitary matrix $Q \in \mathbb{C}^{m \times m}$ and an upper triangular matrix $\mathcal{R} \in \mathbb{C}^{m \times n}$ such that $A = Q\mathcal{R}$. If $\text{rank}(A) = n$ then \mathcal{R} is of the form $\mathcal{R} = \begin{bmatrix} R \\ 0 \end{bmatrix}$ for some nonsingular upper triangular matrix $R \in \mathbb{C}^{n \times n}$. Let $Q = [Q_n \quad Q_{m-n}]$ with $Q_n \in \mathbb{C}^{m \times n}$. Then Q_n is an isometry and

$$A = Q\mathcal{R} = [Q_n \quad Q_{m-n}] \begin{bmatrix} R \\ 0 \end{bmatrix} = Q_n R. \blacksquare$$

Remark: The factorization $A = Q\mathcal{R}$ is called a **full QR factorization** and $A = Q_n R$ is called a **compact (or economy size) QR factorization** of A . MATLAB commands: $[Q,R] = \text{qr}(A)$ and $[Q,R] = \text{qr}(A,0)$, respectively, compute full and compact QR factorization of A

Example:

$$\underbrace{\begin{bmatrix} 1 & 3 & 1 \\ 1 & 3 & 7 \\ 1 & -1 & -4 \\ 1 & -1 & 2 \end{bmatrix}}_A = \frac{1}{2} \underbrace{\begin{bmatrix} 1 & 1 & -1 & -1 \\ 1 & 1 & 1 & 1 \\ 1 & -1 & -1 & 1 \\ 1 & -1 & 1 & -1 \end{bmatrix}}_Q \underbrace{\begin{bmatrix} 2 & 2 & 3 \\ 0 & 4 & 5 \\ 0 & 0 & 6 \\ 0 & 0 & 0 \end{bmatrix}}_{\mathcal{R}}.$$

QR method for LSP

A QR factorization of A provides an efficient method for solution of the LSP $Ax \approx b$.

QR method for LSP

A QR factorization of A provides an efficient method for solution of the LSP $Ax \approx b$.

Suppose that $\text{rank}(A) = n$. Set $\begin{bmatrix} c \\ d \end{bmatrix} := Q^*b$, where $c \in \mathbb{C}^n$ and $d \in \mathbb{C}^{m-n}$. Then

$$\|Ax - b\|_2 = \|Q^*(Ax - b)\|_2 = \left\| \begin{bmatrix} R \\ 0 \end{bmatrix} x - \begin{bmatrix} c \\ d \end{bmatrix} \right\|_2 = \sqrt{\|Rx - c\|_2^2 + \|d\|_2^2}.$$

QR method for LSP

A QR factorization of A provides an efficient method for solution of the LSP $Ax \approx b$.

Suppose that $\text{rank}(A) = n$. Set $\begin{bmatrix} c \\ d \end{bmatrix} := Q^*b$, where $c \in \mathbb{C}^n$ and $d \in \mathbb{C}^{m-n}$. Then

$$\|Ax - b\|_2 = \|Q^*(Ax - b)\|_2 = \left\| \begin{bmatrix} R \\ 0 \end{bmatrix} x - \begin{bmatrix} c \\ d \end{bmatrix} \right\|_2 = \sqrt{\|Rx - c\|_2^2 + \|d\|_2^2}.$$

This shows that $\min \|Ax - b\|_2 = \|d\|_2 \iff Rx = c$. Hence $x = R^{-1}c$ is a unique solution of the LSP and $\|d\|_2$ is the residual.

QR method for LSP

A QR factorization of A provides an efficient method for solution of the LSP $Ax \approx b$.

Suppose that $\text{rank}(A) = n$. Set $\begin{bmatrix} c \\ d \end{bmatrix} := Q^*b$, where $c \in \mathbb{C}^n$ and $d \in \mathbb{C}^{m-n}$. Then

$$\|Ax - b\|_2 = \|Q^*(Ax - b)\|_2 = \left\| \begin{bmatrix} R \\ 0 \end{bmatrix} x - \begin{bmatrix} c \\ d \end{bmatrix} \right\|_2 = \sqrt{\|Rx - c\|_2^2 + \|d\|_2^2}.$$

This shows that $\min \|Ax - b\|_2 = \|d\|_2 \iff Rx = c$. Hence $x = R^{-1}c$ is a unique solution of the LSP and $\|d\|_2$ is the residual. [If $Q \in \mathbb{C}^{n \times n}$ is unitary then $\|Qx\|_2 = \|x\|_2$ for all x .]

QR method for LSP

A QR factorization of A provides an efficient method for solution of the LSP $Ax \approx b$.

Suppose that $\text{rank}(A) = n$. Set $\begin{bmatrix} c \\ d \end{bmatrix} := Q^*b$, where $c \in \mathbb{C}^n$ and $d \in \mathbb{C}^{m-n}$. Then

$$\|Ax - b\|_2 = \|Q^*(Ax - b)\|_2 = \left\| \begin{bmatrix} R \\ 0 \end{bmatrix} x - \begin{bmatrix} c \\ d \end{bmatrix} \right\|_2 = \sqrt{\|Rx - c\|_2^2 + \|d\|_2^2}.$$

This shows that $\min \|Ax - b\|_2 = \|d\|_2 \iff Rx = c$. Hence $x = R^{-1}c$ is a unique solution of the LSP and $\|d\|_2$ is the residual. [If $Q \in \mathbb{C}^{n \times n}$ is unitary then $\|Qx\|_2 = \|x\|_2$ for all x .]

Algorithm: Solution of LSP $Ax \approx b$ when $\text{rank}(A) = n$.

1. Compute QR factorization $A = Q \begin{bmatrix} R \\ 0 \end{bmatrix}$.
2. Set $\begin{bmatrix} c \\ d \end{bmatrix} := Q^*b$, where $c \in \mathbb{C}^n$ and $d \in \mathbb{C}^{m-n}$.
3. Solve upper triangular system $Rx = c$.
4. Compute the residual $\|d\|_2$.

Example

Given $A = \begin{bmatrix} 3 & -6 \\ 4 & -8 \\ 0 & 1 \end{bmatrix}$ and $b = \begin{bmatrix} -1 \\ 7 \\ 2 \end{bmatrix}$, solve the LSP $Ax \approx b$.

1. Compute QR factorization: $A = \begin{bmatrix} -\frac{3}{5} & 0 & \frac{4}{5} \\ -\frac{4}{5} & 0 & -\frac{3}{5} \\ 0 & -1 & 0 \end{bmatrix} \begin{bmatrix} -5 & 10 \\ 0 & -1 \\ \hline 0 & 0 \end{bmatrix} = Q \begin{bmatrix} R \\ 0 \end{bmatrix}.$

Example

Given $A = \begin{bmatrix} 3 & -6 \\ 4 & -8 \\ 0 & 1 \end{bmatrix}$ and $b = \begin{bmatrix} -1 \\ 7 \\ 2 \end{bmatrix}$, solve the LSP $Ax \approx b$.

1. Compute QR factorization: $A = \begin{bmatrix} -\frac{3}{5} & 0 & \frac{4}{5} \\ -\frac{4}{5} & 0 & -\frac{3}{5} \\ 0 & -1 & 0 \end{bmatrix} \begin{bmatrix} -5 & 10 \\ 0 & -1 \\ \hline 0 & 0 \end{bmatrix} = Q \begin{bmatrix} R \\ 0 \end{bmatrix}.$
2. Compute $Q^T b = \begin{bmatrix} -5 \\ -2 \\ -5 \end{bmatrix}.$

Example

Given $A = \begin{bmatrix} 3 & -6 \\ 4 & -8 \\ 0 & 1 \end{bmatrix}$ and $b = \begin{bmatrix} -1 \\ 7 \\ 2 \end{bmatrix}$, solve the LSP $Ax \approx b$.

1. Compute QR factorization: $A = \begin{bmatrix} -\frac{3}{5} & 0 & \frac{4}{5} \\ -\frac{4}{5} & 0 & -\frac{3}{5} \\ 0 & -1 & 0 \end{bmatrix} \begin{bmatrix} -5 & 10 \\ 0 & -1 \\ 0 & 0 \end{bmatrix} = Q \begin{bmatrix} R \\ 0 \end{bmatrix}$.
2. Compute $Q^T b = \begin{bmatrix} -5 \\ -2 \\ -5 \end{bmatrix}$.
3. Solve $\begin{bmatrix} -5 & 10 \\ 0 & -1 \end{bmatrix} x = \begin{bmatrix} -5 \\ -2 \end{bmatrix} \Rightarrow x = \begin{bmatrix} 5 \\ 2 \end{bmatrix}$.
4. The residual $\|d\|_2 = 5$.

QR factorization of augmented matrix

If the matrix Q in the QR factorization $A = QR$ is not required, then the LSP $Ax \approx b$ can be solved more efficiently by computing QR factorization of the augmented matrix $\begin{bmatrix} A & b \end{bmatrix}$.

QR factorization of augmented matrix

If the matrix Q in the QR factorization $A = QR$ is not required, then the LSP $Ax \approx b$ can be solved more efficiently by computing QR factorization of the augmented matrix $[A \ b]$.

Suppose that $\text{rank}(A) = n$. Then

$$Ax - b = [A \ b] \begin{bmatrix} x \\ -1 \end{bmatrix} \text{ and } [A \ b] = Q \left[\begin{array}{c|c} R & c \\ \hline \mathbf{0} & d \end{array} \right] = QR, \text{ where } d \in \mathbb{C}.$$

QR factorization of augmented matrix

If the matrix Q in the QR factorization $A = QR$ is not required, then the LSP $Ax \approx b$ can be solved more efficiently by computing QR factorization of the augmented matrix $[A \ b]$.

Suppose that $\text{rank}(A) = n$. Then

$$Ax - b = [A \ b] \begin{bmatrix} x \\ -1 \end{bmatrix} \text{ and } [A \ b] = Q \left[\begin{array}{c|c} R & c \\ 0 & d \\ \hline & 0 \end{array} \right] = QR, \text{ where } d \in \mathbb{C}.$$

Hence $\|Ax - b\|_2 = \sqrt{\|Rx - c\|_2^2 + |d|^2} \implies \min \|Ax - b\|_2 = |d| \iff Rx = c$.

QR factorization of augmented matrix

If the matrix Q in the QR factorization $A = QR$ is not required, then the LSP $Ax \approx b$ can be solved more efficiently by computing QR factorization of the augmented matrix $[A \ b]$.

Suppose that $\text{rank}(A) = n$. Then

$$Ax - b = [A \ b] \begin{bmatrix} x \\ -1 \end{bmatrix} \text{ and } [A \ b] = Q \left[\begin{array}{c|c} R & c \\ 0 & d \\ \hline & 0 \end{array} \right] = QR, \text{ where } d \in \mathbb{C}.$$

Hence $\|Ax - b\|_2 = \sqrt{\|Rx - c\|_2^2 + |d|^2} \implies \min \|Ax - b\|_2 = |d| \iff Rx = c$.

Hence the LSP $Ax \approx b$ can be solved in three steps:

QR factorization of augmented matrix

If the matrix Q in the QR factorization $A = QR$ is not required, then the LSP $Ax \approx b$ can be solved more efficiently by computing QR factorization of the augmented matrix $[A \ b]$.

Suppose that $\text{rank}(A) = n$. Then

$$Ax - b = [A \ b] \begin{bmatrix} x \\ -1 \end{bmatrix} \text{ and } [A \ b] = Q \left[\begin{array}{c|c} R & c \\ \hline 0 & d \end{array} \right] = QR, \text{ where } d \in \mathbb{C}.$$

Hence $\|Ax - b\|_2 = \sqrt{\|Rx - c\|_2^2 + |d|^2} \implies \min \|Ax - b\|_2 = |d| \iff Rx = c$.

Hence the LSP $Ax \approx b$ can be solved in three steps:

- Compute QR factorization $[A, \ b] = QR$.

QR factorization of augmented matrix

If the matrix Q in the QR factorization $A = QR$ is not required, then the LSP $Ax \approx b$ can be solved more efficiently by computing QR factorization of the augmented matrix $[A \ b]$.

Suppose that $\text{rank}(A) = n$. Then

$$Ax - b = [A \ b] \begin{bmatrix} x \\ -1 \end{bmatrix} \text{ and } [A \ b] = Q \left[\begin{array}{c|c} R & c \\ \hline 0 & d \end{array} \right] = QR, \text{ where } d \in \mathbb{C}.$$

Hence $\|Ax - b\|_2 = \sqrt{\|Rx - c\|_2^2 + |d|^2} \implies \min \|Ax - b\|_2 = |d| \iff Rx = c$.

Hence the LSP $Ax \approx b$ can be solved in three steps:

- Compute QR factorization $[A, \ b] = QR$.
- Solve the upper triangular system $R(1:n, 1:n)x = R(1:n, n+1)$.

QR factorization of augmented matrix

If the matrix Q in the QR factorization $A = QR$ is not required, then the LSP $Ax \approx b$ can be solved more efficiently by computing QR factorization of the augmented matrix $[A \ b]$.

Suppose that $\text{rank}(A) = n$. Then

$$Ax - b = [A \ b] \begin{bmatrix} x \\ -1 \end{bmatrix} \text{ and } [A \ b] = Q \left[\begin{array}{c|c} R & c \\ 0 & d \\ \hline & 0 \end{array} \right] = QR, \text{ where } d \in \mathbb{C}.$$

Hence $\|Ax - b\|_2 = \sqrt{\|Rx - c\|_2^2 + |d|^2} \implies \min \|Ax - b\|_2 = |d| \iff Rx = c$.

Hence the LSP $Ax \approx b$ can be solved in three steps:

- Compute QR factorization $[A, \ b] = QR$.
- Solve the upper triangular system $R(1:n, 1:n)x = R(1:n, n+1)$.
- Compute residual norm $|d| = \text{abs}(R(n+1, n+1))$.

QR method for rank deficient LSP

Theorem: Let $A \in \mathbb{C}^{m \times n}$. Suppose that $\text{rank}(A) = r$. Then there is a unitary matrix $Q \in \mathbb{C}^{m \times m}$ and a nonsingular upper triangular matrix $R_{11} \in \mathbb{C}^{r \times r}$ such that

$$AP = Q \begin{bmatrix} R_{11} & R_{12} \\ 0 & 0 \end{bmatrix} = QR,$$

where $P \in \mathbb{R}^{n \times n}$ is a permutation matrix and $R_{12} \in \mathbb{C}^{r \times (n-r)}$. ■

QR method for rank deficient LSP

Theorem: Let $A \in \mathbb{C}^{m \times n}$. Suppose that $\text{rank}(A) = r$. Then there is a unitary matrix $Q \in \mathbb{C}^{m \times m}$ and a nonsingular upper triangular matrix $R_{11} \in \mathbb{C}^{r \times r}$ such that

$$AP = Q \begin{bmatrix} R_{11} & R_{12} \\ 0 & 0 \end{bmatrix} = QR,$$

where $P \in \mathbb{R}^{n \times n}$ is a permutation matrix and $R_{12} \in \mathbb{C}^{r \times (n-r)}$. ■

Set $\begin{bmatrix} c \\ d \end{bmatrix} := Q^* b$, where $c \in \mathbb{C}^r$ and $d \in \mathbb{C}^{m-r}$. Then

$$\begin{aligned} \|Ax - b\|_2 &= \|Q^*(APP^T x - b)\|_2 = \left\| \begin{bmatrix} R_{11} & R_{12} \\ 0 & 0 \end{bmatrix} \begin{bmatrix} y \\ z \end{bmatrix} - \begin{bmatrix} c \\ d \end{bmatrix} \right\|_2 \\ &= \sqrt{\|R_{11}y + R_{12}z - c\|_2^2 + \|d\|_2^2}. \end{aligned}$$

This shows that $\min \|Ax - b\|_2 = \|d\|_2 \iff R_{11}y = c - R_{12}z$.

QR method for rank deficient LSP

Theorem: Let $A \in \mathbb{C}^{m \times n}$. Suppose that $\text{rank}(A) = r$. Then there is a unitary matrix $Q \in \mathbb{C}^{m \times m}$ and a nonsingular upper triangular matrix $R_{11} \in \mathbb{C}^{r \times r}$ such that

$$AP = Q \begin{bmatrix} R_{11} & R_{12} \\ 0 & 0 \end{bmatrix} = QR,$$

where $P \in \mathbb{R}^{n \times n}$ is a permutation matrix and $R_{12} \in \mathbb{C}^{r \times (n-r)}$. ■

Set $\begin{bmatrix} c \\ d \end{bmatrix} := Q^*b$, where $c \in \mathbb{C}^r$ and $d \in \mathbb{C}^{m-r}$. Then

$$\begin{aligned} \|Ax - b\|_2 &= \|Q^*(APP^T x - b)\|_2 = \left\| \begin{bmatrix} R_{11} & R_{12} \\ 0 & 0 \end{bmatrix} \begin{bmatrix} y \\ z \end{bmatrix} - \begin{bmatrix} c \\ d \end{bmatrix} \right\|_2 \\ &= \sqrt{\|R_{11}y + R_{12}z - c\|_2^2 + \|d\|_2^2}. \end{aligned}$$

This shows that $\min \|Ax - b\|_2 = \|d\|_2 \iff R_{11}y = c - R_{12}z$. Hence $x = P \begin{bmatrix} y \\ z \end{bmatrix}$ is a solution of the LSP for any $z \in \mathbb{C}^{n-r}$ and $\|d\|_2$ is the residual. Setting $z = 0$ we obtain a unique solution with smallest norm.

QR method for rank deficient LSP

Algorithm: Solution of LSP $Ax \approx b$ when $\text{rank}(A) = r$.

1. Compute QR factorization $AP = Q \begin{bmatrix} R_{11} & R_{12} \\ 0 & 0 \end{bmatrix}$, where $Q \in \mathbb{C}^{m \times m}$ is unitary, $R_{11} \in \mathbb{C}^{r \times r}$ is nonsingular and upper triangular.
2. Set $\begin{bmatrix} c \\ d \end{bmatrix} := Q^* b$, where $c \in \mathbb{C}^r$ and $d \in \mathbb{C}^{m-r}$.

QR method for rank deficient LSP

Algorithm: Solution of LSP $Ax \approx b$ when $\text{rank}(A) = r$.

1. Compute QR factorization $AP = Q \begin{bmatrix} R_{11} & R_{12} \\ 0 & 0 \end{bmatrix}$, where $Q \in \mathbb{C}^{m \times m}$ is unitary, $R_{11} \in \mathbb{C}^{r \times r}$ is nonsingular and upper triangular.
2. Set $\begin{bmatrix} c \\ d \end{bmatrix} := Q^* b$, where $c \in \mathbb{C}^r$ and $d \in \mathbb{C}^{m-r}$.
3. Solve upper triangular system $R_{11}y = c$.
4. Set $x = P \begin{bmatrix} y \\ 0 \end{bmatrix}$. Then x is a unique solution of $Ax \approx b$.
5. Compute the residual $\|d\|_2$.

QR method for rank deficient LSP

Algorithm: Solution of LSP $Ax \approx b$ when $\text{rank}(A) = r$.

1. Compute QR factorization $AP = Q \begin{bmatrix} R_{11} & R_{12} \\ 0 & 0 \end{bmatrix}$, where $Q \in \mathbb{C}^{m \times m}$ is unitary, $R_{11} \in \mathbb{C}^{r \times r}$ is nonsingular and upper triangular.
2. Set $\begin{bmatrix} c \\ d \end{bmatrix} := Q^* b$, where $c \in \mathbb{C}^r$ and $d \in \mathbb{C}^{m-r}$.
3. Solve upper triangular system $R_{11}y = c$.
4. Set $x = P \begin{bmatrix} y \\ 0 \end{bmatrix}$. Then x is a unique solution of $Ax \approx b$.
5. Compute the residual $\|d\|_2$.

Remark: If x is a solution of LSP $Ax \approx b$ then $x + z$ is also a solution for any $z \in N(A)$. Hence the LSP has $n - r$ linearly independent solutions.

QR method for rank deficient LSP

Algorithm: Solution of LSP $Ax \approx b$ when $\text{rank}(A) = r$.

1. Compute QR factorization $AP = Q \begin{bmatrix} R_{11} & R_{12} \\ 0 & 0 \end{bmatrix}$, where $Q \in \mathbb{C}^{m \times m}$ is unitary, $R_{11} \in \mathbb{C}^{r \times r}$ is nonsingular and upper triangular.
2. Set $\begin{bmatrix} c \\ d \end{bmatrix} := Q^* b$, where $c \in \mathbb{C}^r$ and $d \in \mathbb{C}^{m-r}$.
3. Solve upper triangular system $R_{11}y = c$.
4. Set $x = P \begin{bmatrix} y \\ 0 \end{bmatrix}$. Then x is a unique solution of $Ax \approx b$.
5. Compute the residual $\|d\|_2$.

Remark: If x is a solution of LSP $Ax \approx b$ then $x + z$ is also a solution for any $z \in N(A)$. Hence the LSP has $n - r$ linearly independent solutions.

A rank deficient LSP is an ill-posed problem and solutions are strongly dependent on the rank of A . Numerical rank determination is a tricky problem.

QR method for rank deficient LSP

Algorithm: Solution of LSP $Ax \approx b$ when $\text{rank}(A) = r$.

1. Compute QR factorization $AP = Q \begin{bmatrix} R_{11} & R_{12} \\ 0 & 0 \end{bmatrix}$, where $Q \in \mathbb{C}^{m \times m}$ is unitary, $R_{11} \in \mathbb{C}^{r \times r}$ is nonsingular and upper triangular.
2. Set $\begin{bmatrix} c \\ d \end{bmatrix} := Q^* b$, where $c \in \mathbb{C}^r$ and $d \in \mathbb{C}^{m-r}$.
3. Solve upper triangular system $R_{11}y = c$.
4. Set $x = P \begin{bmatrix} y \\ 0 \end{bmatrix}$. Then x is a unique solution of $Ax \approx b$.
5. Compute the residual $\|d\|_2$.

Remark: If x is a solution of LSP $Ax \approx b$ then $x + z$ is also a solution for any $z \in N(A)$. Hence the LSP has $n - r$ linearly independent solutions.

A rank deficient LSP is an ill-posed problem and solutions are strongly dependent on the rank of A . Numerical rank determination is a tricky problem.

The MATLAB command `[Q,R,P] = qr(A)` computes a QR factorization $AP = QR$.

Uniqueness of QR factorization

Theorem: Let $A \in \mathbb{R}^{m \times n}$ and $\text{rank}(A) = n$. Then there exist unique isometry $Q \in \mathbb{R}^{m \times n}$ and an upper triangular matrix $R \in \mathbb{R}^{n \times n}$ with positive diagonal entries such that $A = QR$.

Uniqueness of QR factorization

Theorem: Let $A \in \mathbb{R}^{m \times n}$ and $\text{rank}(A) = n$. Then there exist unique isometry $Q \in \mathbb{R}^{m \times n}$ and an upper triangular matrix $R \in \mathbb{R}^{n \times n}$ with positive diagonal entries such that $A = QR$.

Proof: Let $A = \hat{Q}\hat{R}$ be a QR factorization of A , where $\hat{Q} \in \mathbb{R}^{m \times n}$ is an isometry and $\hat{R} \in \mathbb{R}^{n \times n}$ is upper triangular.

Uniqueness of QR factorization

Theorem: Let $A \in \mathbb{R}^{m \times n}$ and $\text{rank}(A) = n$. Then there exist unique isometry $Q \in \mathbb{R}^{m \times n}$ and an upper triangular matrix $R \in \mathbb{R}^{n \times n}$ with positive diagonal entries such that $A = QR$.

Proof: Let $A = \hat{Q}\hat{R}$ be a QR factorization of A , where $\hat{Q} \in \mathbb{R}^{m \times n}$ is an isometry and $\hat{R} \in \mathbb{R}^{n \times n}$ is upper triangular. Let r_{11}, \dots, r_{nn} be the diagonal entries of \hat{R} . Define $D := \text{diag}(\text{sign}(r_{11}), \dots, \text{sign}(r_{nn}))$.

Uniqueness of QR factorization

Theorem: Let $A \in \mathbb{R}^{m \times n}$ and $\text{rank}(A) = n$. Then there exist unique isometry $Q \in \mathbb{R}^{m \times n}$ and an upper triangular matrix $R \in \mathbb{R}^{n \times n}$ with positive diagonal entries such that $A = QR$.

Proof: Let $A = \hat{Q}\hat{R}$ be a QR factorization of A , where $\hat{Q} \in \mathbb{R}^{m \times n}$ is an isometry and $\hat{R} \in \mathbb{R}^{n \times n}$ is upper triangular. Let r_{11}, \dots, r_{nn} be the diagonal entries of \hat{R} . Define $D := \text{diag}(\text{sign}(r_{11}), \dots, \text{sign}(r_{nn}))$.

Then D is unitary, $Q := \hat{Q}D$ is an isometry, $R := D\hat{R}$ is upper triangular with positive diagonal entries, and $A = QR$.

Uniqueness of QR factorization

Theorem: Let $A \in \mathbb{R}^{m \times n}$ and $\text{rank}(A) = n$. Then there exist unique isometry $Q \in \mathbb{R}^{m \times n}$ and an upper triangular matrix $R \in \mathbb{R}^{n \times n}$ with positive diagonal entries such that $A = QR$.

Proof: Let $A = \hat{Q}\hat{R}$ be a QR factorization of A , where $\hat{Q} \in \mathbb{R}^{m \times n}$ is an isometry and $\hat{R} \in \mathbb{R}^{n \times n}$ is upper triangular. Let r_{11}, \dots, r_{nn} be the diagonal entries of \hat{R} . Define $D := \text{diag}(\text{sign}(r_{11}), \dots, \text{sign}(r_{nn}))$.

Then D is unitary, $Q := \hat{Q}D$ is an isometry, $R := D\hat{R}$ is upper triangular with positive diagonal entries, and $A = QR$. This proves the existence.

Uniqueness of QR factorization

Theorem: Let $A \in \mathbb{R}^{m \times n}$ and $\text{rank}(A) = n$. Then there exist unique isometry $Q \in \mathbb{R}^{m \times n}$ and an upper triangular matrix $R \in \mathbb{R}^{n \times n}$ with positive diagonal entries such that $A = QR$.

Proof: Let $A = \hat{Q}\hat{R}$ be a QR factorization of A , where $\hat{Q} \in \mathbb{R}^{m \times n}$ is an isometry and $\hat{R} \in \mathbb{R}^{n \times n}$ is upper triangular. Let r_{11}, \dots, r_{nn} be the diagonal entries of \hat{R} . Define $D := \text{diag}(\text{sign}(r_{11}), \dots, \text{sign}(r_{nn}))$.

Then D is unitary, $Q := \hat{Q}D$ is an isometry, $R := D\hat{R}$ is upper triangular with positive diagonal entries, and $A = QR$. This proves the existence.

Now suppose that $A = Q_1R_1 = Q_2R_2$, where Q_1, Q_2 are isometry and R_1, R_2 are upper triangular with positive diagonal entries.

Uniqueness of QR factorization

Theorem: Let $A \in \mathbb{R}^{m \times n}$ and $\text{rank}(A) = n$. Then there exist unique isometry $Q \in \mathbb{R}^{m \times n}$ and an upper triangular matrix $R \in \mathbb{R}^{n \times n}$ with positive diagonal entries such that $A = QR$.

Proof: Let $A = \hat{Q}\hat{R}$ be a QR factorization of A , where $\hat{Q} \in \mathbb{R}^{m \times n}$ is an isometry and $\hat{R} \in \mathbb{R}^{n \times n}$ is upper triangular. Let r_{11}, \dots, r_{nn} be the diagonal entries of \hat{R} . Define $D := \text{diag}(\text{sign}(r_{11}), \dots, \text{sign}(r_{nn}))$.

Then D is unitary, $Q := \hat{Q}D$ is an isometry, $R := D\hat{R}$ is upper triangular with positive diagonal entries, and $A = QR$. This proves the existence.

Now suppose that $A = Q_1R_1 = Q_2R_2$, where Q_1, Q_2 are isometry and R_1, R_2 are upper triangular with positive diagonal entries.

Then $A^\top A = R_1^\top R_1 = R_2^\top R_2$ are Cholesky factorizations of $A^\top A$. By uniqueness of Cholesky factorization, $R_1 = R_2$ which gives $Q_1 = Q_2$. ■
