

# Eigenvalue Problems

Rafikul Alam  
Department of Mathematics  
Indian Institute of Technology Guwahati  
Guwahati - 781039, INDIA

# Outline

- Eigenvalues and eigenvectors
- Multiplicity and diagonalizability
- Schur Theorem
- Eigenvector of triangular matrix
- Block Schur form
- Trace and eigenvalues

# Eigenvalue problem

Let  $A$  be an  $n \times n$  matrix. Then the **eigenvalue problem** for  $A$  is to solve

$$Av = \lambda v$$

for a **nonzero vector**  $v$  and a **scalar**  $\lambda$ . The scalar  $\lambda$  is called an **eigenvalue** of  $A$  and the vector  $v$  is called an **eigenvector** of  $A$  corresponding to  $\lambda$ .

# Eigenvalue problem

Let  $A$  be an  $n \times n$  matrix. Then the **eigenvalue problem** for  $A$  is to solve

$$Av = \lambda v$$

for a **nonzero vector**  $v$  and a **scalar**  $\lambda$ . The scalar  $\lambda$  is called an **eigenvalue** of  $A$  and the vector  $v$  is called an **eigenvector** of  $A$  corresponding to  $\lambda$ .

**Example:** Consider the matrix  $A = \begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix}$ . Then 4 is an eigenvalue of  $A$  with  $[1, 1]^T$  is a corresponding eigenvector.

# Eigenvalue problem

Let  $A$  be an  $n \times n$  matrix. Then the **eigenvalue problem** for  $A$  is to solve

$$Av = \lambda v$$

for a **nonzero vector**  $v$  and a **scalar**  $\lambda$ . The scalar  $\lambda$  is called an **eigenvalue** of  $A$  and the vector  $v$  is called an **eigenvector** of  $A$  corresponding to  $\lambda$ .

**Example:** Consider the matrix  $A = \begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix}$ . Then 4 is an eigenvalue of  $A$  with  $[1, 1]^T$  is a corresponding eigenvector.

Next, consider the rotation  $A = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$ . Then  $Av \neq \lambda v$  for any  $\lambda \in \mathbb{R}$  and  $0 \neq v \in \mathbb{R}^2$ . Hence we must consider eigenvalue problems over the field  $\mathbb{C}$ .

# Eigenvalue problem

Let  $A$  be an  $n \times n$  matrix. Then the **eigenvalue problem** for  $A$  is to solve

$$Av = \lambda v$$

for a **nonzero vector**  $v$  and a **scalar**  $\lambda$ . The scalar  $\lambda$  is called an **eigenvalue** of  $A$  and the vector  $v$  is called an **eigenvector** of  $A$  corresponding to  $\lambda$ .

**Example:** Consider the matrix  $A = \begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix}$ . Then 4 is an eigenvalue of  $A$  with  $[1, 1]^T$  is a corresponding eigenvector.

Next, consider the rotation  $A = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$ . Then  $Av \neq \lambda v$  for any  $\lambda \in \mathbb{R}$  and  $0 \neq v \in \mathbb{R}^2$ . Hence we must consider eigenvalue problems over the field  $\mathbb{C}$ .

Eigenvalue problems arise in many applications. For instance, **Google search engine** solves an eigenvalue problem for ranking web pages. Computation of eigenvalues and eigenvectors of a matrix is a major task.

# Eigenspace

**Definition:** Let  $\lambda$  be an eigenvalue of a matrix  $A \in \mathbb{C}^{n \times n}$ .

# Eigenspace

**Definition:** Let  $\lambda$  be an eigenvalue of a matrix  $A \in \mathbb{C}^{n \times n}$ . Then  $N(A - \lambda I) = \{\mathbf{x} : A\mathbf{x} = \lambda\mathbf{x}\}$  is called the **eigenspace of  $A$  corresponding to  $\lambda$** . Note that the eigenspace of  $A$  corresponding to  $\lambda$  consists of all eigenvectors of  $A$  corresponding to  $\lambda$  together with the **zero vector**.



# Eigenspace

**Definition:** Let  $\lambda$  be an eigenvalue of a matrix  $A \in \mathbb{C}^{n \times n}$ . Then  $N(A - \lambda I) = \{\mathbf{x} : A\mathbf{x} = \lambda\mathbf{x}\}$  is called the **eigenspace of  $A$  corresponding to  $\lambda$** . Note that the eigenspace of  $A$  corresponding to  $\lambda$  consists of all eigenvectors of  $A$  corresponding to  $\lambda$  together with the **zero vector**.

**Theorem:** Let  $A \in \mathbb{C}^{n \times n}$ . Then the following conditions are equivalent.

- $\lambda \in \mathbb{C}$  is an eigenvalue of  $A$ .

# Eigenspace

**Definition:** Let  $\lambda$  be an eigenvalue of a matrix  $A \in \mathbb{C}^{n \times n}$ . Then  $N(A - \lambda I) = \{\mathbf{x} : A\mathbf{x} = \lambda\mathbf{x}\}$  is called the **eigenspace of  $A$  corresponding to  $\lambda$** . Note that the eigenspace of  $A$  corresponding to  $\lambda$  consists of all eigenvectors of  $A$  corresponding to  $\lambda$  together with the **zero vector**.

**Theorem:** Let  $A \in \mathbb{C}^{n \times n}$ . Then the following conditions are equivalent.

- $\lambda \in \mathbb{C}$  is an eigenvalue of  $A$ .
- $N(A - \lambda I)$  is a **nontrivial subspace** of  $\mathbb{C}^n$ .

# Eigenspace

**Definition:** Let  $\lambda$  be an eigenvalue of a matrix  $A \in \mathbb{C}^{n \times n}$ . Then  $N(A - \lambda I) = \{\mathbf{x} : A\mathbf{x} = \lambda\mathbf{x}\}$  is called the **eigenspace of  $A$  corresponding to  $\lambda$** . Note that the eigenspace of  $A$  corresponding to  $\lambda$  consists of all eigenvectors of  $A$  corresponding to  $\lambda$  together with the **zero vector**.

**Theorem:** Let  $A \in \mathbb{C}^{n \times n}$ . Then the following conditions are equivalent.

- $\lambda \in \mathbb{C}$  is an eigenvalue of  $A$ .
- $N(A - \lambda I)$  is a **nontrivial subspace** of  $\mathbb{C}^n$ .
- $\text{rank}(A - \lambda I) < n$ .

# Eigenspace

**Definition:** Let  $\lambda$  be an eigenvalue of a matrix  $A \in \mathbb{C}^{n \times n}$ . Then  $N(A - \lambda I) = \{\mathbf{x} : A\mathbf{x} = \lambda\mathbf{x}\}$  is called the **eigenspace of  $A$  corresponding to  $\lambda$** . Note that the eigenspace of  $A$  corresponding to  $\lambda$  consists of all eigenvectors of  $A$  corresponding to  $\lambda$  together with the **zero vector**.

**Theorem:** Let  $A \in \mathbb{C}^{n \times n}$ . Then the following conditions are equivalent.

- $\lambda \in \mathbb{C}$  is an eigenvalue of  $A$ .
- $N(A - \lambda I)$  is a **nontrivial subspace** of  $\mathbb{C}^n$ .
- $\text{rank}(A - \lambda I) < n$ .
- $\det(A - \lambda I) = 0$ .

# Eigenspace

**Definition:** Let  $\lambda$  be an eigenvalue of a matrix  $A \in \mathbb{C}^{n \times n}$ . Then  $N(A - \lambda I) = \{\mathbf{x} : A\mathbf{x} = \lambda\mathbf{x}\}$  is called the **eigenspace of  $A$  corresponding to  $\lambda$** . Note that the eigenspace of  $A$  corresponding to  $\lambda$  consists of all eigenvectors of  $A$  corresponding to  $\lambda$  together with the **zero vector**.

**Theorem:** Let  $A \in \mathbb{C}^{n \times n}$ . Then the following conditions are equivalent.

- $\lambda \in \mathbb{C}$  is an eigenvalue of  $A$ .
- $N(A - \lambda I)$  is a **nontrivial subspace** of  $\mathbb{C}^n$ .
- $\text{rank}(A - \lambda I) < n$ .
- $\det(A - \lambda I) = 0$ .

**Definition:** Let  $A$  be an  $n \times n$  matrix. Then

- $p(\lambda) := \det(\lambda I - A)$  is called the **characteristic polynomial** of  $A$ .

# Eigenspace

**Definition:** Let  $\lambda$  be an eigenvalue of a matrix  $A \in \mathbb{C}^{n \times n}$ . Then  $N(A - \lambda I) = \{\mathbf{x} : A\mathbf{x} = \lambda\mathbf{x}\}$  is called the **eigenspace of  $A$  corresponding to  $\lambda$** . Note that the eigenspace of  $A$  corresponding to  $\lambda$  consists of all eigenvectors of  $A$  corresponding to  $\lambda$  together with the **zero vector**.

**Theorem:** Let  $A \in \mathbb{C}^{n \times n}$ . Then the following conditions are equivalent.

- $\lambda \in \mathbb{C}$  is an eigenvalue of  $A$ .
- $N(A - \lambda I)$  is a **nontrivial subspace** of  $\mathbb{C}^n$ .
- $\text{rank}(A - \lambda I) < n$ .
- $\det(A - \lambda I) = 0$ .

**Definition:** Let  $A$  be an  $n \times n$  matrix. Then

- $p(\lambda) := \det(\lambda I - A)$  is called the **characteristic polynomial** of  $A$ .
- $\det(\lambda I - A) = 0$  is called the **characteristic equation** of  $A$ .

# Eigenspace

**Definition:** Let  $\lambda$  be an eigenvalue of a matrix  $A \in \mathbb{C}^{n \times n}$ . Then  $N(A - \lambda I) = \{\mathbf{x} : A\mathbf{x} = \lambda\mathbf{x}\}$  is called the **eigenspace of  $A$  corresponding to  $\lambda$** . Note that the eigenspace of  $A$  corresponding to  $\lambda$  consists of all eigenvectors of  $A$  corresponding to  $\lambda$  together with the **zero vector**.

**Theorem:** Let  $A \in \mathbb{C}^{n \times n}$ . Then the following conditions are equivalent.

- $\lambda \in \mathbb{C}$  is an eigenvalue of  $A$ .
- $N(A - \lambda I)$  is a **nontrivial subspace** of  $\mathbb{C}^n$ .
- $\text{rank}(A - \lambda I) < n$ .
- $\det(A - \lambda I) = 0$ .

**Definition:** Let  $A$  be an  $n \times n$  matrix. Then

- $p(\lambda) := \det(\lambda I - A)$  is called the **characteristic polynomial** of  $A$ .
- $\det(\lambda I - A) = 0$  is called the **characteristic equation** of  $A$ .

**Definition:** The set  $\Lambda(A) := \{\lambda \in \mathbb{C} : \text{rank}(A - \lambda I) < n\}$  is called the **spectrum** of  $A$ .

# Eigenvalues

- The eigenvalues of  $A$  are the **roots** of the characteristic polynomial  $p(\lambda) := \det(\lambda I - A)$ . Hence  $A$  has at most  $n$  eigenvalues.



# Eigenvalues

- The eigenvalues of  $A$  are the **roots** of the characteristic polynomial  $p(\lambda) := \det(\lambda I - A)$ . Hence  $A$  has at most  $n$  eigenvalues.
- The eigenvalues of a **triangular** matrix are its **diagonal** entries.

# Eigenvalues

- The eigenvalues of  $A$  are the **roots** of the characteristic polynomial  $p(\lambda) := \det(\lambda I - A)$ . Hence  $A$  has at most  $n$  eigenvalues.
- The eigenvalues of a **triangular** matrix are its **diagonal** entries.
- $A$  is **invertible** iff  $0$  is not an eigenvalue of  $A$ .

# Eigenvalues

- The eigenvalues of  $A$  are the **roots** of the characteristic polynomial  $p(\lambda) := \det(\lambda I - A)$ . Hence  $A$  has at most  $n$  eigenvalues.
- The eigenvalues of a **triangular** matrix are its **diagonal** entries.
- $A$  is **invertible** iff  $0$  is not an eigenvalue of  $A$ .

**Example:** Let  $A := \begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix}$ . Then we have

$$\det(A - \lambda I) = \det \begin{bmatrix} 1 - \lambda & 3 \\ 3 & 1 - \lambda \end{bmatrix} = 0 \implies \lambda = 4, -2.$$

# Eigenvalues

- The eigenvalues of  $A$  are the **roots** of the characteristic polynomial  $p(\lambda) := \det(\lambda I - A)$ . Hence  $A$  has at most  $n$  eigenvalues.
- The eigenvalues of a **triangular** matrix are its **diagonal** entries.
- $A$  is **invertible** iff  $0$  is not an eigenvalue of  $A$ .

**Example:** Let  $A := \begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix}$ . Then we have

$$\det(A - \lambda I) = \det \begin{bmatrix} 1 - \lambda & 3 \\ 3 & 1 - \lambda \end{bmatrix} = 0 \implies \lambda = 4, -2.$$

$$N(A - 4I) = \left\{ \mathbf{x} : \begin{bmatrix} -3 & 3 \\ 3 & -3 \end{bmatrix} \mathbf{x} = \mathbf{0} \right\} = \text{span}([1, 1]^T),$$

# Eigenvalues

- The eigenvalues of  $A$  are the **roots** of the characteristic polynomial  $p(\lambda) := \det(\lambda I - A)$ . Hence  $A$  has at most  $n$  eigenvalues.
- The eigenvalues of a triangular matrix are its diagonal entries.
- $A$  is invertible iff 0 is not an eigenvalue of  $A$ .  $\Rightarrow \det(A - 0 \cdot I) = 0 \Rightarrow \det(A) = 0$ . So,  $A$  is singular, hence not invertible.

**Example:** Let  $A := \begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix}$ . Then we have

$$\det(A - \lambda I) = \det \begin{bmatrix} 1 - \lambda & 3 \\ 3 & 1 - \lambda \end{bmatrix} = 0 \implies \lambda = 4, -2.$$

$$N(A - 4I) = \left\{ \mathbf{x} : \begin{bmatrix} -3 & 3 \\ 3 & -3 \end{bmatrix} \mathbf{x} = \mathbf{0} \right\} = \text{span}([1, 1]^T),$$

$$N(A + 2I) = \left\{ \mathbf{x} : \begin{bmatrix} 3 & 3 \\ 3 & 3 \end{bmatrix} \mathbf{x} = \mathbf{0} \right\} = \text{span}([1, -1]^T).$$

# Algebraic and geometric multiplicities

**Definition:** Let  $\lambda$  be an eigenvalue of  $A \in \mathbb{C}^{n \times n}$ .

# Algebraic and geometric multiplicities

**Definition:** Let  $\lambda$  be an eigenvalue of  $A \in \mathbb{C}^{n \times n}$ .

- The **algebraic multiplicity** of  $\lambda$  is the multiplicity of  $\lambda$  as a root of the characteristic polynomial  $p(\lambda) := \det(\lambda I - A)$ .

# Algebraic and geometric multiplicities

**Definition:** Let  $\lambda$  be an eigenvalue of  $A \in \mathbb{C}^{n \times n}$ .

- The **algebraic multiplicity** of  $\lambda$  is the multiplicity of  $\lambda$  as a root of the characteristic polynomial  $p(\lambda) := \det(\lambda I - A)$ .
- The **geometric multiplicity** of  $\lambda$  is the dimension of  $N(A - \lambda I)$ .



# Algebraic and geometric multiplicities

**Definition:** Let  $\lambda$  be an eigenvalue of  $A \in \mathbb{C}^{n \times n}$ .

- The **algebraic multiplicity** of  $\lambda$  is the multiplicity of  $\lambda$  as a root of the characteristic polynomial  $p(\lambda) := \det(\lambda I - A)$ .
- The **geometric multiplicity** of  $\lambda$  is the dimension of  $N(A - \lambda I)$ .

**Theorem:** Let  $\lambda$  be an eigenvalue of  $A$ . Then

$$\text{geometric multiplicity of } \lambda \leq \text{algebraic multiplicity of } \lambda.$$

# Algebraic and geometric multiplicities

**Definition:** Let  $\lambda$  be an eigenvalue of  $A \in \mathbb{C}^{n \times n}$ .

- The **algebraic multiplicity** of  $\lambda$  is the multiplicity of  $\lambda$  as a root of the characteristic polynomial  $p(\lambda) := \det(\lambda I - A)$ .
- The **geometric multiplicity** of  $\lambda$  is the dimension of  $N(A - \lambda I)$ .

**Theorem:** Let  $\lambda$  be an eigenvalue of  $A$ . Then

$$\text{geometric multiplicity of } \lambda \leq \text{algebraic multiplicity of } \lambda.$$

**Example:** Let  $A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$ . Then 1 is the eigenvalue of  $A$  of **algebraic multiplicity** 3 and **geometric multiplicity** 1.

# Algebraic and geometric multiplicities

**Definition:** Let  $\lambda$  be an eigenvalue of  $A \in \mathbb{C}^{n \times n}$ .

- The **algebraic multiplicity** of  $\lambda$  is the multiplicity of  $\lambda$  as a root of the characteristic polynomial  $p(\lambda) := \det(\lambda I - A)$ .
- The **geometric multiplicity** of  $\lambda$  is the dimension of  $N(A - \lambda I)$ .

**Theorem:** Let  $\lambda$  be an eigenvalue of  $A$ . Then

$$\text{geometric multiplicity of } \lambda \leq \text{algebraic multiplicity of } \lambda.$$

**Example:** Let  $A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$ . Then 1 is the eigenvalue of  $A$  of **algebraic multiplicity** 3 and **geometric multiplicity** 1. Indeed, we have

$$N(A - I) = N\left(\begin{bmatrix} 0 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}\right) = \text{span}\left(\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}\right) \implies \dim(N(A - I)) = 1.$$

## Eigenvalues via characteristic polynomial

Let  $A \in \mathbb{C}^{n \times n}$  and  $p(x) := \det(xI - A)$  be the characteristic polynomial of  $A$ . Then  $\lambda \in \mathbb{C}$  is an eigenvalue of  $A \iff p(\lambda) = 0$ . Thus eigenvalues of  $A$  can be computed by finding the roots of  $p(x)$ . To see the efficacy of this method, consider  $A := \text{diag}(1, 2, \dots, 22)$ .

## Eigenvalues via characteristic polynomial

Let  $A \in \mathbb{C}^{n \times n}$  and  $p(x) := \det(xI - A)$  be the characteristic polynomial of  $A$ . Then  $\lambda \in \mathbb{C}$  is an eigenvalue of  $A \iff p(\lambda) = 0$ . Thus eigenvalues of  $A$  can be computed by finding the roots of  $p(x)$ . To see the efficacy of this method, consider  $A := \text{diag}(1, 2, \dots, 22)$ .

```
>> A = diag(1:22) ; % eigenvalues 1,2,...,22
>> p = charpoly(A); % coefficients of the poly p(x)
>> rt = roots(p); % roots of p(x)
>> rt(6:9) % displays four roots
```

## Eigenvalues via characteristic polynomial

Let  $A \in \mathbb{C}^{n \times n}$  and  $p(x) := \det(xI - A)$  be the characteristic polynomial of  $A$ . Then  $\lambda \in \mathbb{C}$  is an eigenvalue of  $A \iff p(\lambda) = 0$ . Thus eigenvalues of  $A$  can be computed by finding the roots of  $p(x)$ . To see the efficacy of this method, consider  $A := \text{diag}(1, 2, \dots, 22)$ .

```
>> A = diag(1:22) ; % eigenvalues 1,2,...,22
>> p = charpoly(A); % coefficients of the poly p(x)
>> rt = roots(p); % roots of p(x)
>> rt(6:9) % displays four roots
ans =
17.564435730130054 + 0.661474607910510i
17.564435730130054 - 0.661474607910510i
15.388471193084563 + 0.581348923952655i
15.388471193084563 - 0.581348923952655i
```

These roots do not have even a single correct digit! Never compute roots of  $\det(\lambda I - A)$  for computing eigenvalues of  $A$ .

# Similar matrices

The **diagonal** and **triangular** matrices are nice in the sense that their **eigenvalues** are displayed on the diagonals.

# Similar matrices

The **diagonal** and **triangular** matrices are nice in the sense that their **eigenvalues** are displayed on the diagonals.

It would be desirable to **transform** a square matrix to a **diagonal** or **triangular** matrix in such a way that they have **exactly the same eigenvalues**.



# Similar matrices

The **diagonal** and **triangular** matrices are nice in the sense that their **eigenvalues** are displayed on the diagonals.

It would be desirable to **transform** a square matrix to a **diagonal** or **triangular** matrix in such a way that they have **exactly the same eigenvalues**.

**Gaussian elimination** reduces a matrix to a triangular matrix. Unfortunately, this process **does not preserve the eigenvalues**.

# Similar matrices

The **diagonal** and **triangular** matrices are nice in the sense that their **eigenvalues** are displayed on the diagonals.

It would be desirable to **transform** a square matrix to a **diagonal** or **triangular** matrix in such a way that they have **exactly the same eigenvalues**.

**Gaussian elimination** reduces a matrix to a triangular matrix. Unfortunately, this process **does not preserve the eigenvalues**.

**Definition:** Let  $A, B \in \mathbb{C}^{n \times n}$ . Then  $A$  is said to be **similar** to  $B$  if there is an **invertible** matrix  $P \in \mathbb{C}^{n \times n}$  such that  $P^{-1}AP = B$ .

# Similar matrices

The **diagonal** and **triangular** matrices are nice in the sense that their **eigenvalues** are displayed on the diagonals.

It would be desirable to **transform** a square matrix to a **diagonal** or **triangular** matrix in such a way that they have **exactly the same eigenvalues**.

**Gaussian elimination** reduces a matrix to a triangular matrix. Unfortunately, this process **does not preserve the eigenvalues**.

**Definition:** Let  $A, B \in \mathbb{C}^{n \times n}$ . Then  $A$  is said to be **similar** to  $B$  if there is an **invertible** matrix  $P \in \mathbb{C}^{n \times n}$  such that  $P^{-1}AP = B$ .

The map  $A \mapsto P^{-1}AP$  is called a **similarity transformation** of  $A$ .

# Similar matrices

The **diagonal** and **triangular** matrices are nice in the sense that their **eigenvalues** are displayed on the diagonals.

It would be desirable to **transform** a square matrix to a **diagonal** or **triangular** matrix in such a way that they have **exactly the same eigenvalues**.

**Gaussian elimination** reduces a matrix to a triangular matrix. Unfortunately, this process **does not preserve the eigenvalues**.

**Definition:** Let  $A, B \in \mathbb{C}^{n \times n}$ . Then  $A$  is said to be **similar** to  $B$  if there is an **invertible** matrix  $P \in \mathbb{C}^{n \times n}$  such that  $P^{-1}AP = B$ .

The map  $A \mapsto P^{-1}AP$  is called a **similarity transformation** of  $A$ .

Note that  $\det(P^{-1}AP - \lambda I) = \det(A - \lambda I)$ . Consequently, similar matrices have the same **characteristic polynomials** and hence have the same eigenvalues.

PROOF:

<https://math.stackexchange.com/questions/8339/similar-matrices-have-the-same-eigenvalues-with-the-same-geometric-multiplicity>

# Diagonalization

Note that  $A$  and  $B := P^{-1}AP$  have the same eigenvalues. Indeed, we have  $\det(\lambda I - A) = \det(\lambda I - B)$ . Moreover,  $B\mathbf{v} = \lambda\mathbf{v} \iff A\mathbf{u} = \lambda\mathbf{u}$ , where  $\mathbf{u} := P\mathbf{v}$ .

# Diagonalization

Note that  $A$  and  $B := P^{-1}AP$  have the same eigenvalues. Indeed, we have  $\det(\lambda I - A) = \det(\lambda I - B)$ . Moreover,  $B\mathbf{v} = \lambda\mathbf{v} \iff A\mathbf{u} = \lambda\mathbf{u}$ , where  $\mathbf{u} := P\mathbf{v}$ .

**Example:** Let  $A = \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}$  and  $B = \begin{bmatrix} 1 & 0 \\ -2 & -1 \end{bmatrix}$ . Then  $AP = PB$ , where

$$P = \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}. \blacksquare$$

# Diagonalization

Note that  $A$  and  $B := P^{-1}AP$  have the same eigenvalues. Indeed, we have  $\det(\lambda I - A) = \det(\lambda I - B)$ . Moreover,  $B\mathbf{v} = \lambda\mathbf{v} \iff A\mathbf{u} = \lambda\mathbf{u}$ , where  $\mathbf{u} := P\mathbf{v}$ .

**Example:** Let  $A = \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}$  and  $B = \begin{bmatrix} 1 & 0 \\ -2 & -1 \end{bmatrix}$ . Then  $AP = PB$ , where

$$P = \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}. \blacksquare$$

**Definition:** An matrix  $A \in \mathbb{C}^{n \times n}$  is said to be **diagonalizable** if there is an **invertible** matrix  $P \in \mathbb{C}^{n \times n}$  such that  $D := P^{-1}AP$  is a **diagonal matrix**.

# Diagonalization

Note that  $A$  and  $B := P^{-1}AP$  have the same eigenvalues. Indeed, we have  $\det(\lambda I - A) = \det(\lambda I - B)$ . Moreover,  $B\mathbf{v} = \lambda\mathbf{v} \iff A\mathbf{u} = \lambda\mathbf{u}$ , where  $\mathbf{u} := P\mathbf{v}$ .

**Example:** Let  $A = \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}$  and  $B = \begin{bmatrix} 1 & 0 \\ -2 & -1 \end{bmatrix}$ . Then  $AP = PB$ , where  $P = \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$ . ■

**Definition:** An matrix  $A \in \mathbb{C}^{n \times n}$  is said to be **diagonalizable** if there is an **invertible** matrix  $P \in \mathbb{C}^{n \times n}$  such that  $D := P^{-1}AP$  is a **diagonal matrix**.

**Theorem:** Let  $A \in \mathbb{C}^{n \times n}$ . Then  $A$  is diagonalizable  $\iff A$  has  $n$  **linearly independent eigenvectors**.



# Diagonalization

Note that  $A$  and  $B := P^{-1}AP$  have the same eigenvalues. Indeed, we have  $\det(\lambda I - A) = \det(\lambda I - B)$ . Moreover,  $B\mathbf{v} = \lambda\mathbf{v} \iff A\mathbf{u} = \lambda\mathbf{u}$ , where  $\mathbf{u} := P\mathbf{v}$ .

**Example:** Let  $A = \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}$  and  $B = \begin{bmatrix} 1 & 0 \\ -2 & -1 \end{bmatrix}$ . Then  $AP = PB$ , where  $P = \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$ . ■

**Definition:** An matrix  $A \in \mathbb{C}^{n \times n}$  is said to be **diagonalizable** if there is an **invertible** matrix  $P \in \mathbb{C}^{n \times n}$  such that  $D := P^{-1}AP$  is a **diagonal matrix**.

**Theorem:** Let  $A \in \mathbb{C}^{n \times n}$ . Then  $A$  is diagonalizable  $\iff A$  has  **$n$  linearly independent eigenvectors**. In particular, if  $A$  has  **$n$  distinct eigenvalues** then  $A$  is diagonalizable.

**Proof:** If  $P^{-1}AP = \text{diag}(\lambda_1, \dots, \lambda_m)$  then  $A\mathbf{v}_j = \lambda_j\mathbf{v}_j$ , where  $\mathbf{v}_j := Pe_j$ .

# Diagonalization

Note that  $A$  and  $B := P^{-1}AP$  have the same eigenvalues. Indeed, we have  $\det(\lambda I - A) = \det(\lambda I - B)$ . Moreover,  $B\mathbf{v} = \lambda\mathbf{v} \iff A\mathbf{u} = \lambda\mathbf{u}$ , where  $\mathbf{u} := P\mathbf{v}$ .

**Example:** Let  $A = \begin{bmatrix} 1 & 2 \\ 0 & -1 \end{bmatrix}$  and  $B = \begin{bmatrix} 1 & 0 \\ -2 & -1 \end{bmatrix}$ . Then  $AP = PB$ , where  $P = \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$ . ■

**Definition:** An matrix  $A \in \mathbb{C}^{n \times n}$  is said to be **diagonalizable** if there is an **invertible** matrix  $P \in \mathbb{C}^{n \times n}$  such that  $D := P^{-1}AP$  is a **diagonal matrix**.

**Theorem:** Let  $A \in \mathbb{C}^{n \times n}$ . Then  $A$  is diagonalizable  $\iff A$  has  **$n$  linearly independent eigenvectors**. In particular, if  $A$  has  **$n$  distinct eigenvalues** then  $A$  is diagonalizable.

**Proof:** If  $P^{-1}AP = \text{diag}(\lambda_1, \dots, \lambda_m)$  then  $A\mathbf{v}_j = \lambda_j\mathbf{v}_j$ , where  $\mathbf{v}_j := Pe_j$ . Conversely, if  $A\mathbf{v}_i = \lambda_i\mathbf{v}_i$  for  $i = 1 : n$  and  $\mathbf{v}_1, \dots, \mathbf{v}_n$  are linearly independent then  $\hookrightarrow$  As  $u_j = e_j$

$$A[\mathbf{v}_1, \dots, \mathbf{v}_n] = [\mathbf{v}_1, \dots, \mathbf{v}_n]\text{diag}(\lambda_1, \dots, \lambda_n). \blacksquare$$

# Schur triangularization theorem

**Schur Theorem:** Let  $A \in \mathbb{C}^{n \times n}$ . Then there is a unitary matrix  $U \in \mathbb{C}^{n \times n}$  such that  $U^*AU$  is upper triangular.

# Schur triangularization theorem

**Schur Theorem:** Let  $A \in \mathbb{C}^{n \times n}$ . Then there is a unitary matrix  $U \in \mathbb{C}^{n \times n}$  such that  $U^*AU$  is upper triangular.

**Proof:** Use induction on  $n$ . The result holds for  $n = 1$ . Assume that the result holds for  $n - 1$ .

# Schur triangularization theorem

**Schur Theorem:** Let  $A \in \mathbb{C}^{n \times n}$ . Then there is a unitary matrix  $U \in \mathbb{C}^{n \times n}$  such that  $U^*AU$  is upper triangular.

**Proof:** Use induction on  $n$ . The result holds for  $n = 1$ . Assume that the result holds for  $n - 1$ .

Let  $A\mathbf{u}_1 = \lambda_1\mathbf{u}_1$  be such that  $\|\mathbf{u}_1\|_2 = 1$ . Choose  $\mathbf{u}_2, \dots, \mathbf{u}_n$  such that  $U_1 := [\mathbf{u}_1 \ \mathbf{u}_2 \ \cdots \ \mathbf{u}_n]$  is unitary.

# Schur triangularization theorem

**Schur Theorem:** Let  $A \in \mathbb{C}^{n \times n}$ . Then there is a unitary matrix  $U \in \mathbb{C}^{n \times n}$  such that  $U^*AU$  is upper triangular.

**Proof:** Use induction on  $n$ . The result holds for  $n = 1$ . Assume that the result holds for  $n - 1$ .

Let  $A\mathbf{u}_1 = \lambda_1\mathbf{u}_1$  be such that  $\|\mathbf{u}_1\|_2 = 1$ . Choose  $\mathbf{u}_2, \dots, \mathbf{u}_n$  such that

$U_1 := [\mathbf{u}_1 \ \mathbf{u}_2 \ \cdots \ \mathbf{u}_n]$  is unitary. Then  $AU_1 = [\lambda_1\mathbf{u}_1 \ A\mathbf{u}_2 \ \cdots \ A\mathbf{u}_n]$  and hence

$$U_1^*AU_1 = \begin{bmatrix} \lambda_1 & \mathbf{h}^\top \\ 0 & H \end{bmatrix} \text{ for some } \mathbf{h} \in \mathbb{C}^{n-1} \text{ and } H \in \mathbb{C}^{(n-1) \times (n-1)}.$$

# Schur triangularization theorem

**Schur Theorem:** Let  $A \in \mathbb{C}^{n \times n}$ . Then there is a unitary matrix  $U \in \mathbb{C}^{n \times n}$  such that  $U^*AU$  is upper triangular.

**Proof:** Use induction on  $n$ . The result holds for  $n = 1$ . Assume that the result holds for  $n - 1$ .

Let  $A\mathbf{u}_1 = \lambda_1\mathbf{u}_1$  be such that  $\|\mathbf{u}_1\|_2 = 1$ . Choose  $\mathbf{u}_2, \dots, \mathbf{u}_n$  such that

$U_1 := [\mathbf{u}_1 \ \mathbf{u}_2 \ \cdots \ \mathbf{u}_n]$  is unitary. Then  $AU_1 = [\lambda_1\mathbf{u}_1 \ A\mathbf{u}_2 \ \cdots \ A\mathbf{u}_n]$  and hence

$$U_1^*AU_1 = \begin{bmatrix} \lambda_1 & \mathbf{h}^\top \\ 0 & H \end{bmatrix} \text{ for some } \mathbf{h} \in \mathbb{C}^{n-1} \text{ and } H \in \mathbb{C}^{(n-1) \times (n-1)}.$$

By induction hypothesis, there exists a unitary  $V$  such that  $V^*HV = T$  is upper triangular.

# Schur triangularization theorem

**Schur Theorem:** Let  $A \in \mathbb{C}^{n \times n}$ . Then there is a unitary matrix  $U \in \mathbb{C}^{n \times n}$  such that  $U^*AU$  is upper triangular.

**Proof:** Use induction on  $n$ . The result holds for  $n = 1$ . Assume that the result holds for  $n - 1$ .

Let  $A\mathbf{u}_1 = \lambda_1\mathbf{u}_1$  be such that  $\|\mathbf{u}_1\|_2 = 1$ . Choose  $\mathbf{u}_2, \dots, \mathbf{u}_n$  such that

$U_1 := [\mathbf{u}_1 \ \mathbf{u}_2 \ \cdots \ \mathbf{u}_n]$  is unitary. Then  $AU_1 = [\lambda_1\mathbf{u}_1 \ A\mathbf{u}_2 \ \cdots \ A\mathbf{u}_n]$  and hence

$$U_1^*AU_1 = \begin{bmatrix} \lambda_1 & \mathbf{h}^\top \\ 0 & H \end{bmatrix} \text{ for some } \mathbf{h} \in \mathbb{C}^{n-1} \text{ and } H \in \mathbb{C}^{(n-1) \times (n-1)}.$$

By induction hypothesis, there exists a unitary  $V$  such that  $V^*HV = T$  is upper triangular.

Set  $U := U_1 \begin{bmatrix} 1 & 0 \\ 0 & V \end{bmatrix}$ .



# Schur triangularization theorem

**Schur Theorem:** Let  $A \in \mathbb{C}^{n \times n}$ . Then there is a unitary matrix  $U \in \mathbb{C}^{n \times n}$  such that  $U^*AU$  is upper triangular.

**Proof:** Use induction on  $n$ . The result holds for  $n = 1$ . Assume that the result holds for  $n - 1$ .

Let  $A\mathbf{u}_1 = \lambda_1\mathbf{u}_1$  be such that  $\|\mathbf{u}_1\|_2 = 1$ . Choose  $\mathbf{u}_2, \dots, \mathbf{u}_n$  such that

$U_1 := [\mathbf{u}_1 \ \mathbf{u}_2 \ \cdots \ \mathbf{u}_n]$  is unitary. Then  $AU_1 = [\lambda_1\mathbf{u}_1 \ A\mathbf{u}_2 \ \cdots \ A\mathbf{u}_n]$  and hence

$$U_1^*AU_1 = \begin{bmatrix} \lambda_1 & \mathbf{h}^\top \\ 0 & H \end{bmatrix} \text{ for some } \mathbf{h} \in \mathbb{C}^{n-1} \text{ and } H \in \mathbb{C}^{(n-1) \times (n-1)}.$$

By induction hypothesis, there exists a unitary  $V$  such that  $V^*HV = T$  is upper triangular.

Set  $U := U_1 \begin{bmatrix} 1 & 0 \\ 0 & V \end{bmatrix}$ . Then

$$U^*AU = \begin{bmatrix} \lambda_1 & \mathbf{h}^\top V \\ 0 & T \end{bmatrix} = \text{upper triangular.} \blacksquare$$

# Spectral theorem for Hermitian matrices

**Theorem:** Let  $A \in \mathbb{C}^{n \times n}$ . Then  $A$  is Hermitian  $\iff$  there exist a unitary matrix  $U$  and a real diagonal matrix  $D$  such that

$$U^*AU = D.$$

# Spectral theorem for Hermitian matrices

**Theorem:** Let  $A \in \mathbb{C}^{n \times n}$ . Then  $A$  is Hermitian  $\iff$  there exist a unitary matrix  $U$  and a real diagonal matrix  $D$  such that

$$U^*AU = D.$$

Thus a Hermitian matrix is unitarily diagonalizable and has  $n$  orthonormal eigenvectors.

# Spectral theorem for Hermitian matrices

**Theorem:** Let  $A \in \mathbb{C}^{n \times n}$ . Then  $A$  is Hermitian  $\iff$  there exist a unitary matrix  $U$  and a real diagonal matrix  $D$  such that

$$U^*AU = D.$$

Thus a Hermitian matrix is unitarily diagonalizable and has  $n$  orthonormal eigenvectors.

**Proof:** By Schur theorem, there is a unitary matrix  $U$  and an upper triangular matrix  $D$  such that  $U^*AU = D$ .

# Spectral theorem for Hermitian matrices

**Theorem:** Let  $A \in \mathbb{C}^{n \times n}$ . Then  $A$  is Hermitian  $\iff$  there exist a unitary matrix  $U$  and a real diagonal matrix  $D$  such that

$$U^*AU = D.$$

Thus a Hermitian matrix is unitarily diagonalizable and has  $n$  orthonormal eigenvectors.

**Proof:** By Schur theorem, there is a unitary matrix  $U$  and an upper triangular matrix  $D$  such that  $U^*AU = D$ .

Now  $A^* = A \iff D^* = D \iff D$  is a diagonal matrix with real diagonals.

# Spectral theorem for Hermitian matrices

**Theorem:** Let  $A \in \mathbb{C}^{n \times n}$ . Then  $A$  is Hermitian  $\iff$  there exist a unitary matrix  $U$  and a real diagonal matrix  $D$  such that

$$U^*AU = D.$$

Thus a Hermitian matrix is unitarily diagonalizable and has  $n$  orthonormal eigenvectors.

**Proof:** By Schur theorem, there is a unitary matrix  $U$  and an upper triangular matrix  $D$  such that  $U^*AU = D$ .

Now  $A^* = A \iff D^* = D \iff D$  is a diagonal matrix with real diagonals. If  $D = \text{diag}(\lambda_1, \dots, \lambda_n)$  and  $U = [\mathbf{u}_1, \dots, \mathbf{u}_n]$  then

# Spectral theorem for Hermitian matrices

**Theorem:** Let  $A \in \mathbb{C}^{n \times n}$ . Then  $A$  is Hermitian  $\iff$  there exist a unitary matrix  $U$  and a real diagonal matrix  $D$  such that

$$U^*AU = D.$$

Thus a Hermitian matrix is unitarily diagonalizable and has  $n$  orthonormal eigenvectors.

**Proof:** By Schur theorem, there is a unitary matrix  $U$  and an upper triangular matrix  $D$  such that  $U^*AU = D$ .

Now  $A^* = A \iff D^* = D \iff D$  is a diagonal matrix with real diagonals. If  $D = \text{diag}(\lambda_1, \dots, \lambda_n)$  and  $U = [\mathbf{u}_1, \dots, \mathbf{u}_n]$  then

$$A\mathbf{u}_j = \lambda_j\mathbf{u}_j \text{ for } j = 1 : n$$

# Spectral theorem for Hermitian matrices

**Theorem:** Let  $A \in \mathbb{C}^{n \times n}$ . Then  $A$  is Hermitian  $\iff$  there exist a unitary matrix  $U$  and a real diagonal matrix  $D$  such that

$$U^*AU = D.$$

★ Thus a Hermitian matrix is unitarily diagonalizable and has  $n$  orthonormal eigenvectors.

**Proof:** By Schur theorem, there is a unitary matrix  $U$  and an upper triangular matrix  $D$  such that  $U^*AU = D$ .

Now  $A^* = A \iff D^* = D \iff D$  is a diagonal matrix with real diagonals. If  $D = \text{diag}(\lambda_1, \dots, \lambda_n)$  and  $U = [\mathbf{u}_1, \dots, \mathbf{u}_n]$  then

$$A\mathbf{u}_j = \lambda_j\mathbf{u}_j \text{ for } j = 1 : n$$

and  $\mathbf{u}_1, \dots, \mathbf{u}_n$  are orthonormal. ■



# Eigenvectors of triangular matrix

By Schur's theorem, an  $n \times n$  matrix can be transformed into triangular (Schur) form by unitary similarity and eigenvalues of a triangular matrix are the diagonal entries.

# Eigenvectors of triangular matrix

By Schur's theorem, an  $n \times n$  matrix can be transformed into triangular (Schur) form by unitary similarity and eigenvalues of a triangular matrix are the diagonal entries.

Eigenvectors of a triangular matrix are less obvious, but still easy to compute. Suppose that  $A$  is upper triangular and  $\lambda$  is a simple eigenvalue of  $A$ .

# Eigenvectors of triangular matrix

By Schur's theorem, an  $n \times n$  matrix can be transformed into triangular (Schur) form by unitary similarity and eigenvalues of a triangular matrix are the diagonal entries.

Eigenvectors of a triangular matrix are less obvious, but still easy to compute. Suppose that  $A$  is upper triangular and  $\lambda$  is a simple eigenvalue of  $A$ . Then  $A - \lambda I$  will be of the form

$$A - \lambda I = \begin{bmatrix} A_{11} & \mathbf{b} & A_{13} \\ & 0 & \mathbf{h}^\top \\ & & A_{33} \end{bmatrix}.$$

## Eigenvectors of triangular matrix

By Schur's theorem, an  $n \times n$  matrix can be transformed into triangular (Schur) form by unitary similarity and eigenvalues of a triangular matrix are the diagonal entries.

Eigenvectors of a triangular matrix are less obvious, but still easy to compute. Suppose that  $A$  is upper triangular and  $\lambda$  is a simple eigenvalue of  $A$ . Then  $A - \lambda I$  will be of the form

$$A - \lambda I = \begin{bmatrix} A_{11} & \mathbf{b} & A_{13} \\ & 0 & \mathbf{h}^\top \\ & & A_{33} \end{bmatrix}.$$

Then solving the upper triangular system  $A_{11}\mathbf{x} = \mathbf{b}$  and defining

$$\mathbf{v} := \begin{bmatrix} \mathbf{x} \\ -1 \\ \mathbf{0} \end{bmatrix}$$

it follows that  $\mathbf{v}$  is an eigenvector of  $A$  corresponding to  $\lambda$ , that is,  $A\mathbf{v} = \lambda\mathbf{v}$ .

# Block upper triangular form

**Real Schur Theorem:** Let  $A \in \mathbb{R}^{n \times n}$ . Then there is an orthogonal matrix  $U \in \mathbb{R}^{n \times n}$  such that  $U^*AU$  is **block upper triangular** given by

# Block upper triangular form

**Real Schur Theorem:** Let  $A \in \mathbb{R}^{n \times n}$ . Then there is an orthogonal matrix  $U \in \mathbb{R}^{n \times n}$  such that  $U^*AU$  is **block upper triangular** given by

$$U^*AU = \begin{bmatrix} T_{11} & T_{12} & \cdots & T_{1m} \\ & T_{22} & \cdots & T_{2m} \\ & & \ddots & \vdots \\ & & & T_{mm} \end{bmatrix},$$

where  $T_{jj} = \lambda_j \in \mathbb{R}$  or  $T_{jj} = \begin{bmatrix} \alpha_j & -\beta_j \\ \beta_j & \alpha_j \end{bmatrix}$  with  $\alpha_j, \beta_j \in \mathbb{R}$  for  $j = 1 : m$ .

# Block upper triangular form

**Real Schur Theorem:** Let  $A \in \mathbb{R}^{n \times n}$ . Then there is an orthogonal matrix  $U \in \mathbb{R}^{n \times n}$  such that  $U^*AU$  is **block upper triangular** given by

$$U^*AU = \begin{bmatrix} T_{11} & T_{12} & \cdots & T_{1m} \\ & T_{22} & \cdots & T_{2m} \\ & & \ddots & \vdots \\ & & & T_{mm} \end{bmatrix},$$

where  $T_{jj} = \lambda_j \in \mathbb{R}$  or  $T_{jj} = \begin{bmatrix} \alpha_j & -\beta_j \\ \beta_j & \alpha_j \end{bmatrix}$  with  $\alpha_j, \beta_j \in \mathbb{R}$  for  $j = 1 : m$ .

Further, each  $2 \times 2$  block  $T_{jj} = \begin{bmatrix} \alpha_j & -\beta_j \\ \beta_j & \alpha_j \end{bmatrix}$  corresponds to a **pair of complex conjugate eigenvalues**  $\alpha_j \pm i\beta_j$  of  $A$ .

# Trace and eigenvalues

**Theorem:** Let  $A \in \mathbb{C}^{n \times n}$  with eigenvalues  $\lambda_1, \dots, \lambda_n$  (counted according to their algebraic multiplicities).



# Trace and eigenvalues

**Theorem:** Let  $A \in \mathbb{C}^{n \times n}$  with eigenvalues  $\lambda_1, \dots, \lambda_n$  (counted according to their algebraic multiplicities). Then

$$\text{Trace}(A) = \lambda_1 + \dots + \lambda_n \text{ and } \det(A) = \prod_{j=1}^n \lambda_j.$$

**Proof:** By Schur theorem, there is a unitary matrix  $U$  such that  $T := U^*AU$  is upper triangular.

# Trace and eigenvalues

**Theorem:** Let  $A \in \mathbb{C}^{n \times n}$  with eigenvalues  $\lambda_1, \dots, \lambda_n$  (counted according to their algebraic multiplicities). Then

$$\text{Trace}(A) = \lambda_1 + \dots + \lambda_n \text{ and } \det(A) = \prod_{j=1}^n \lambda_j.$$

**Proof:** By Schur theorem, there is a unitary matrix  $U$  such that  $T := U^*AU$  is upper triangular. Hence

$$\det(A) = \det(T) = \prod_{j=1}^n \lambda_j.$$

# Trace and eigenvalues

**Theorem:** Let  $A \in \mathbb{C}^{n \times n}$  with eigenvalues  $\lambda_1, \dots, \lambda_n$  (counted according to their algebraic multiplicities). Then

$$\text{Trace}(A) = \lambda_1 + \dots + \lambda_n \text{ and } \det(A) = \prod_{j=1}^n \lambda_j.$$

**Proof:** By Schur theorem, there is a unitary matrix  $U$  such that  $T := U^*AU$  is upper triangular. Hence

$$\det(A) = \det(T) = \prod_{j=1}^n \lambda_j.$$

Now  $\text{Trace}(A) = \text{Trace}(UTU^*) = \text{Trace}(T) = \lambda_1 + \dots + \lambda_n$ . ■

**Exercise:** Let  $A, B \in \mathbb{C}^{n \times n}$ . Show that  $\text{Trace}(AB) = \text{Trace}(BA)$ .

