

Implicit QR Algorithm for Eigenvalue Problems

Rafikul Alam
Department of Mathematics
Indian Institute of Technology Guwahati
Guwahati - 781039, INDIA

Outline

- Implicit QR algorithm and bulge chasing
- Implicit single and double shift QR algorithm

Explicit QR step

Let A be proper Hessenberg. Then a **shifted explicit QR step with shift μ** is given by

- $A - \mu I = Q_1 Q_2 \cdots Q_{n-1} R$ % Gives QR factorization of $A - \mu I$.

Explicit QR step

Let A be proper Hessenberg. Then a **shifted explicit QR step with shift μ** is given by

- $A - \mu I = Q_1 Q_2 \cdots Q_{n-1} R$ % Gives QR factorization of $A - \mu I$.
- $A_1 = R Q_1 Q_2 \cdots Q_{n-1} + \mu I$

Explicit QR step

Let A be proper Hessenberg. Then a **shifted explicit QR step with shift μ** is given by

- $A - \mu I = Q_1 Q_2 \cdots Q_{n-1} R$ % Gives QR factorization of $A - \mu I$.
- $A_1 = R Q_1 Q_2 \cdots Q_{n-1} + \mu I = (Q_1 Q_2 \cdots Q_{n-1})^* A Q_1 Q_2 \cdots Q_{n-1}$.

Explicit QR step

Let A be proper Hessenberg. Then a **shifted explicit QR step with shift μ** is given by

- $A - \mu I = Q_1 Q_2 \cdots Q_{n-1} R$ % Gives QR factorization of $A - \mu I$.
- $A_1 = R Q_1 Q_2 \cdots Q_{n-1} + \mu I = (Q_1 Q_2 \cdots Q_{n-1})^* A Q_1 Q_2 \cdots Q_{n-1}$.

Note that A_1 is Hessenberg. Let us take a close look at A_1 :

$$A_1 = (Q_1 Q_2 \cdots Q_{n-1})^* A Q_1 Q_2 \cdots Q_{n-1}$$

Explicit QR step

Let A be proper Hessenberg. Then a **shifted explicit QR step with shift μ** is given by

- $A - \mu I = Q_1 Q_2 \cdots Q_{n-1} R$ % Gives QR factorization of $A - \mu I$.
- $A_1 = R Q_1 Q_2 \cdots Q_{n-1} + \mu I = (Q_1 Q_2 \cdots Q_{n-1})^* A Q_1 Q_2 \cdots Q_{n-1}$.

Note that A_1 is Hessenberg. Let us take a close look at A_1 :

$$A_1 = (Q_1 Q_2 \cdots Q_{n-1})^* A Q_1 Q_2 \cdots Q_{n-1} = Q_{n-1}^* \cdots Q_2^* (Q_1^* A Q_1) Q_2 \cdots Q_{n-1}.$$

Explicit QR step

Let A be proper Hessenberg. Then a **shifted explicit QR step with shift μ** is given by

- $A - \mu I = Q_1 Q_2 \cdots Q_{n-1} R$ % Gives QR factorization of $A - \mu I$.
- $A_1 = R Q_1 Q_2 \cdots Q_{n-1} + \mu I = (Q_1 Q_2 \cdots Q_{n-1})^* A Q_1 Q_2 \cdots Q_{n-1}$.

Note that A_1 is Hessenberg. Let us take a close look at A_1 :

$$A_1 = (Q_1 Q_2 \cdots Q_{n-1})^* A Q_1 Q_2 \cdots Q_{n-1} = Q_{n-1}^* \cdots Q_2^* (Q_1^* A Q_1) Q_2 \cdots Q_{n-1}.$$

Next, observe that the similarity transformation $A \mapsto Q_1^* A Q_1$ destroys the Hessenberg structure:

Explicit QR step

Let A be proper Hessenberg. Then a **shifted explicit QR step with shift μ** is given by

- $A - \mu I = Q_1 Q_2 \cdots Q_{n-1} R$ % Gives QR factorization of $A - \mu I$.
- $A_1 = R Q_1 Q_2 \cdots Q_{n-1} + \mu I = (Q_1 Q_2 \cdots Q_{n-1})^* A Q_1 Q_2 \cdots Q_{n-1}$.

Note that A_1 is Hessenberg. Let us take a close look at A_1 :

$$A_1 = (Q_1 Q_2 \cdots Q_{n-1})^* A Q_1 Q_2 \cdots Q_{n-1} = Q_{n-1}^* \cdots Q_2^* (Q_1^* A Q_1) Q_2 \cdots Q_{n-1}.$$

Next, observe that the similarity transformation $A \mapsto Q_1^* A Q_1$ destroys the Hessenberg structure:

$$Q_1^* \begin{bmatrix} \times & \times & \times & \times & \times \\ \times & \times & \times & \times & \times \\ & \times & \times & \times & \times \\ & & \times & \times & \times \\ & & & \times & \times \end{bmatrix} Q_1 = \begin{bmatrix} * & * & * & * & * \\ & * & * & * & * \\ & \times & \times & \times & \times \\ & & \times & \times & \times \\ & & & \times & \times \end{bmatrix} Q_1$$

Explicit QR step

Let A be proper Hessenberg. Then a **shifted explicit QR step with shift μ** is given by

- $A - \mu I = Q_1 Q_2 \cdots Q_{n-1} R$ % Gives QR factorization of $A - \mu I$.
- $A_1 = R Q_1 Q_2 \cdots Q_{n-1} + \mu I = (Q_1 Q_2 \cdots Q_{n-1})^* A Q_1 Q_2 \cdots Q_{n-1}$.

Note that A_1 is Hessenberg. Let us take a close look at A_1 :

$$A_1 = (Q_1 Q_2 \cdots Q_{n-1})^* A Q_1 Q_2 \cdots Q_{n-1} = Q_{n-1}^* \cdots Q_2^* (Q_1^* A Q_1) Q_2 \cdots Q_{n-1}.$$

Next, observe that the similarity transformation $A \mapsto Q_1^* A Q_1$ destroys the Hessenberg structure:

$$Q_1^* \begin{bmatrix} \times & \times & \times & \times & \times \\ \times & \times & \times & \times & \times \\ & \times & \times & \times & \times \\ & & \times & \times & \times \\ & & & \times & \times \end{bmatrix} Q_1 = \begin{bmatrix} * & * & * & * & * \\ & * & * & * & * \\ & \times & \times & \times & \times \\ & & \times & \times & \times \\ & & & \times & \times \end{bmatrix} Q_1 = \begin{bmatrix} * & * & \times & \times & \times \\ * & * & \times & \times & \times \\ * & * & \times & \times & \times \\ & & \times & \times & \times \\ & & & \times & \times \end{bmatrix}$$

Bulge chasing implicit QR step

Since $A_1 = Q_{n-1}^* \cdots Q_2^* (Q_1^* A Q_1) Q_2 \cdots Q_{n-1}$ is Hessenberg but $Q_1^* A Q_1$ is NOT Hessenberg,

Bulge chasing implicit QR step

Since $A_1 = Q_{n-1}^* \cdots Q_2^* (Q_1^* A Q_1) Q_2 \cdots Q_{n-1}$ is Hessenberg but $Q_1^* A Q_1$ is NOT Hessenberg, it follows that the rotations Q_2, \dots, Q_{n-1} reduces $Q_1^* A Q_1$ to Hessenberg form.

Bulge chasing implicit QR step

Since $A_1 = Q_{n-1}^* \cdots Q_2^* (Q_1^* A Q_1) Q_2 \cdots Q_{n-1}$ is Hessenberg but $Q_1^* A Q_1$ is NOT Hessenberg, it follows that the rotations Q_2, \dots, Q_{n-1} reduces $Q_1^* A Q_1$ to Hessenberg form. However, we can independently reduce $Q_1^* A Q_1$ to Hessenberg form without using the rotations Q_2, \dots, Q_{n-1} .

Bulge chasing implicit QR step

Since $A_1 = Q_{n-1}^* \cdots Q_2^* (Q_1^* A Q_1) Q_2 \cdots Q_{n-1}$ is Hessenberg but $Q_1^* A Q_1$ is NOT Hessenberg, it follows that the rotations Q_2, \dots, Q_{n-1} reduces $Q_1^* A Q_1$ to Hessenberg form. However, we can independently reduce $Q_1^* A Q_1$ to Hessenberg form without using the rotations Q_2, \dots, Q_{n-1} .

$$\underbrace{\begin{bmatrix} \times & \times & \times & \times & \times \\ \times & \times & \times & \times & \times \\ \color{blue}{\times} & \times & \times & \times & \times \\ & & \times & \times & \times \\ & & & \times & \times \end{bmatrix}}_{Q_1^* A Q_1} \longrightarrow \underbrace{\begin{bmatrix} \times & \times & \times & \times & \times \\ * & * & * & * & * \\ & * & * & * & * \\ & & \times & \times & \times \\ & & & \times & \times \end{bmatrix}}_{G_2^* Q_1^* A Q_1}$$

Bulge chasing implicit QR step

Since $A_1 = Q_{n-1}^* \cdots Q_2^* (Q_1^* A Q_1) Q_2 \cdots Q_{n-1}$ is Hessenberg but $Q_1^* A Q_1$ is NOT Hessenberg, it follows that the rotations Q_2, \dots, Q_{n-1} reduces $Q_1^* A Q_1$ to Hessenberg form. However, we can independently reduce $Q_1^* A Q_1$ to Hessenberg form without using the rotations Q_2, \dots, Q_{n-1} .

$$\underbrace{\begin{bmatrix} \times & \times & \times & \times & \times \\ \times & \times & \times & \times & \times \\ \textcolor{blue}{\times} & \times & \times & \times & \times \\ & & \times & \times & \times \\ & & & \times & \times \end{bmatrix}}_{Q_1^* A Q_1} \longrightarrow \underbrace{\begin{bmatrix} \times & \times & \times & \times & \times \\ * & * & * & * & * \\ & * & * & * & * \\ & & \times & \times & \times \\ & & & \times & \times \end{bmatrix}}_{G_2^* Q_1^* A Q_1} \longrightarrow \underbrace{\begin{bmatrix} \times & * & * & \times & \times \\ \times & * & * & \times & \times \\ & * & * & \times & \times \\ & * & * & \times & \times \\ & & & \times & \times \end{bmatrix}}_{G_2^* Q_1^* A Q_1 G_2}$$


Bulge chasing implicit QR step

Since $A_1 = Q_{n-1}^* \cdots Q_2^* (Q_1^* A Q_1) Q_2 \cdots Q_{n-1}$ is Hessenberg but $Q_1^* A Q_1$ is NOT Hessenberg, it follows that the rotations Q_2, \dots, Q_{n-1} reduces $Q_1^* A Q_1$ to Hessenberg form. However, we can independently reduce $Q_1^* A Q_1$ to Hessenberg form without using the rotations Q_2, \dots, Q_{n-1} .

$$\begin{array}{ccc}
 \underbrace{\begin{bmatrix} \times & \times & \times & \times & \times \\ \times & \times & \times & \times & \times \\ \textcolor{blue}{\times} & \times & \times & \times & \times \\ & & \times & \times & \times \\ & & & \times & \times \end{bmatrix}}_{Q_1^* A Q_1} & \longrightarrow & \underbrace{\begin{bmatrix} \times & \times & \times & \times & \times \\ * & * & * & * & * \\ & * & * & * & * \\ & & \times & \times & \times \\ & & & \times & \times \end{bmatrix}}_{G_2^* Q_1^* A Q_1} & \longrightarrow & \underbrace{\begin{bmatrix} \times & * & * & \times & \times \\ \times & * & * & \times & \times \\ & * & * & \times & \times \\ & * & * & \times & \times \\ & & & \times & \times \end{bmatrix}}_{G_2^* Q_1^* A Q_1 G_2} \\
 & & \longrightarrow & \underbrace{\begin{bmatrix} \times & \times & \times & \times & \times \\ \times & \times & \times & \times & \times \\ & * & * & * & * \\ & & * & * & * \\ & & & \times & \times \end{bmatrix}}_{G_3^* G_2^* Q_1^* A Q_1 G_2}
 \end{array}$$

Bulge chasing implicit QR step

Since $A_1 = Q_{n-1}^* \cdots Q_2^* (Q_1^* A Q_1) Q_2 \cdots Q_{n-1}$ is Hessenberg but $Q_1^* A Q_1$ is NOT Hessenberg, it follows that the rotations Q_2, \dots, Q_{n-1} reduces $Q_1^* A Q_1$ to Hessenberg form. However, we can independently reduce $Q_1^* A Q_1$ to Hessenberg form without using the rotations Q_2, \dots, Q_{n-1} .



Bulge chasing implicit QR step

Let G_2, \dots, G_{n-1} be Givens rotations such that $\hat{A}_1 := G_{n-1}^* \cdots G_2^* (Q_1^* A Q_1) G_2 \cdots G_{n-1}$ is in Hessenberg form. Set $Q := Q_1 Q_2 \cdots Q_{n-1}$ and $G := Q_1 G_2 \cdots G_{n-1}$.

Bulge chasing implicit QR step

Let G_2, \dots, G_{n-1} be Givens rotations such that $\hat{A}_1 := G_{n-1}^* \cdots G_2^* (Q_1^* A Q_1) G_2 \cdots G_{n-1}$ is in Hessenberg form. Set $Q := Q_1 Q_2 \cdots Q_{n-1}$ and $G := Q_1 G_2 \cdots G_{n-1}$.

Question: How are Q and G related? How are A_1 and \hat{A}_1 related?

Bulge chasing implicit QR step

Let G_2, \dots, G_{n-1} be Givens rotations such that $\hat{A}_1 := G_{n-1}^* \cdots G_2^* (Q_1^* A Q_1) G_2 \cdots G_{n-1}$ is in Hessenberg form. Set $Q := Q_1 Q_2 \cdots Q_{n-1}$ and $G := Q_1 G_2 \cdots G_{n-1}$.

Question: How are Q and G related? How are A_1 and \hat{A}_1 related? Since A proper Hessenberg, **Implicit Q-Theorem** there is a diagonal unitary matrix D such that $G = QD$ and $\hat{A}_1 = D^* A_1 D$.

Bulge chasing implicit QR step

Let G_2, \dots, G_{n-1} be Givens rotations such that $\hat{A}_1 := G_{n-1}^* \cdots G_2^* (Q_1^* A Q_1) G_2 \cdots G_{n-1}$ is in Hessenberg form. Set $Q := Q_1 Q_2 \cdots Q_{n-1}$ and $G := Q_1 G_2 \cdots G_{n-1}$.

Question: How are Q and G related? How are A_1 and \hat{A}_1 related? Since A proper Hessenberg, **Implicit Q-Theorem** there is a diagonal unitary matrix D such that $G = QD$ and $\hat{A}_1 = D^* A_1 D$.

Implicit Q-Theorem: Let $A \in \mathbb{C}^{n \times n}$. Let U and V be unitary matrices such that $U^* A U$ and $V^* A V$ are proper upper Hessenberg. If $U e_1 = V e_1$ then there is a diagonal unitary matrix D such that $V = U D$.

Bulge chasing implicit QR step

Let G_2, \dots, G_{n-1} be Givens rotations such that $\hat{A}_1 := G_{n-1}^* \cdots G_2^* (Q_1^* A Q_1) G_2 \cdots G_{n-1}$ is in Hessenberg form. Set $Q := Q_1 Q_2 \cdots Q_{n-1}$ and $G := Q_1 G_2 \cdots G_{n-1}$.

Question: How are Q and G related? How are A_1 and \hat{A}_1 related? Since A proper Hessenberg, **Implicit Q-Theorem** there is a diagonal unitary matrix D such that $G = QD$ and $\hat{A}_1 = D^* A_1 D$.

Implicit Q-Theorem: Let $A \in \mathbb{C}^{n \times n}$. Let U and V be unitary matrices such that $U^* A U$ and $V^* A V$ are proper upper Hessenberg. If $U e_1 = V e_1$ then there is a diagonal unitary matrix D such that $V = U D$.

Now observe that $A_1 = Q^* A Q$ and $\hat{A}_1 = G^* A G$ are proper upper Hessenberg.

Bulge chasing implicit QR step

Let G_2, \dots, G_{n-1} be Givens rotations such that $\hat{A}_1 := G_{n-1}^* \cdots G_2^* (Q_1^* A Q_1) G_2 \cdots G_{n-1}$ is in Hessenberg form. Set $Q := Q_1 Q_2 \cdots Q_{n-1}$ and $G := Q_1 G_2 \cdots G_{n-1}$.

Question: How are Q and G related? How are A_1 and \hat{A}_1 related? Since A proper Hessenberg, **Implicit Q-Theorem** there is a diagonal unitary matrix D such that $G = QD$ and $\hat{A}_1 = D^* A_1 D$.

Implicit Q-Theorem: Let $A \in \mathbb{C}^{n \times n}$. Let U and V be unitary matrices such that $U^* A U$ and $V^* A V$ are proper upper Hessenberg. If $U e_1 = V e_1$ then there is a diagonal unitary matrix D such that $V = U D$.

Now observe that $A_1 = Q^* A Q$ and $\hat{A}_1 = G^* A G$ are proper upper Hessenberg. Next, note that Q_j and G_j are Givens rotations in x_j - x_{j+1} plane.

Bulge chasing implicit QR step

Let G_2, \dots, G_{n-1} be Givens rotations such that $\hat{A}_1 := G_{n-1}^* \cdots G_2^* (Q_1^* A Q_1) G_2 \cdots G_{n-1}$ is in Hessenberg form. Set $Q := Q_1 Q_2 \cdots Q_{n-1}$ and $G := Q_1 G_2 \cdots G_{n-1}$.

Question: How are Q and G related? How are A_1 and \hat{A}_1 related? Since A proper Hessenberg, **Implicit Q-Theorem** there is a diagonal unitary matrix D such that $G = QD$ and $\hat{A}_1 = D^* A_1 D$.

Implicit Q-Theorem: Let $A \in \mathbb{C}^{n \times n}$. Let U and V be unitary matrices such that $U^* A U$ and $V^* A V$ are proper upper Hessenberg. If $U e_1 = V e_1$ then there is a diagonal unitary matrix D such that $V = U D$.

Now observe that $A_1 = Q^* A Q$ and $\hat{A}_1 = G^* A G$ are proper upper Hessenberg. Next, note that Q_j and G_j are Givens rotations in x_j - x_{j+1} plane. Hence $Q_j e_1 = e_1 = G_j e_1$ for $j = 2 : n-1$.

Bulge chasing implicit QR step

Let G_2, \dots, G_{n-1} be Givens rotations such that $\hat{A}_1 := G_{n-1}^* \cdots G_2^* (Q_1^* A Q_1) G_2 \cdots G_{n-1}$ is in Hessenberg form. Set $Q := Q_1 Q_2 \cdots Q_{n-1}$ and $G := Q_1 G_2 \cdots G_{n-1}$.

Question: How are Q and G related? How are A_1 and \hat{A}_1 related? Since A proper Hessenberg, **Implicit Q-Theorem** there is a diagonal unitary matrix D such that $G = QD$ and $\hat{A}_1 = D^* A_1 D$.

Implicit Q-Theorem: Let $A \in \mathbb{C}^{n \times n}$. Let U and V be unitary matrices such that $U^* A U$ and $V^* A V$ are proper upper Hessenberg. If $U e_1 = V e_1$ then there is a diagonal unitary matrix D such that $V = U D$.

Now observe that $A_1 = Q^* A Q$ and $\hat{A}_1 = G^* A G$ are proper upper Hessenberg. Next, note that Q_j and G_j are Givens rotations in x_j - x_{j+1} plane. Hence $Q_j e_1 = e_1 = G_j e_1$ for $j = 2 : n-1$. Thus

$$Q e_1 = Q_1 Q_2 \cdots Q_{n-1} e_1 = Q_1 e_1 = Q_1 G_2 \cdots G_{n-1} e_1 = G e_1.$$

Bulge chasing implicit QR step

Let G_2, \dots, G_{n-1} be Givens rotations such that $\hat{A}_1 := G_{n-1}^* \cdots G_2^* (Q_1^* A Q_1) G_2 \cdots G_{n-1}$ is in Hessenberg form. Set $Q := Q_1 Q_2 \cdots Q_{n-1}$ and $G := Q_1 G_2 \cdots G_{n-1}$.

Question: How are Q and G related? How are A_1 and \hat{A}_1 related? Since A proper Hessenberg, **Implicit Q-Theorem** there is a diagonal unitary matrix D such that $G = QD$ and $\hat{A}_1 = D^* A_1 D$.

Implicit Q-Theorem: Let $A \in \mathbb{C}^{n \times n}$. Let U and V be unitary matrices such that $U^* A U$ and $V^* A V$ are proper upper Hessenberg. If $U e_1 = V e_1$ then there is a diagonal unitary matrix D such that $V = U D$.

Now observe that $A_1 = Q^* A Q$ and $\hat{A}_1 = G^* A G$ are proper upper Hessenberg. Next, note that Q_j and G_j are Givens rotations in x_j - x_{j+1} plane. Hence $Q_j e_1 = e_1 = G_j e_1$ for $j = 2 : n-1$. Thus

$$Q e_1 = Q_1 Q_2 \cdots Q_{n-1} e_1 = Q_1 e_1 = Q_1 G_2 \cdots G_{n-1} e_1 = G e_1.$$

Therefore, by Implicit-Q Theorem, $G = QD$ and $\hat{A}_1 = G^* A G = D^* Q^* A Q D = D^* A_1 D$.

Single shift Implicit QR algorithm

Algorithm. (Single shift implicit QR algorithm)

Input: An $n \times n$ Hessenberg matrix A

Output: Upper triangular matrix $T = Q^*AQ$

Repeat until convergence

(i) Choose a shift μ and construct a Givens rotation Q_1 such that

$$Q_1^*(A - \mu I)e_1 = Q_1^* \begin{bmatrix} a_{11} - \mu \\ a_{21} \\ 0 \\ \vdots \\ 0 \end{bmatrix} = \begin{bmatrix} * \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

and compute $Q_1^*AQ_1$. This destroys Hessenberg form.

Single shift Implicit QR algorithm

Algorithm. (Single shift implicit QR algorithm)

Input: An $n \times n$ Hessenberg matrix A

Output: Upper triangular matrix $T = Q^* A Q$

Repeat until convergence

(i) Choose a shift μ and construct a Givens rotation Q_1 such that

$$Q_1^*(A - \mu I)e_1 = Q_1^* \begin{bmatrix} a_{11} - \mu \\ a_{21} \\ 0 \\ \vdots \\ 0 \end{bmatrix} = \begin{bmatrix} * \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

and compute $Q_1^* A Q_1$. This destroys Hessenberg form.

(ii) Construct Givens rotations Q_2, \dots, Q_{n-1} such that

$$A \leftarrow Q_{n-1}^* \cdots Q_2^* Q_1^* A Q_1 Q_2 \cdots Q_{n-1}$$

is upper Hessenberg.

Double shift explicit QR Algorithm

Let A be Hessenberg and let μ_1 and μ_2 be scalars. Consider two steps of QR iteration on A .

- $A - \mu_1 I = Q_1 R_1$, $A_1 = R_1 Q_1 + \mu_1 I$ % first QR step

Double shift explicit QR Algorithm

Let A be Hessenberg and let μ_1 and μ_2 be scalars. Consider two steps of QR iteration on A .

- $A - \mu_1 I = Q_1 R_1, \quad A_1 = R_1 Q_1 + \mu_1 I$ % first QR step
- $A_1 - \mu_2 I = Q_2 R_2, \quad A_2 = R_2 Q_2 + \mu_2 I$ % 2nd QR step

Double shift explicit QR Algorithm

Let A be Hessenberg and let μ_1 and μ_2 be scalars. Consider two steps of QR iteration on A .

- $A - \mu_1 I = Q_1 R_1, \quad A_1 = R_1 Q_1 + \mu_1 I$ % first QR step
- $A_1 - \mu_2 I = Q_2 R_2, \quad A_2 = R_2 Q_2 + \mu_2 I$ % 2nd QR step

Define $Q := Q_1 Q_2$ and $R := R_2 R_1$.

Double shift explicit QR Algorithm

Let A be Hessenberg and let μ_1 and μ_2 be scalars. Consider two steps of QR iteration on A .

- $A - \mu_1 I = Q_1 R_1, \quad A_1 = R_1 Q_1 + \mu_1 I \quad \% \text{ first QR step}$
- $A_1 - \mu_2 I = Q_2 R_2, \quad A_2 = R_2 Q_2 + \mu_2 I \quad \% \text{ 2nd QR step}$

Define $Q := Q_1 Q_2$ and $R := R_2 R_1$. Then $A_2 = Q_2^* A_1 Q_2 = Q_2^* Q_1^* A Q_1 Q_2 = Q^* A Q$ and

Double shift explicit QR Algorithm

Let A be Hessenberg and let μ_1 and μ_2 be scalars. Consider two steps of QR iteration on A .

- $A - \mu_1 I = Q_1 R_1, \quad A_1 = R_1 Q_1 + \mu_1 I$ % first QR step
- $A_1 - \mu_2 I = Q_2 R_2, \quad A_2 = R_2 Q_2 + \mu_2 I$ % 2nd QR step

Define $Q := Q_1 Q_2$ and $R := R_2 R_1$. Then $A_2 = Q_2^* A_1 Q_2 = Q_2^* Q_1^* A Q_1 Q_2 = Q^* A Q$ and

$$A^2 - (\mu_1 + \mu_2)A + \mu_1 \mu_2 I = (A - \mu_2 I)(A - \mu_1 I) = (A - \mu_2 I)Q_1 R_1$$

Double shift explicit QR Algorithm

Let A be Hessenberg and let μ_1 and μ_2 be scalars. Consider two steps of QR iteration on A .

- $A - \mu_1 I = Q_1 R_1$, $A_1 = R_1 Q_1 + \mu_1 I$ % first QR step
- $A_1 - \mu_2 I = Q_2 R_2$, $A_2 = R_2 Q_2 + \mu_2 I$ % 2nd QR step

Define $Q := Q_1 Q_2$ and $R := R_2 R_1$. Then $A_2 = Q_2^* A_1 Q_2 = Q_2^* Q_1^* A Q_1 Q_2 = Q^* A Q$ and

$$\begin{aligned} A^2 - (\mu_1 + \mu_2)A + \mu_1 \mu_2 I &= (A - \mu_2 I)(A - \mu_1 I) = (A - \mu_2 I)Q_1 R_1 \\ &= Q_1 Q_1^* (A - \mu_2 I)Q_1 R_1 = Q_1 (A_1 - \mu_2 I)R_1 \end{aligned}$$

Double shift explicit QR Algorithm

Let A be Hessenberg and let μ_1 and μ_2 be scalars. Consider two steps of QR iteration on A .

- $A - \mu_1 I = Q_1 R_1$, $A_1 = R_1 Q_1 + \mu_1 I$ % first QR step
- $A_1 - \mu_2 I = Q_2 R_2$, $A_2 = R_2 Q_2 + \mu_2 I$ % 2nd QR step

Define $Q := Q_1 Q_2$ and $R := R_2 R_1$. Then $A_2 = Q_2^* A_1 Q_2 = Q_2^* Q_1^* A Q_1 Q_2 = Q^* A Q$ and

$$\begin{aligned} A^2 - (\mu_1 + \mu_2)A + \mu_1 \mu_2 I &= (A - \mu_2 I)(A - \mu_1 I) = (A - \mu_2 I)Q_1 R_1 \\ &= Q_1 Q_1^* (A - \mu_2 I)Q_1 R_1 = Q_1 (A_1 - \mu_2 I)R_1 \\ &= Q_1 Q_2 R_2 R_1 = QR. \end{aligned}$$

Double shift explicit QR Algorithm

Let A be Hessenberg and let μ_1 and μ_2 be scalars. Consider two steps of QR iteration on A .

- $A - \mu_1 I = Q_1 R_1$, $A_1 = R_1 Q_1 + \mu_1 I$ % first QR step
- $A_1 - \mu_2 I = Q_2 R_2$, $A_2 = R_2 Q_2 + \mu_2 I$ % 2nd QR step

Define $Q := Q_1 Q_2$ and $R := R_2 R_1$. Then $A_2 = Q_2^* A_1 Q_2 = Q_2^* Q_1^* A Q_1 Q_2 = Q^* A Q$ and

$$\begin{aligned} A^2 - (\mu_1 + \mu_2)A + \mu_1 \mu_2 I &= (A - \mu_2 I)(A - \mu_1 I) = (A - \mu_2 I)Q_1 R_1 \\ &= Q_1 Q_1^* (A - \mu_2 I)Q_1 R_1 = Q_1 (A_1 - \mu_2 I)R_1 \\ &= Q_1 Q_2 R_2 R_1 = QR. \end{aligned}$$

This yields double shift QR algorithm.

Double shift explicit QR Algorithm:

- Choose shifts μ_1 and μ_2
- $A^2 - (\mu_1 + \mu_2)A + \mu_1 \mu_2 I = QR = Q_1 Q_2 \cdots Q_{n-1} R$ % QR factorization

Double shift explicit QR Algorithm

Let A be Hessenberg and let μ_1 and μ_2 be scalars. Consider two steps of QR iteration on A .

- $A - \mu_1 I = Q_1 R_1$, $A_1 = R_1 Q_1 + \mu_1 I$ % first QR step
- $A_1 - \mu_2 I = Q_2 R_2$, $A_2 = R_2 Q_2 + \mu_2 I$ % 2nd QR step

Define $Q := Q_1 Q_2$ and $R := R_2 R_1$. Then $A_2 = Q_2^* A_1 Q_2 = Q_2^* Q_1^* A Q_1 Q_2 = Q^* A Q$ and

$$\begin{aligned} A^2 - (\mu_1 + \mu_2)A + \mu_1 \mu_2 I &= (A - \mu_2 I)(A - \mu_1 I) = (A - \mu_2 I)Q_1 R_1 \\ &= Q_1 Q_1^* (A - \mu_2 I)Q_1 R_1 = Q_1 (A_1 - \mu_2 I)R_1 \\ &= Q_1 Q_2 R_2 R_1 = QR. \end{aligned}$$

This yields double shift QR algorithm.

Double shift explicit QR Algorithm:

- Choose shifts μ_1 and μ_2
- $A^2 - (\mu_1 + \mu_2)A + \mu_1 \mu_2 I = QR = Q_1 Q_2 \cdots Q_{n-1} R$ % QR factorization
- $A \leftarrow Q^* A Q = Q_{n-1}^* \cdots Q_2^* (Q_1^* A Q_1) Q_2 \cdots Q_{n-1}$ % similarity transformation

Double shift Implicit QR algorithm

Algorithm. (Double shift implicit QR algorithm)

Input: An $n \times n$ Hessenberg matrix A

Output: Upper triangular matrix $T = Q^*AQ$

(i) Choose shifts μ_1 and μ_2 and construct a Householder reflector Q_1 such that

$$Q_1^*(A^2 - (\mu_1 + \mu_2)A + \mu_1\mu_2 I)e_1 = Q_1^* \begin{bmatrix} a_{21} \left[\frac{a_{11}^2 - (\mu_1 + \mu_2)a_{11} + \mu_1\mu_2}{a_{21}} + a_{12} \right] \\ a_{21}[a_{11} + a_{22} - (\mu_1 + \mu_2)] \\ a_{21}a_{32} \\ 0 \\ \vdots \\ 0 \end{bmatrix} = \begin{bmatrix} * \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

and compute $Q_1^*AQ_1$. This destroys Hessenberg form.

Double shift Implicit QR algorithm

Algorithm. (Double shift implicit QR algorithm)

Input: An $n \times n$ Hessenberg matrix A

Output: Upper triangular matrix $T = Q^* A Q$

(i) Choose shifts μ_1 and μ_2 and construct a Householder reflector Q_1 such that

$$Q_1^*(A^2 - (\mu_1 + \mu_2)A + \mu_1\mu_2 I)e_1 = Q_1^* \begin{bmatrix} a_{21} \left[\frac{a_{11}^2 - (\mu_1 + \mu_2)a_{11} + \mu_1\mu_2}{a_{21}} + a_{12} \right] \\ a_{21}[a_{11} + a_{22} - (\mu_1 + \mu_2)] \\ a_{21}a_{32} \\ 0 \\ \vdots \\ 0 \end{bmatrix} = \begin{bmatrix} * \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

and compute $Q_1^* A Q_1$. This destroys Hessenberg form.

(ii) Construct Householder reflectors Q_2, \dots, Q_{n-1} such that

$$A \leftarrow Q_{n-1}^* \cdots Q_2^* Q_1^* A Q_1 Q_2 \cdots Q_{n-1}$$

is upper Hessenberg.