

# QR factorization by Givens rotations

Rafikul Alam  
Department of Mathematics  
Indian Institute of Technology Guwahati  
Guwahati - 781039, INDIA

# Outline

- Rotation in  $\mathbb{R}^2$
- Givens rotation in  $\mathbb{R}^n$
- QR factorization by Givens rotations

# Rotation in $\mathbb{R}^2$

**Definition:** Let  $\theta \in [0, 2\pi]$ . A **rotation** in  $\mathbb{R}^2$  is a matrix  $G(\theta) \in \mathbb{R}^{2 \times 2}$  that rotates each vector in  $\mathbb{R}^2$  by an angle  $\theta$  in the **anti-clock-wise direction**.

# Rotation in $\mathbb{R}^2$

**Definition:** Let  $\theta \in [0, 2\pi]$ . A **rotation** in  $\mathbb{R}^2$  is a matrix  $G(\theta) \in \mathbb{R}^{2 \times 2}$  that rotates each vector in  $\mathbb{R}^2$  by an angle  $\theta$  in the **anti-clock-wise direction**.

Obviously  $G(\theta) : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$  is a linear transformation and  $G(\theta)v$  is the anti-clock-wise rotation of  $v$  by an angle  $\theta$ .

# Rotation in $\mathbb{R}^2$

**Definition:** Let  $\theta \in [0, 2\pi]$ . A **rotation** in  $\mathbb{R}^2$  is a matrix  $G(\theta) \in \mathbb{R}^{2 \times 2}$  that rotates each vector in  $\mathbb{R}^2$  by an angle  $\theta$  in the **anti-clock-wise direction**.

Obviously  $G(\theta) : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$  is a linear transformation and  $G(\theta)v$  is the anti-clock-wise rotation of  $v$  by an angle  $\theta$ . The rotation  $G(\theta)$  is given by

$$G(\theta) = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} = \begin{bmatrix} c & -s \\ s & c \end{bmatrix}.$$

# Rotation in $\mathbb{R}^2$

**Definition:** Let  $\theta \in [0, 2\pi]$ . A **rotation** in  $\mathbb{R}^2$  is a matrix  $G(\theta) \in \mathbb{R}^{2 \times 2}$  that rotates each vector in  $\mathbb{R}^2$  by an angle  $\theta$  in the **anti-clock-wise direction**.

Obviously  $G(\theta) : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$  is a linear transformation and  $G(\theta)v$  is the anti-clock-wise rotation of  $v$  by an angle  $\theta$ . The rotation  $G(\theta)$  is given by

$$G(\theta) = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} = \begin{bmatrix} c & -s \\ s & c \end{bmatrix}.$$

Note that  $e_1$  and  $G(\theta)e_1$  are points on the unit circle  $x^2 + y^2 = 1$ .

# Rotation in $\mathbb{R}^2$

**Definition:** Let  $\theta \in [0, 2\pi]$ . A **rotation** in  $\mathbb{R}^2$  is a matrix  $G(\theta) \in \mathbb{R}^{2 \times 2}$  that rotates each vector in  $\mathbb{R}^2$  by an angle  $\theta$  in the **anti-clock-wise direction**.

Obviously  $G(\theta) : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$  is a linear transformation and  $G(\theta)v$  is the anti-clock-wise rotation of  $v$  by an angle  $\theta$ . The rotation  $G(\theta)$  is given by

$$G(\theta) = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} = \begin{bmatrix} c & -s \\ s & c \end{bmatrix}.$$

Note that  $e_1$  and  $G(\theta)e_1$  are points on the unit circle  $x^2 + y^2 = 1$ . Hence  $G(\theta)e_1 = \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix}$ . Also, note that  $G(\theta)e_2 = G(\pi/2 + \theta)e_1$ .

# Rotation in $\mathbb{R}^2$

**Definition:** Let  $\theta \in [0, 2\pi]$ . A **rotation** in  $\mathbb{R}^2$  is a matrix  $G(\theta) \in \mathbb{R}^{2 \times 2}$  that rotates each vector in  $\mathbb{R}^2$  by an angle  $\theta$  in the **anti-clock-wise direction**.

Obviously  $G(\theta) : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$  is a linear transformation and  $G(\theta)v$  is the anti-clock-wise rotation of  $v$  by an angle  $\theta$ . The rotation  $G(\theta)$  is given by

$$G(\theta) = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} = \begin{bmatrix} c & -s \\ s & c \end{bmatrix}.$$

Note that  $e_1$  and  $G(\theta)e_1$  are points on the unit circle  $x^2 + y^2 = 1$ . Hence  $G(\theta)e_1 = \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix}$ . Also, note that  $G(\theta)e_2 = G(\pi/2 + \theta)e_1$ .

Consequently,  $G(\theta)e_2 = G(\pi/2 + \theta)e_1 = \begin{bmatrix} \cos(\pi/2 + \theta) \\ \sin(\pi/2 + \theta) \end{bmatrix} = \begin{bmatrix} -\sin \theta \\ \cos \theta \end{bmatrix}$ .



## Rotation in $\mathbb{R}^2$

**Definition:** Let  $\theta \in [0, 2\pi]$ . A **rotation** in  $\mathbb{R}^2$  is a matrix  $G(\theta) \in \mathbb{R}^{2 \times 2}$  that rotates each vector in  $\mathbb{R}^2$  by an angle  $\theta$  in the **anti-clock-wise direction**.

Obviously  $G(\theta) : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$  is a linear transformation and  $G(\theta)v$  is the anti-clock-wise rotation of  $v$  by an angle  $\theta$ . The rotation  $G(\theta)$  is given by

$$G(\theta) = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} = \begin{bmatrix} c & -s \\ s & c \end{bmatrix}.$$

Note that  $e_1$  and  $G(\theta)e_1$  are points on the unit circle  $x^2 + y^2 = 1$ . Hence  $G(\theta)e_1 = \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix}$ . Also, note that  $G(\theta)e_2 = G(\pi/2 + \theta)e_1$ .

Consequently,  $G(\theta)e_2 = G(\pi/2 + \theta)e_1 = \begin{bmatrix} \cos(\pi/2 + \theta) \\ \sin(\pi/2 + \theta) \end{bmatrix} = \begin{bmatrix} -\sin \theta \\ \cos \theta \end{bmatrix}$ .


Hence we have  $G(\theta) = [G(\theta)e_1 \quad G(\theta)e_2] = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$ .

## Plane rotation in $\mathbb{R}^2$

We have  $G(\theta) \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} x_1 \cos \theta - x_2 \sin \theta \\ x_1 \sin \theta + x_2 \cos \theta \end{bmatrix}.$


## Plane rotation in $\mathbb{R}^2$

We have  $G(\theta) \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} x_1 \cos \theta - x_2 \sin \theta \\ x_1 \sin \theta + x_2 \cos \theta \end{bmatrix}$ .




## Plane rotation in $\mathbb{R}^2$

We have  $G(\theta) \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} x_1 \cos \theta - x_2 \sin \theta \\ x_1 \sin \theta + x_2 \cos \theta \end{bmatrix}$ .




## Plane rotation in $\mathbb{R}^2$

We have  $G(\theta) \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} x_1 \cos \theta - x_2 \sin \theta \\ x_1 \sin \theta + x_2 \cos \theta \end{bmatrix}$ .




## Plane rotation in $\mathbb{R}^2$

We have  $G(\theta) \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} x_1 \cos \theta - x_2 \sin \theta \\ x_1 \sin \theta + x_2 \cos \theta \end{bmatrix}$ .



## Plane rotation in $\mathbb{R}^2$

We have  $G(\theta) \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} x_1 \cos \theta - x_2 \sin \theta \\ x_1 \sin \theta + x_2 \cos \theta \end{bmatrix}$ .



The rotation  $G(\theta)$ . Note that  $G(\theta)e_2 = G(\pi/2 + \theta)e_1$ .

# Properties of rotation

- Consider  $G(\theta) := \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$ . Then  $G(0) = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I$  and  $G(\theta)^\top = G(-\theta)$ . Thus  $G(\theta)^\top$  is a **clock-wise rotation** by an angle  $\theta$ .



# Properties of rotation

- Consider  $G(\theta) := \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$ . Then  $G(0) = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I$  and  $G(\theta)^\top = G(-\theta)$ . Thus  $G(\theta)^\top$  is a **clock-wise rotation** by an angle  $\theta$ .
- Fact:** Let  $G(\theta)$  and  $G(\phi)$  be rotations in  $\mathbb{R}^2$ . Then  $G(\phi)G(\theta)$  is a rotation in  $\mathbb{R}^2$  and  **$G(\theta)G(\phi) = G(\phi)G(\theta) = G(\theta + \phi)$** .

# Properties of rotation

- Consider  $G(\theta) := \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$ . Then  $G(0) = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I$  and  $G(\theta)^\top = G(-\theta)$ . Thus  $G(\theta)^\top$  is a **clock-wise rotation** by an angle  $\theta$ .
- **Fact:** Let  $G(\theta)$  and  $G(\phi)$  be rotations in  $\mathbb{R}^2$ . Then  $G(\phi)G(\theta)$  is a rotation in  $\mathbb{R}^2$  and  $G(\theta)G(\phi) = G(\phi)G(\theta) = G(\theta + \phi)$ .
- **Proof:** Note that  $G(\phi)G(\theta)v$  is the anti-clock-wise rotation of  $v$  by an angle  $\theta$  followed by an angle  $\phi$ . Thus  $G(\phi)G(\theta)v = G(\phi + \theta)v$  for all  $v \in \mathbb{R}^2$ . Hence  $G(\phi)G(\theta) = G(\theta + \phi)$ . ■

# Properties of rotation

- Consider  $G(\theta) := \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$ . Then  $G(0) = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I$  and  $G(\theta)^\top = G(-\theta)$ . Thus  $G(\theta)^\top$  is a **clock-wise rotation** by an angle  $\theta$ .
- **Fact:** Let  $G(\theta)$  and  $G(\phi)$  be rotations in  $\mathbb{R}^2$ . Then  $G(\phi)G(\theta)$  is a rotation in  $\mathbb{R}^2$  and  $G(\theta)G(\phi) = G(\phi)G(\theta) = G(\theta + \phi)$ .
- **Proof:** Note that  $G(\phi)G(\theta)v$  is the anti-clock-wise rotation of  $v$  by an angle  $\theta$  followed by an angle  $\phi$ . Thus  $G(\phi)G(\theta)v = G(\phi + \theta)v$  for all  $v \in \mathbb{R}^2$ . Hence  $G(\phi)G(\theta) = G(\theta + \phi)$ . ■
- We have  $G(\theta)G(\theta)^\top = G(\theta)G(-\theta) = G(0) = I$  which shows that  $G(\theta)$  is **unitary** and  $G(\theta)^{-1} = G(\theta)^\top = G(-\theta)$ .

# Properties of rotation

- Consider  $G(\theta) := \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$ . Then  $G(0) = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I$  and  $G(\theta)^\top = G(-\theta)$ . Thus  $G(\theta)^\top$  is a **clock-wise rotation** by an angle  $\theta$ .
- **Fact:** Let  $G(\theta)$  and  $G(\phi)$  be rotations in  $\mathbb{R}^2$ . Then  $G(\phi)G(\theta)$  is a rotation in  $\mathbb{R}^2$  and  $G(\theta)G(\phi) = G(\phi)G(\theta) = G(\theta + \phi)$ .
- **Proof:** Note that  $G(\phi)G(\theta)v$  is the anti-clock-wise rotation of  $v$  by an angle  $\theta$  followed by an angle  $\phi$ . Thus  $G(\phi)G(\theta)v = G(\phi + \theta)v$  for all  $v \in \mathbb{R}^2$ . Hence  $G(\phi)G(\theta) = G(\theta + \phi)$ . ■
- We have  $G(\theta)G(\theta)^\top = G(\theta)G(-\theta) = G(0) = I$  which shows that  $G(\theta)$  is **unitary** and  $G(\theta)^{-1} = G(\theta)^\top = G(-\theta)$ .

**Example:**  $G(\pi/6) = \begin{bmatrix} \frac{\sqrt{3}}{2} & -\frac{1}{2} \\ \frac{1}{2} & \frac{\sqrt{3}}{2} \end{bmatrix}$  and  $G(\pi/2) = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$ .

## Creating zero by rotation

For simplicity, we write  $G := \begin{bmatrix} c & -s \\ s & c \end{bmatrix}$ , where  $c := \cos \theta$  and  $s = \sin \theta$ .

## Creating zero by rotation

For simplicity, we write  $G := \begin{bmatrix} c & -s \\ s & c \end{bmatrix}$ , where  $c := \cos \theta$  and  $s = \sin \theta$ .

Moreover, we require only  $c$  and  $s$  and not the angle  $\theta$ . Hence in order to generate  $G$ , we need  $c$  and  $s$  such that  $c^2 + s^2 = 1$ .

## Creating zero by rotation

For simplicity, we write  $G := \begin{bmatrix} c & -s \\ s & c \end{bmatrix}$ , where  $c := \cos \theta$  and  $s = \sin \theta$ .

Moreover, we require only  $c$  and  $s$  and not the angle  $\theta$ . Hence in order to generate  $G$ , we need  $c$  and  $s$  such that  $c^2 + s^2 = 1$ .

Let  $v := \begin{bmatrix} v_1 \\ v_2 \end{bmatrix}$ . Define  $c := \frac{v_1}{\sqrt{v_1^2 + v_2^2}}$  and  $s := \frac{v_2}{\sqrt{v_1^2 + v_2^2}}$ . Consider the rotation  $G := \begin{bmatrix} c & -s \\ s & c \end{bmatrix}$ . Then we have  $G^T v = \begin{bmatrix} \|v\|_2 \\ 0 \end{bmatrix} = \|v\|_2 e_1$ .

## Creating zero by rotation

For simplicity, we write  $G := \begin{bmatrix} c & -s \\ s & c \end{bmatrix}$ , where  $c := \cos \theta$  and  $s = \sin \theta$ .

Moreover, we require only  $c$  and  $s$  and not the angle  $\theta$ . Hence in order to generate  $G$ , we need  $c$  and  $s$  such that  $c^2 + s^2 = 1$ .

Let  $v := \begin{bmatrix} v_1 \\ v_2 \end{bmatrix}$ . Define  $c := \frac{v_1}{\sqrt{v_1^2 + v_2^2}}$  and  $s := \frac{v_2}{\sqrt{v_1^2 + v_2^2}}$ . Consider the rotation  $G := \begin{bmatrix} c & -s \\ s & c \end{bmatrix}$ . Then we have  $G^\top v = \begin{bmatrix} \|v\|_2 \\ 0 \end{bmatrix} = \|v\|_2 e_1$ .

Note that  $v$  makes an angle  $\theta := \arctan(v_2/v_1)$  with  $e_1$  (the  $x$ -axis) and  $G^\top$  rotates  $v$  clock-wise by an angle  $\theta$  to align with  $e_1$ .



## Creating zero by rotation

For simplicity, we write  $G := \begin{bmatrix} c & -s \\ s & c \end{bmatrix}$ , where  $c := \cos \theta$  and  $s = \sin \theta$ .

Moreover, we require only  $c$  and  $s$  and not the angle  $\theta$ . Hence in order to generate  $G$ , we need  $c$  and  $s$  such that  $c^2 + s^2 = 1$ .

Let  $v := \begin{bmatrix} v_1 \\ v_2 \end{bmatrix}$ . Define  $c := \frac{v_1}{\sqrt{v_1^2 + v_2^2}}$  and  $s := \frac{v_2}{\sqrt{v_1^2 + v_2^2}}$ . Consider the rotation  $G := \begin{bmatrix} c & -s \\ s & c \end{bmatrix}$ . Then we have  $G^\top v = \begin{bmatrix} \|v\|_2 \\ 0 \end{bmatrix} = \|v\|_2 e_1$ .

Note that  $v$  makes an angle  $\theta := \arctan(v_2/v_1)$  with  $e_1$  (the  $x$ -axis) and  $G^\top$  rotates  $v$  clock-wise by an angle  $\theta$  to align with  $e_1$ . Also, we can construct a rotation  $G$  such that  $Gv = -\|v\|_2 e_1$ .

## Creating zero by rotation

For simplicity, we write  $G := \begin{bmatrix} c & -s \\ s & c \end{bmatrix}$ , where  $c := \cos \theta$  and  $s = \sin \theta$ .

Moreover, we require only  $c$  and  $s$  and not the angle  $\theta$ . Hence in order to generate  $G$ , we need  $c$  and  $s$  such that  $c^2 + s^2 = 1$ .

Let  $v := \begin{bmatrix} v_1 \\ v_2 \end{bmatrix}$ . Define  $c := \frac{v_1}{\sqrt{v_1^2 + v_2^2}}$  and  $s := \frac{v_2}{\sqrt{v_1^2 + v_2^2}}$ . Consider the rotation  $G := \begin{bmatrix} c & -s \\ s & c \end{bmatrix}$ . Then we have  $G^\top v = \begin{bmatrix} \|v\|_2 \\ 0 \end{bmatrix} = \|v\|_2 e_1$ .

Note that  $v$  makes an angle  $\theta := \arctan(v_2/v_1)$  with  $e_1$  (the  $x$ -axis) and  $G^\top$  rotates  $v$  clock-wise by an angle  $\theta$  to align with  $e_1$ . Also, we can construct a rotation  $G$  such that  $Gv = -\|v\|_2 e_1$ .

Similarly, we can construct a rotation  $G$  such that  $Gv = \|v\|_2 e_2$  or  $G^\top v = -\|v\|_2 e_2$ .

## Creating zero by rotation

For simplicity, we write  $G := \begin{bmatrix} c & -s \\ s & c \end{bmatrix}$ , where  $c := \cos \theta$  and  $s = \sin \theta$ .

Moreover, we require only  $c$  and  $s$  and not the angle  $\theta$ . Hence in order to generate  $G$ , we need  $c$  and  $s$  such that  $c^2 + s^2 = 1$ .

Let  $v := \begin{bmatrix} v_1 \\ v_2 \end{bmatrix}$ . Define  $c := \frac{v_1}{\sqrt{v_1^2 + v_2^2}}$  and  $s := \frac{v_2}{\sqrt{v_1^2 + v_2^2}}$ . Consider the rotation  $G := \begin{bmatrix} c & -s \\ s & c \end{bmatrix}$ . Then we have  $G^T v = \begin{bmatrix} \|v\|_2 \\ 0 \end{bmatrix} = \|v\|_2 e_1$ .

Note that  $v$  makes an angle  $\theta := \arctan(v_2/v_1)$  with  $e_1$  (the  $x$ -axis) and  $G^T$  rotates  $v$  clock-wise by an angle  $\theta$  to align with  $e_1$ . Also, we can construct a rotation  $G$  such that  $Gv = -\|v\|_2 e_1$ . Take  $c := -v_1/(\sqrt{v_1^2 + v_2^2})$ ,  $s := v_2/(\sqrt{v_1^2 + v_2^2})$ .

Similarly, we can construct a rotation  $G$  such that  $Gv = \|v\|_2 e_2$  or  $G^T v = -\|v\|_2 e_2$ .

**Example:** Consider  $v := \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ . Then  $\begin{bmatrix} c & -s \\ s & c \end{bmatrix}^T \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} \sqrt{2} \\ 0 \end{bmatrix}$ , where  $c = 1/\sqrt{2} = s$ . Note that  $\theta = \pi/4$ .

# Givens rotations in $\mathbb{R}^3$

**Definition:** A **Givens rotation** in  $\mathbb{R}^3$  is a rotation in the coordinate plane, that is, in the plane **spanned by two coordinate axes**.

## Givens rotations in $\mathbb{R}^3$

**Definition:** A **Givens rotation** in  $\mathbb{R}^3$  is a rotation in the coordinate plane, that is, in the plane **spanned by two coordinate axes**.

Let  $\theta \in [0, 2\pi]$ . Then a **Givens rotation** in the  $x_1$ - $x_2$  plane in  $\mathbb{R}^3$  is a matrix  $G_{12}(\theta) \in \mathbb{R}^{3 \times 3}$  that rotates each vector in the  $x_1$ - $x_2$  plane in  $\mathbb{R}^3$  by an angle  $\theta$  in the **anti-clock-wise direction** and  $G_{12}(\theta)e_3 = e_3$ .

## Givens rotations in $\mathbb{R}^3$

**Definition:** A **Givens rotation** in  $\mathbb{R}^3$  is a rotation in the coordinate plane, that is, in the plane **spanned by two coordinate axes**.

Let  $\theta \in [0, 2\pi]$ . Then a **Givens rotation** in the  $x_1$ - $x_2$  plane in  $\mathbb{R}^3$  is a matrix  $G_{12}(\theta) \in \mathbb{R}^{3 \times 3}$  that rotates each vector in the  $x_1$ - $x_2$  plane in  $\mathbb{R}^3$  by an angle  $\theta$  in the **anti-clock-wise direction** and  $G_{12}(\theta)e_3 = e_3$ .

Consequently, the rotation  $G_{12}(\theta)$  is given by

$$G_{12}(\theta) = \left[ \begin{array}{cc|c} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ \hline 0 & 0 & 1 \end{array} \right] = \left[ \begin{array}{cc|c} c & -s & 0 \\ s & c & 0 \\ \hline 0 & 0 & 1 \end{array} \right].$$

## Givens rotations in $\mathbb{R}^3$

**Definition:** A **Givens rotation** in  $\mathbb{R}^3$  is a rotation in the coordinate plane, that is, in the plane **spanned by two coordinate axes**.

Let  $\theta \in [0, 2\pi]$ . Then a **Givens rotation** in the  $x_1$ - $x_2$  plane in  $\mathbb{R}^3$  is a matrix  $G_{12}(\theta) \in \mathbb{R}^{3 \times 3}$  that rotates each vector in the  $x_1$ - $x_2$  plane in  $\mathbb{R}^3$  by an angle  $\theta$  in the **anti-clock-wise direction** and  $G_{12}(\theta)e_3 = e_3$ .

Consequently, the rotation  $G_{12}(\theta)$  is given by

$$G_{12}(\theta) = \left[ \begin{array}{cc|c} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ \hline 0 & 0 & 1 \end{array} \right] = \left[ \begin{array}{cc|c} c & -s & 0 \\ s & c & 0 \\ \hline 0 & 0 & 1 \end{array} \right].$$

Similarly, the Givens rotation  $G_{23}(\theta) \in \mathbb{R}^{3 \times 3}$  in the  $x_2$ - $x_3$  plane leaves  $e_1$  unchanged, that is,  $G_{23}(\theta)e_1 = e_1$ , and is given by

## Givens rotations in $\mathbb{R}^3$

**Definition:** A **Givens rotation** in  $\mathbb{R}^3$  is a rotation in the coordinate plane, that is, in the plane **spanned by two coordinate axes**.

Let  $\theta \in [0, 2\pi]$ . Then a **Givens rotation** in the  $x_1$ - $x_2$  plane in  $\mathbb{R}^3$  is a matrix  $G_{12}(\theta) \in \mathbb{R}^{3 \times 3}$  that rotates each vector in the  $x_1$ - $x_2$  plane in  $\mathbb{R}^3$  by an angle  $\theta$  in the **anti-clock-wise direction** and  $G_{12}(\theta)e_3 = e_3$ .

Consequently, the rotation  $G_{12}(\theta)$  is given by

$$G_{12}(\theta) = \left[ \begin{array}{cc|c} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ \hline 0 & 0 & 1 \end{array} \right] = \left[ \begin{array}{cc|c} c & -s & 0 \\ s & c & 0 \\ \hline 0 & 0 & 1 \end{array} \right].$$

Similarly, the Givens rotation  $G_{23}(\theta) \in \mathbb{R}^{3 \times 3}$  in the  $x_2$ - $x_3$  plane leaves  $e_1$  unchanged, that is,  $G_{23}(\theta)e_1 = e_1$ , and is given by

$$G_{23}(\theta) = \left[ \begin{array}{c|cc} 1 & 0 & 0 \\ \hline 0 & \cos \theta & -\sin \theta \\ 0 & \sin \theta & \cos \theta \end{array} \right] = \left[ \begin{array}{c|cc} 1 & 0 & 0 \\ \hline 0 & c & -s \\ 0 & s & c \end{array} \right].$$



## Givens rotations in $\mathbb{R}^3$

The Givens rotation  $G_{13}(\theta)$  is a rotation in  $x_1$ - $x_3$  plane. Hence we have

$$G_{13}(\theta)e_1 = \begin{bmatrix} \cos \theta \\ 0 \\ \sin \theta \end{bmatrix} \text{ and } G_{13}(\theta)e_3 = \begin{bmatrix} -\sin \theta \\ 0 \\ \cos \theta \end{bmatrix}.$$

## Givens rotations in $\mathbb{R}^3$

The Givens rotation  $G_{13}(\theta)$  is a rotation in  $x_1$ - $x_3$  plane. Hence we have

$$G_{13}(\theta)e_1 = \begin{bmatrix} \cos \theta \\ 0 \\ \sin \theta \end{bmatrix} \text{ and } G_{13}(\theta)e_3 = \begin{bmatrix} -\sin \theta \\ 0 \\ \cos \theta \end{bmatrix}.$$

Since  $G_{13}(\theta)$  leaves  $e_2$  unchanged, that is,  $G_{13}(\theta)e_2 = e_2$ , we have

## Givens rotations in $\mathbb{R}^3$

The Givens rotation  $G_{13}(\theta)$  is a rotation in  $x_1$ - $x_3$  plane. Hence we have

$$G_{13}(\theta)e_1 = \begin{bmatrix} \cos \theta \\ 0 \\ \sin \theta \end{bmatrix} \text{ and } G_{13}(\theta)e_3 = \begin{bmatrix} -\sin \theta \\ 0 \\ \cos \theta \end{bmatrix}.$$

Since  $G_{13}(\theta)$  leaves  $e_2$  unchanged, that is,  $G_{13}(\theta)e_2 = e_2$ , we have

$$G_{13}(\theta) = \begin{bmatrix} \cos \theta & 0 & -\sin \theta \\ 0 & 1 & 0 \\ \sin \theta & 0 & \cos \theta \end{bmatrix} = \begin{bmatrix} c & 0 & -s \\ 0 & 1 & 0 \\ s & 0 & c \end{bmatrix}.$$

## Givens rotations in $\mathbb{R}^3$

The Givens rotation  $G_{13}(\theta)$  is a rotation in  $x_1$ - $x_3$  plane. Hence we have

$$G_{13}(\theta)e_1 = \begin{bmatrix} \cos \theta \\ 0 \\ \sin \theta \end{bmatrix} \text{ and } G_{13}(\theta)e_3 = \begin{bmatrix} -\sin \theta \\ 0 \\ \cos \theta \end{bmatrix}.$$

Since  $G_{13}(\theta)$  leaves  $e_2$  unchanged, that is,  $G_{13}(\theta)e_2 = e_2$ , we have

$$G_{13}(\theta) = \begin{bmatrix} \cos \theta & 0 & -\sin \theta \\ 0 & 1 & 0 \\ \sin \theta & 0 & \cos \theta \end{bmatrix} = \begin{bmatrix} c & 0 & -s \\ 0 & 1 & 0 \\ s & 0 & c \end{bmatrix}.$$

- It follows that  $G_{13}(\theta)G_{13}(\phi) = G_{13}(\theta + \phi)$  and  $G_{13}(0) = I$  which shows that  $G_{13}(\theta)$  is unitary and  $G_{13}(\theta)^{-1} = G_{13}(-\theta) = G_{13}(\theta)^\top$ .

## Givens rotations in $\mathbb{R}^3$

The Givens rotation  $G_{13}(\theta)$  is a rotation in  $x_1$ - $x_3$  plane. Hence we have

$$G_{13}(\theta)e_1 = \begin{bmatrix} \cos \theta \\ 0 \\ \sin \theta \end{bmatrix} \text{ and } G_{13}(\theta)e_3 = \begin{bmatrix} -\sin \theta \\ 0 \\ \cos \theta \end{bmatrix}.$$

Since  $G_{13}(\theta)$  leaves  $e_2$  unchanged, that is,  $G_{13}(\theta)e_2 = e_2$ , we have

$$G_{13}(\theta) = \begin{bmatrix} \cos \theta & 0 & -\sin \theta \\ 0 & 1 & 0 \\ \sin \theta & 0 & \cos \theta \end{bmatrix} = \begin{bmatrix} c & 0 & -s \\ 0 & 1 & 0 \\ s & 0 & c \end{bmatrix}.$$

- It follows that  $G_{13}(\theta)G_{13}(\phi) = G_{13}(\theta + \phi)$  and  $G_{13}(0) = I$  which shows that  $G_{13}(\theta)$  is unitary and  $G_{13}(\theta)^{-1} = G_{13}(-\theta) = G_{13}(\theta)^\top$ .
- Ditto for the rotations  $G_{12}(\theta)$  and  $G_{23}(\theta)$ .

## Givens rotations in $\mathbb{R}^3$

The Givens rotation  $G_{13}(\theta)$  is a rotation in  $x_1$ - $x_3$  plane. Hence we have

$$G_{13}(\theta)e_1 = \begin{bmatrix} \cos \theta \\ 0 \\ \sin \theta \end{bmatrix} \text{ and } G_{13}(\theta)e_3 = \begin{bmatrix} -\sin \theta \\ 0 \\ \cos \theta \end{bmatrix}.$$

Since  $G_{13}(\theta)$  leaves  $e_2$  unchanged, that is,  $G_{13}(\theta)e_2 = e_2$ , we have

$$G_{13}(\theta) = \begin{bmatrix} \cos \theta & 0 & -\sin \theta \\ 0 & 1 & 0 \\ \sin \theta & 0 & \cos \theta \end{bmatrix} = \begin{bmatrix} c & 0 & -s \\ 0 & 1 & 0 \\ s & 0 & c \end{bmatrix}.$$

- It follows that  $G_{13}(\theta)G_{13}(\phi) = G_{13}(\theta + \phi)$  and  $G_{13}(0) = I$  which shows that  $G_{13}(\theta)$  is unitary and  $G_{13}(\theta)^{-1} = G_{13}(-\theta) = G_{13}(\theta)^\top$ .
- Ditto for the rotations  $G_{12}(\theta)$  and  $G_{23}(\theta)$ .
- There are three Givens rotations in  $\mathbb{R}^3$ , namely,  $G_{12}(\theta)$ ,  $G_{23}(\theta)$  and  $G_{13}(\theta)$ .

# Givens rotations in $\mathbb{R}^n$

**Definition:** A **Givens rotation** in  $\mathbb{R}^n$  is a rotation in the coordinate plane, that is, in the plane **spanned by two coordinate axes**.

# Givens rotations in $\mathbb{R}^n$

**Definition:** A **Givens rotation** in  $\mathbb{R}^n$  is a rotation in the coordinate plane, that is, in the plane **spanned by two coordinate axes**.

Let  $\theta \in [0, 2\pi]$ . Then a **Givens rotation** in the  $x_i$ - $x_j$  plane in  $\mathbb{R}^n$  is a matrix  $G_{ij}(\theta) \in \mathbb{R}^{n \times n}$  that rotates each vector in the  $x_i$ - $x_j$  plane in  $\mathbb{R}^n$  by an angle  $\theta$  in the **anti-clock-wise direction** and  $G_{ij}(\theta)e_k = e_k$  for  $k \notin \{i, j\}$ .



## Givens rotations in $\mathbb{R}^n$

**Definition:** A **Givens rotation** in  $\mathbb{R}^n$  is a rotation in the coordinate plane, that is, in the plane **spanned by two coordinate axes**.

Let  $\theta \in [0, 2\pi]$ . Then a **Givens rotation** in the  $x_i$ - $x_j$  plane in  $\mathbb{R}^n$  is a matrix  $G_{ij}(\theta) \in \mathbb{R}^{n \times n}$  that rotates each vector in the  $x_i$ - $x_j$  plane in  $\mathbb{R}^n$  by an angle  $\theta$  in the **anti-clock-wise direction** and  $G_{ij}(\theta)e_k = e_k$  for  $k \notin \{i, j\}$ .

Thus the Givens rotation  $G_{ij}(\theta)$  differs from the identity matrix  $I$  only in the  $i$ -th and the  $j$ -th columns.

## Givens rotations in $\mathbb{R}^n$

**Definition:** A **Givens rotation** in  $\mathbb{R}^n$  is a rotation in the coordinate plane, that is, in the plane **spanned by two coordinate axes**.

Let  $\theta \in [0, 2\pi]$ . Then a **Givens rotation** in the  $x_i$ - $x_j$  plane in  $\mathbb{R}^n$  is a matrix  $G_{ij}(\theta) \in \mathbb{R}^{n \times n}$  that rotates each vector in the  $x_i$ - $x_j$  plane in  $\mathbb{R}^n$  by an angle  $\theta$  in the **anti-clock-wise direction** and  $G_{ij}(\theta)e_k = e_k$  for  $k \notin \{i, j\}$ .

Thus the Givens rotation  $G_{ij}(\theta)$  differs from the identity matrix  $I$  only in the  $i$ -th and the  $j$ -th columns.

Note that if  $x := [x_1 \ \cdots \ x_n]^\top \in \mathbb{R}^n$  then the map  $x \mapsto G_{ij}(\theta)x$  alters only the components  $x_i$  and  $x_j$  by their linear combinations

## Givens rotations in $\mathbb{R}^n$

**Definition:** A **Givens rotation** in  $\mathbb{R}^n$  is a rotation in the coordinate plane, that is, in the plane **spanned by two coordinate axes**.

Let  $\theta \in [0, 2\pi]$ . Then a **Givens rotation** in the  $x_i$ - $x_j$  plane in  $\mathbb{R}^n$  is a matrix  $G_{ij}(\theta) \in \mathbb{R}^{n \times n}$  that rotates each vector in the  $x_i$ - $x_j$  plane in  $\mathbb{R}^n$  by an angle  $\theta$  in the **anti-clock-wise direction** and  $G_{ij}(\theta)e_k = e_k$  for  $k \notin \{i, j\}$ .

Thus the Givens rotation  $G_{ij}(\theta)$  differs from the identity matrix  $I$  only in the  $i$ -th and the  $j$ -th columns.

Note that if  $x := [x_1 \ \cdots \ x_n]^\top \in \mathbb{R}^n$  then the map  $x \mapsto G_{ij}(\theta)x$  alters only the components  $x_i$  and  $x_j$  by their linear combinations

$$\begin{bmatrix} x_i \\ x_j \end{bmatrix} \longrightarrow \begin{bmatrix} x_i \cos \theta - x_j \sin \theta \\ x_i \sin \theta + x_j \cos \theta \end{bmatrix}$$

and leaves other components of  $x$  unchanged.

# Givens rotations in $\mathbb{R}^n$

**Definition:** A **Givens rotation** in  $\mathbb{R}^n$  is a rotation in the coordinate plane, that is, in the plane **spanned by two coordinate axes**.

Let  $\theta \in [0, 2\pi]$ . Then a **Givens rotation** in the  $x_i$ - $x_j$  plane in  $\mathbb{R}^n$  is a matrix  $G_{ij}(\theta) \in \mathbb{R}^{n \times n}$  that rotates each vector in the  $x_i$ - $x_j$  plane in  $\mathbb{R}^n$  by an angle  $\theta$  in the **anti-clock-wise direction** and  $G_{ij}(\theta)e_k = e_k$  for  $k \notin \{i, j\}$ .

Thus the Givens rotation  $G_{ij}(\theta)$  differs from the identity matrix  $I$  only in the  $i$ -th and the  $j$ -th columns.

Note that if  $x := [x_1 \ \cdots \ x_n]^\top \in \mathbb{R}^n$  then the map  $x \mapsto G_{ij}(\theta)x$  alters only the components  $x_i$  and  $x_j$  by their linear combinations

$$\begin{bmatrix} x_i \\ x_j \end{bmatrix} \longrightarrow \begin{bmatrix} x_i \cos \theta - x_j \sin \theta \\ x_i \sin \theta + x_j \cos \theta \end{bmatrix}$$

and leaves other components of  $x$  unchanged.

Hence  $G_{ij}(\theta)e_i = e_i \cos \theta + e_j \sin \theta$  and  $G_{ij}(\theta)e_j = -e_i \sin \theta + e_j \cos \theta$ .

# Givens rotations in $\mathbb{R}^n$

For simplicity, we denote a Givens rotation in the  $x_i$ - $x_j$  plane in  $\mathbb{R}^n$  by  $G_{ij}$ . For  $i < j$ , we have

$$G_{ij} = \begin{bmatrix} 1 & & & & & \\ & \ddots & & & & \\ & & 1 & & & \\ & & & c & & -s \\ & & & & 1 & \\ & & & & & \ddots \\ & & & s & & c \\ & & & & & & 1 \\ & & & & & & & \ddots \\ & & & & & & & & 1 \end{bmatrix}.$$

## Givens rotations in $\mathbb{R}^n$

For simplicity, we denote a Givens rotation in the  $x_i$ - $x_j$  plane in  $\mathbb{R}^n$  by  $G_{ij}$ . For  $i < j$ , we have

$$G_{ij} = \begin{bmatrix} 1 & & & & & \\ & \ddots & & & & \\ & & 1 & & & \\ & & & c & & -s \\ & & & & 1 & \\ & & & s & & c \\ & & & & & & 1 \\ & & & & & & & \ddots \\ & & & & & & & & 1 \end{bmatrix}.$$

Note that  $G_{ij}$  differs from the identity matrix  $I$  in four entries  $(i, i)$ ,  $(i, j)$ ,  $(j, i)$  and  $(j, j)$ . These entries are  $c$ ,  $-s$ ,  $s$  and  $c$ , respectively.

# Properties of Givens rotations in $\mathbb{R}^n$

- Let  $G_{ij}(\theta)$  and  $G_{ij}(\phi)$  be Givens rotations in  $\mathbb{R}^n$ . Then  $G_{ij}(\theta)G_{ij}(\phi) = G_{ij}(\phi)G_{ij}(\theta) = G_{ij}(\theta + \phi)$  and  $G_{ij}(0) = I$ .

# Properties of Givens rotations in $\mathbb{R}^n$

- Let  $G_{ij}(\theta)$  and  $G_{ij}(\phi)$  be Givens rotations in  $\mathbb{R}^n$ . Then  $G_{ij}(\theta)G_{ij}(\phi) = G_{ij}(\phi)G_{ij}(\theta) = G_{ij}(\theta + \phi)$  and  $G_{ij}(0) = I$ .
- $G_{ij}(\theta)$  is unitary and  $G_{ij}(\theta)^{-1} = G_{ij}(-\theta) = G_{ij}(\theta)^\top$ .



# Properties of Givens rotations in $\mathbb{R}^n$

- Let  $G_{ij}(\theta)$  and  $G_{ij}(\phi)$  be Givens rotations in  $\mathbb{R}^n$ . Then  $G_{ij}(\theta)G_{ij}(\phi) = G_{ij}(\phi)G_{ij}(\theta) = G_{ij}(\theta + \phi)$  and  $G_{ij}(0) = I$ .
- $G_{ij}(\theta)$  is unitary and  $G_{ij}(\theta)^{-1} = G_{ij}(-\theta) = G_{ij}(\theta)^\top$ .
- The map  $x \mapsto G_{ij}(\theta)x$  alters only the  $i$ -th and  $j$ -th components of

$$x \in \mathbb{R}^n \text{ and requires 8 flops as } G_{ij}(\theta) \begin{bmatrix} x_1 \\ \vdots \\ x_i \\ \vdots \\ x_j \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} x_1 \\ \vdots \\ cx_i - sx_j \\ \vdots \\ sx_i + cx_j \\ \vdots \\ x_n \end{bmatrix}.$$

# Properties of Givens rotations in $\mathbb{R}^n$

- Let  $G_{ij}(\theta)$  and  $G_{ij}(\phi)$  be Givens rotations in  $\mathbb{R}^n$ . Then  $G_{ij}(\theta)G_{ij}(\phi) = G_{ij}(\phi)G_{ij}(\theta) = G_{ij}(\theta + \phi)$  and  $G_{ij}(0) = I$ .
- $G_{ij}(\theta)$  is unitary and  $G_{ij}(\theta)^{-1} = G_{ij}(-\theta) = G_{ij}(\theta)^\top$ .
- The map  $x \mapsto G_{ij}(\theta)x$  alters only the  $i$ -th and  $j$ -th components of

$$x \in \mathbb{R}^n \text{ and requires } 8 \text{ flops as } G_{ij}(\theta) \begin{bmatrix} x_1 \\ \vdots \\ x_i \\ \vdots \\ x_j \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} x_1 \\ \vdots \\ cx_i - sx_j \\ \vdots \\ sx_i + cx_j \\ \vdots \\ x_n \end{bmatrix}.$$

4 multiplication ops, c with  $x_i$ ,  $x_j$  & s with  $x_i$ ,  $x_j$   
 2 subtraction ops  
 2 ops to calculate c and s.

- The transformation  $A \mapsto G_{ij}(\theta)A$  alters only the  $i$ -th and  $j$ -th rows of  $A \in \mathbb{R}^{n \times p}$  and requires  $8p$  flops as

$$e_i^\top (G_{ij}(\theta)A) = c(e_i^\top A) - s(e_j^\top A) \text{ and } e_j^\top (G_{ij}(\theta)A) = s(e_i^\top A) + c(e_j^\top A).$$

## Creating zero by Givens rotations

Let  $x := [x_1 \ \cdots \ x_n]^\top \in \mathbb{R}^n$ . We can choose a rotation to transform  $x$  to a vector  $y$  with a specified component zero.

## Creating zero by Givens rotations

Let  $x := [x_1 \ \cdots \ x_n]^\top \in \mathbb{R}^n$ . We can choose a rotation to transform  $x$  to a vector  $y$  with a specified component zero. For example, consider the rotation  $G_{ij}$  in the  $x_i$ - $x_j$  plane with  $c := \frac{x_i}{\sqrt{x_i^2 + x_j^2}}$  and  $s := \frac{x_j}{\sqrt{x_i^2 + x_j^2}}$ .

## Creating zero by Givens rotations

Let  $x := [x_1 \ \cdots \ x_n]^\top \in \mathbb{R}^n$ . We can choose a rotation to transform  $x$  to a vector  $y$  with a specified component zero. For example, consider the rotation  $G_{ij}$  in the  $x_i$ - $x_j$  plane with  $c := \frac{x_i}{\sqrt{x_i^2 + x_j^2}}$  and  $s := \frac{x_j}{\sqrt{x_i^2 + x_j^2}}$ .

Then the  $j$ -th component of  $y := G_{ij}^\top x$  is zero. In fact, we have

$$y_j = 0, y_i = \sqrt{x_i^2 + x_j^2} \text{ and } y_k = x_k \text{ for } k \notin \{i, j\}.$$

## Creating zero by Givens rotations

Let  $x := [x_1 \ \cdots \ x_n]^\top \in \mathbb{R}^n$ . We can choose a rotation to transform  $x$  to a vector  $y$  with a specified component zero. For example, consider the rotation  $G_{ij}$  in the  $x_i$ - $x_j$  plane with  $c := \frac{x_i}{\sqrt{x_i^2 + x_j^2}}$  and  $s := \frac{x_j}{\sqrt{x_i^2 + x_j^2}}$ .

Then the  $j$ -th component of  $y := G_{ij}^\top x$  is zero. In fact, we have

$$y_j = 0, \ y_i = \sqrt{x_i^2 + x_j^2} \text{ and } y_k = x_k \text{ for } k \notin \{i, j\}.$$

Several rotations can be used to transform  $x$  to  $\|x\|_2 e_1$ .

## Creating zero by Givens rotations

Let  $x := [x_1 \ \cdots \ x_n]^\top \in \mathbb{R}^n$ . We can choose a rotation to transform  $x$  to a vector  $y$  with a specified component zero. For example, consider the rotation  $G_{ij}$  in the  $x_i$ - $x_j$  plane with  $c := \frac{x_i}{\sqrt{x_i^2 + x_j^2}}$  and  $s := \frac{x_j}{\sqrt{x_i^2 + x_j^2}}$ .

Then the  $j$ -th component of  $y := G_{ij}^\top x$  is zero. In fact, we have

$$y_j = 0, y_i = \sqrt{x_i^2 + x_j^2} \text{ and } y_k = x_k \text{ for } k \notin \{i, j\}.$$

Several rotations can be used to transform  $x$  to  $\|x\|_2 e_1$ . Choose  $G_{12}$  so that the 2nd component of  $G_{12}^\top x$  is zero.

## Creating zero by Givens rotations

Let  $x := [x_1 \ \cdots \ x_n]^\top \in \mathbb{R}^n$ . We can choose a rotation to transform  $x$  to a vector  $y$  with a specified component zero. For example, consider the rotation  $G_{ij}$  in the  $x_i$ - $x_j$  plane with  $c := \frac{x_i}{\sqrt{x_i^2 + x_j^2}}$  and  $s := \frac{x_j}{\sqrt{x_i^2 + x_j^2}}$ .

Then the  $j$ -th component of  $y := G_{ij}^\top x$  is zero. In fact, we have

$$y_j = 0, y_i = \sqrt{x_i^2 + x_j^2} \text{ and } y_k = x_k \text{ for } k \notin \{i, j\}.$$

Several rotations can be used to transform  $x$  to  $\|x\|_2 e_1$ . Choose  $G_{12}$  so that the 2nd component of  $G_{12}^\top x$  is zero. Then choose  $G_{13}$  so that the 3rd component of  $G_{13}^\top (G_{12}^\top x)$  is zero and so on.



## Creating zero by Givens rotations

Let  $x := [x_1 \ \cdots \ x_n]^\top \in \mathbb{R}^n$ . We can choose a rotation to transform  $x$  to a vector  $y$  with a specified component zero. For example, consider the rotation  $G_{ij}$  in the  $x_i$ - $x_j$  plane with  $c := \frac{x_i}{\sqrt{x_i^2 + x_j^2}}$  and  $s := \frac{x_j}{\sqrt{x_i^2 + x_j^2}}$ .

Then the  $j$ -th component of  $y := G_{ij}^\top x$  is zero. In fact, we have

$$y_j = 0, y_i = \sqrt{x_i^2 + x_j^2} \text{ and } y_k = x_k \text{ for } k \notin \{i, j\}.$$

Several rotations can be used to transform  $x$  to  $\|x\|_2 e_1$ . Choose  $G_{12}$  so that the 2nd component of  $G_{12}^\top x$  is zero. Then choose  $G_{13}$  so that the 3rd component of  $G_{13}^\top (G_{12}^\top x)$  is zero and so on. **Cost:  $8n$  flops.**

## Creating zero by Givens rotations

Let  $x := [x_1 \ \cdots \ x_n]^\top \in \mathbb{R}^n$ . We can choose a rotation to transform  $x$  to a vector  $y$  with a specified component zero. For example, consider the rotation  $G_{ij}$  in the  $x_i$ - $x_j$  plane with  $c := \frac{x_i}{\sqrt{x_i^2 + x_j^2}}$  and  $s := \frac{x_j}{\sqrt{x_i^2 + x_j^2}}$ .

Then the  $j$ -th component of  $y := G_{ij}^\top x$  is zero. In fact, we have

$$y_j = 0, \ y_i = \sqrt{x_i^2 + x_j^2} \text{ and } y_k = x_k \text{ for } k \notin \{i, j\}.$$

Several rotations can be used to transform  $x$  to  $\|x\|_2 e_1$ . Choose  $G_{12}$  so that the 2nd component of  $G_{12}^\top x$  is zero. Then choose  $G_{13}$  so that the 3rd component of  $G_{13}^\top (G_{12}^\top x)$  is zero and so on. **Cost:  $8n$  flops.**

$$G_{12}^\top x = \begin{bmatrix} \sigma_1 \\ 0 \\ x_3 \\ x_4 \\ \vdots \\ x_n \end{bmatrix}$$

## Creating zero by Givens rotations

Let  $x := [x_1 \ \cdots \ x_n]^\top \in \mathbb{R}^n$ . We can choose a rotation to transform  $x$  to a vector  $y$  with a specified component zero. For example, consider the rotation  $G_{ij}$  in the  $x_i$ - $x_j$  plane with  $c := \frac{x_i}{\sqrt{x_i^2 + x_j^2}}$  and  $s := \frac{x_j}{\sqrt{x_i^2 + x_j^2}}$ .

Then the  $j$ -th component of  $y := G_{ij}^\top x$  is zero. In fact, we have

$$y_j = 0, \ y_i = \sqrt{x_i^2 + x_j^2} \text{ and } y_k = x_k \text{ for } k \notin \{i, j\}.$$

Several rotations can be used to transform  $x$  to  $\|x\|_2 e_1$ . Choose  $G_{12}$  so that the 2nd component of  $G_{12}^\top x$  is zero. Then choose  $G_{13}$  so that the 3rd component of  $G_{13}^\top (G_{12}^\top x)$  is zero and so on. **Cost:  $8n$  flops.**

$$G_{12}^\top x = \begin{bmatrix} \sigma_1 \\ 0 \\ x_3 \\ x_4 \\ \vdots \\ x_n \end{bmatrix} \longrightarrow G_{13}^\top G_{12}^\top x = \begin{bmatrix} \sigma_2 \\ 0 \\ 0 \\ x_4 \\ \vdots \\ x_n \end{bmatrix}$$

## Creating zero by Givens rotations

Let  $x := [x_1 \ \cdots \ x_n]^\top \in \mathbb{R}^n$ . We can choose a rotation to transform  $x$  to a vector  $y$  with a specified component zero. For example, consider the rotation  $G_{ij}$  in the  $x_i$ - $x_j$  plane with  $c := \frac{x_i}{\sqrt{x_i^2 + x_j^2}}$  and  $s := \frac{x_j}{\sqrt{x_i^2 + x_j^2}}$ .

Then the  $j$ -th component of  $y := G_{ij}^\top x$  is zero. In fact, we have

$$y_j = 0, \ y_i = \sqrt{x_i^2 + x_j^2} \text{ and } y_k = x_k \text{ for } k \notin \{i, j\}.$$

Several rotations can be used to transform  $x$  to  $\|x\|_2 e_1$ . Choose  $G_{12}$  so that the 2nd component of  $G_{12}^\top x$  is zero. Then choose  $G_{13}$  so that the 3rd component of  $G_{13}^\top (G_{12}^\top x)$  is zero and so on. **Cost:  $8n$  flops.**

$$G_{12}^\top x = \begin{bmatrix} \sigma_1 \\ 0 \\ x_3 \\ x_4 \\ \vdots \\ x_n \end{bmatrix} \longrightarrow G_{13}^\top G_{12}^\top x = \begin{bmatrix} \sigma_2 \\ 0 \\ 0 \\ x_4 \\ \vdots \\ x_n \end{bmatrix} \longrightarrow G_{1n}^\top \cdots G_{12}^\top x = \begin{bmatrix} \sigma_{n-1} \\ 0 \\ 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}.$$

# QR factorization by Givens rotations

We now use rotations for introducing zeros below the diagonal entries of a matrix. For  $i < j$ , we denote a rotation in the  $x_i$ - $x_j$  plane  $G_{ji}$ .

# QR factorization by Givens rotations

We now use rotations for introducing zeros below the diagonal entries of a matrix. For  $i < j$ , we denote a rotation in the  $x_i$ - $x_j$  plane  $G_{ji}$ .

Let  $A \in \mathbb{R}^{n \times n}$ . First, we choose rotations  $G_{21}, \dots, G_{n1}$  that introduce zeros at  $(2,1), (3, 1), \dots, (n, 1)$  entries of  $A$ . Schematically

# QR factorization by Givens rotations

We now use rotations for introducing zeros below the diagonal entries of a matrix. For  $i < j$ , we denote a rotation in the  $x_i$ - $x_j$  plane  $G_{ji}$ .

Let  $A \in \mathbb{R}^{n \times n}$ . First, we choose rotations  $G_{21}, \dots, G_{n1}$  that introduce zeros at  $(2,1), (3,1), \dots, (n,1)$  entries of  $A$ . Schematically

$$\underbrace{\begin{bmatrix} \times & \times & \times \\ \times & \times & \times \\ \times & \times & \times \end{bmatrix}}_A \longrightarrow \underbrace{\begin{bmatrix} \times & \times & \times \\ 0 & \times & \times \\ \times & \times & \times \end{bmatrix}}_{G_{21}^T A} \longrightarrow \underbrace{\begin{bmatrix} \times & \times & \times \\ 0 & \times & \times \\ 0 & \times & \times \end{bmatrix}}_{G_{31}^T G_{21}^T A}.$$

# QR factorization by Givens rotations

We now use rotations for introducing zeros below the diagonal entries of a matrix. For  $i < j$ , we denote a rotation in the  $x_i$ - $x_j$  plane  $G_{ji}$ .

Let  $A \in \mathbb{R}^{n \times n}$ . First, we choose rotations  $G_{21}, \dots, G_{n1}$  that introduce zeros at  $(2,1), (3,1), \dots, (n,1)$  entries of  $A$ . Schematically

$$\underbrace{\begin{bmatrix} \times & \times & \times \\ \times & \times & \times \\ \times & \times & \times \end{bmatrix}}_A \longrightarrow \underbrace{\begin{bmatrix} \times & \times & \times \\ 0 & \times & \times \\ \times & \times & \times \end{bmatrix}}_{G_{21}^T A} \longrightarrow \underbrace{\begin{bmatrix} \times & \times & \times \\ 0 & \times & \times \\ 0 & \times & \times \end{bmatrix}}_{G_{31}^T G_{21}^T A}.$$

Next, choose rotations  $G_{32}, \dots, G_{n2}$  that introduce zeros at  $(3,2), (4,2), \dots, (n,2)$  entries of  $G_{n1}^T \cdots G_{21}^T A$  and so on. Schematically



# QR factorization by Givens rotations

We now use rotations for introducing zeros below the diagonal entries of a matrix. For  $i < j$ , we denote a rotation in the  $x_i$ - $x_j$  plane  $G_{ji}$ .

Let  $A \in \mathbb{R}^{n \times n}$ . First, we choose rotations  $G_{21}, \dots, G_{n1}$  that introduce zeros at  $(2,1), (3,1), \dots, (n,1)$  entries of  $A$ . Schematically

$$\underbrace{\begin{bmatrix} \times & \times & \times \\ \times & \times & \times \\ \times & \times & \times \end{bmatrix}}_A \longrightarrow \underbrace{\begin{bmatrix} \times & \times & \times \\ 0 & \times & \times \\ \times & \times & \times \end{bmatrix}}_{G_{21}^T A} \longrightarrow \underbrace{\begin{bmatrix} \times & \times & \times \\ 0 & \times & \times \\ 0 & \times & \times \end{bmatrix}}_{G_{31}^T G_{21}^T A}.$$

Next, choose rotations  $G_{32}, \dots, G_{n2}$  that introduce zeros at  $(3,2), (4,2), \dots, (n,2)$  entries of  $G_{n1}^T \cdots G_{21}^T A$  and so on. Schematically

$$\underbrace{\begin{bmatrix} \times & \times & \times \\ \times & \times & \times \\ \times & \times & \times \end{bmatrix}}_A \longrightarrow \underbrace{\begin{bmatrix} \times & \times & \times \\ 0 & \times & \times \\ 0 & \times & \times \end{bmatrix}}_{G_{31}^T G_{21}^T A} \longrightarrow \underbrace{\begin{bmatrix} \times & \times & \times \\ 0 & \times & \times \\ 0 & 0 & \times \end{bmatrix}}_{G_{32}^T G_{31}^T G_{21}^T A}.$$

# QR factorization by Givens rotations

**Theorem:** Let  $A \in \mathbb{R}^{m \times n}$  with  $m \geq n$ . Then  $A = QR$ , where  $Q \in \mathbb{R}^{m \times m}$  is unitary and  $R \in \mathbb{R}^{m \times n}$  is upper triangular.

# QR factorization by Givens rotations

**Theorem:** Let  $A \in \mathbb{R}^{m \times n}$  with  $m \geq n$ . Then  $A = QR$ , where  $Q \in \mathbb{R}^{m \times m}$  is unitary and  $R \in \mathbb{R}^{m \times n}$  is upper triangular.

**Proof:** Choose a rotation  $G_{21}$  in the  $x_1$ - $x_2$  plane such that

$$G_{21}^T \begin{bmatrix} a_{11} \\ a_{21} \\ a_{31} \\ \vdots \\ a_{m1} \end{bmatrix} = \begin{bmatrix} \sigma_1 \\ 0 \\ a_{31} \\ \vdots \\ a_{m1} \end{bmatrix}.$$

# QR factorization by Givens rotations

**Theorem:** Let  $A \in \mathbb{R}^{m \times n}$  with  $m \geq n$ . Then  $A = QR$ , where  $Q \in \mathbb{R}^{m \times m}$  is unitary and  $R \in \mathbb{R}^{m \times n}$  is upper triangular.

**Proof:** Choose a rotation  $G_{21}$  in the  $x_1$ - $x_2$  plane such that

$$G_{21}^T \begin{bmatrix} a_{11} \\ a_{21} \\ a_{31} \\ \vdots \\ a_{m1} \end{bmatrix} = \begin{bmatrix} \sigma_1 \\ 0 \\ a_{31} \\ \vdots \\ a_{m1} \end{bmatrix}. \text{ Then } G_{21}^T A = \begin{bmatrix} \sigma_1 & a_{12}^{(1)} & \cdots & a_{1n}^{(1)} \\ 0 & a_{22}^{(1)} & \cdots & a_{2n}^{(1)} \\ a_{31} & a_{32} & \cdots & a_{3n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}.$$

# QR factorization by Givens rotations

**Theorem:** Let  $A \in \mathbb{R}^{m \times n}$  with  $m \geq n$ . Then  $A = QR$ , where  $Q \in \mathbb{R}^{m \times m}$  is unitary and  $R \in \mathbb{R}^{m \times n}$  is upper triangular.

**Proof:** Choose a rotation  $G_{21}$  in the  $x_1$ - $x_2$  plane such that

$$G_{21}^T \begin{bmatrix} a_{11} \\ a_{21} \\ a_{31} \\ \vdots \\ a_{m1} \end{bmatrix} = \begin{bmatrix} \sigma_1 \\ 0 \\ a_{31} \\ \vdots \\ a_{m1} \end{bmatrix}. \text{ Then } G_{21}^T A = \begin{bmatrix} \sigma_1 & a_{12}^{(1)} & \cdots & a_{1n}^{(1)} \\ 0 & a_{22}^{(1)} & \cdots & a_{2n}^{(1)} \\ a_{31} & a_{32} & \cdots & a_{3n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}.$$

Next, choose a rotation  $G_{31}$  that creates a zero at (3, 1) entry of  $G_{21}^T A$ . Since  $G_{31}$  is a rotation in  $x_1$ - $x_3$  plane, it leaves the zero at (2, 1) entry of  $G_{21}^T A$  unchanged.

# QR factorization by Givens rotations

**Theorem:** Let  $A \in \mathbb{R}^{m \times n}$  with  $m \geq n$ . Then  $A = QR$ , where  $Q \in \mathbb{R}^{m \times m}$  is unitary and  $R \in \mathbb{R}^{m \times n}$  is upper triangular.

**Proof:** Choose a rotation  $G_{21}$  in the  $x_1$ - $x_2$  plane such that

$$G_{21}^T \begin{bmatrix} a_{11} \\ a_{21} \\ a_{31} \\ \vdots \\ a_{m1} \end{bmatrix} = \begin{bmatrix} \sigma_1 \\ 0 \\ a_{31} \\ \vdots \\ a_{m1} \end{bmatrix}. \text{ Then } G_{21}^T A = \begin{bmatrix} \sigma_1 & a_{12}^{(1)} & \cdots & a_{1n}^{(1)} \\ 0 & a_{22}^{(1)} & \cdots & a_{2n}^{(1)} \\ a_{31} & a_{32} & \cdots & a_{3n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}.$$

Next, choose a rotation  $G_{31}$  that creates a zero at (3, 1) entry of  $G_{21}^T A$ . Since  $G_{31}$  is a rotation in  $x_1$ - $x_3$  plane, it leaves the zero at (2, 1) entry of  $G_{21}^T A$  unchanged. Hence  $G_{31}^T G_{21}^T A$  has zeros at (2, 1) and (3, 1) entries.

# QR factorization by Givens rotations

**Theorem:** Let  $A \in \mathbb{R}^{m \times n}$  with  $m \geq n$ . Then  $A = QR$ , where  $Q \in \mathbb{R}^{m \times m}$  is unitary and  $R \in \mathbb{R}^{m \times n}$  is upper triangular.

**Proof:** Choose a rotation  $G_{21}$  in the  $x_1$ - $x_2$  plane such that

$$G_{21}^T \begin{bmatrix} a_{11} \\ a_{21} \\ a_{31} \\ \vdots \\ a_{m1} \end{bmatrix} = \begin{bmatrix} \sigma_1 \\ 0 \\ a_{31} \\ \vdots \\ a_{m1} \end{bmatrix}. \text{ Then } G_{21}^T A = \begin{bmatrix} \sigma_1 & a_{12}^{(1)} & \cdots & a_{1n}^{(1)} \\ 0 & a_{22}^{(1)} & \cdots & a_{2n}^{(1)} \\ a_{31} & a_{32} & \cdots & a_{3n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}.$$

Next, choose a rotation  $G_{31}$  that creates a zero at (3, 1) entry of  $G_{21}^T A$ . Since  $G_{31}$  is a rotation in  $x_1$ - $x_3$  plane, it leaves the zero at (2, 1) entry of  $G_{21}^T A$  unchanged. Hence  $G_{31}^T G_{21}^T A$  has zeros at (2, 1) and (3, 1) entries.

Continuing in this manner, choose rotations  $G_{41}, \dots, G_{m1}$  such that  $G_{m1}^T \cdots G_{21}^T A$  has zeros in the first column at (2, 1),  $\dots$ , (m, 1) entries.

## QR factorization by Givens rotations

Similarly, choose rotations  $G_{32}, \dots, G_{m2}$  that create zeros at  $(3, 2), \dots, (m, 2)$  entries of  $G_{m1}^\top \cdots G_{21}^\top A$ .



## QR factorization by Givens rotations

Similarly, choose rotations  $G_{32}, \dots, G_{m2}$  that create zeros at  $(3, 2), \dots, (m, 2)$  entries of  $G_{m1}^\top \cdots G_{21}^\top A$ . Then  $G_{m2}^\top \cdots G_{32}^\top (G_{m1}^\top \cdots G_{21}^\top A)$  has zeros in the first and second columns below the diagonals.

## QR factorization by Givens rotations

Similarly, choose rotations  $G_{32}, \dots, G_{m2}$  that create zeros at  $(3, 2), \dots, (m, 2)$  entries of  $G_{m1}^\top \cdots G_{21}^\top A$ . Then  $G_{m2}^\top \cdots G_{32}^\top (G_{m1}^\top \cdots G_{21}^\top A)$  has zeros in the first and second columns below the diagonals.

Continuing in this manner, choose rotations that introduce zeros in the 3rd, 4th,  $\dots$ ,  $n$ th columns below the diagonals.

## QR factorization by Givens rotations

Similarly, choose rotations  $G_{32}, \dots, G_{m2}$  that create zeros at  $(3, 2), \dots, (m, 2)$  entries of  $G_{m1}^\top \cdots G_{21}^\top A$ . Then  $G_{m2}^\top \cdots G_{32}^\top (G_{m1}^\top \cdots G_{21}^\top A)$  has zeros in the first and second columns below the diagonals.

Continuing in this manner, choose rotations that introduce zeros in the 3rd, 4th,  $\dots$ ,  $n$ th columns below the diagonals. Thus we create rotations  $G_{21}, \dots, G_{m1}, G_{32}, \dots, G_{m2}, \dots, G_{n+1,n}, \dots, G_{mn}$  such that

$$R = G_{mn}^\top G_{m-1,n}^\top \cdots G_{n+1,n}^\top \cdots G_{m1}^\top \cdots G_{21}^\top A$$

is upper triangular.

## QR factorization by Givens rotations

Similarly, choose rotations  $G_{32}, \dots, G_{m2}$  that create zeros at  $(3, 2), \dots, (m, 2)$  entries of  $G_{m1}^\top \cdots G_{21}^\top A$ . Then  $G_{m2}^\top \cdots G_{32}^\top (G_{m1}^\top \cdots G_{21}^\top A)$  has zeros in the first and second columns below the diagonals.

Continuing in this manner, choose rotations that introduce zeros in the 3rd, 4th,  $\dots$ ,  $n$ th columns below the diagonals. Thus we create rotations  $G_{21}, \dots, G_{m1}, G_{32}, \dots, G_{m2}, \dots, G_{n+1,n}, \dots, G_{mn}$  such that

$$R = G_{mn}^\top G_{m-1,n}^\top \cdots G_{n+1,n}^\top \cdots G_{m1}^\top \cdots G_{21}^\top A$$

is upper triangular. Set  $Q := G_{21} \cdots G_{m1} G_{32}, \dots, G_{m2} \cdots G_{n+1,n} \cdots, G_{mn}$ . Then  $Q$  is unitary and  $A = QR$ . ■

## QR factorization by Givens rotations

Similarly, choose rotations  $G_{32}, \dots, G_{m2}$  that create zeros at  $(3, 2), \dots, (m, 2)$  entries of  $G_{m1}^\top \cdots G_{21}^\top A$ . Then  $G_{m2}^\top \cdots G_{32}^\top (G_{m1}^\top \cdots G_{21}^\top A)$  has zeros in the first and second columns below the diagonals.

Continuing in this manner, choose rotations that introduce zeros in the 3rd, 4th,  $\dots$ ,  $n$ th columns below the diagonals. Thus we create rotations  $G_{21}, \dots, G_{m1}, G_{32}, \dots, G_{m2}, \dots, G_{n+1,n}, \dots, G_{mn}$  such that

$$R = G_{mn}^\top G_{m-1,n}^\top \cdots G_{n+1,n}^\top \cdots G_{m1}^\top \cdots G_{21}^\top A$$

is upper triangular. Set  $Q := G_{21} \cdots G_{m1} G_{32}, \dots, G_{m2} \cdots G_{n+1,n} \cdots, G_{mn}$ . Then  $Q$  is unitary and  $A = QR$ . ■

**Cost:** At the 1st step, the cost of applying  $m - 1$  rotators  $G_{21}^\top, \dots, G_{m1}^\top$  is  $8(m - 1)n$ .

## QR factorization by Givens rotations

Similarly, choose rotations  $G_{32}, \dots, G_{m2}$  that create zeros at  $(3, 2), \dots, (m, 2)$  entries of  $G_{m1}^\top \cdots G_{21}^\top A$ . Then  $G_{m2}^\top \cdots G_{32}^\top (G_{m1}^\top \cdots G_{21}^\top A)$  has zeros in the first and second columns below the diagonals.

Continuing in this manner, choose rotations that introduce zeros in the 3rd, 4th,  $\dots$ ,  $n$ th columns below the diagonals. Thus we create rotations  $G_{21}, \dots, G_{m1}, G_{32}, \dots, G_{m2}, \dots, G_{n+1,n}, \dots, G_{mn}$  such that

$$R = G_{mn}^\top G_{m-1,n}^\top \cdots G_{n+1,n}^\top \cdots G_{m1}^\top \cdots G_{21}^\top A$$

is upper triangular. Set  $Q := G_{21} \cdots G_{m1} G_{32}, \dots, G_{m2} \cdots G_{n+1,n} \cdots, G_{mn}$ . Then  $Q$  is unitary and  $A = QR$ . ■

**Cost:** At the 1st step, the cost of applying  $m - 1$  rotators  $G_{21}^\top, \dots, G_{m1}^\top$  is  $8(m - 1)n$ . At the 2nd step, the cost is  $8(m - 2)(n - 1)$  and, at the  $j$ -th step, the cost is  $8(m - j)(n - j + 1)$  flops.

## QR factorization by Givens rotations

Similarly, choose rotations  $G_{32}, \dots, G_{m2}$  that create zeros at  $(3, 2), \dots, (m, 2)$  entries of  $G_{m1}^\top \cdots G_{21}^\top A$ . Then  $G_{m2}^\top \cdots G_{32}^\top (G_{m1}^\top \cdots G_{21}^\top A)$  has zeros in the first and second columns below the diagonals.

Continuing in this manner, choose rotations that introduce zeros in the 3rd, 4th,  $\dots$ ,  $n$ th columns below the diagonals. Thus we create rotations  $G_{21}, \dots, G_{m1}, G_{32}, \dots, G_{m2}, \dots, G_{n+1,n}, \dots, G_{mn}$  such that

$$R = G_{mn}^\top G_{m-1,n}^\top \cdots G_{n+1,n}^\top \cdots G_{m1}^\top \cdots G_{21}^\top A$$

is upper triangular. Set  $Q := G_{21} \cdots G_{m1} G_{32}, \dots, G_{m2} \cdots G_{n+1,n} \cdots, G_{mn}$ . Then  $Q$  is unitary and  $A = QR$ . ■

**Cost:** At the 1st step, the cost of applying  $m - 1$  rotators  $G_{21}^\top, \dots, G_{m1}^\top$  is  $8(m - 1)n$ . At the 2nd step, the cost is  $8(m - 2)(n - 1)$  and, at the  $j$ -th step, the cost is  $8(m - j)(n - j + 1)$  flops. Hence the total cost is

$$\sum_{j=1}^n 8(m - j)(n - j + 1) = 8 \int_0^n (m - x)(n - x) dx = 4(mn^2 - \frac{n^3}{3}).$$

## Stable generation of rotations

A naive method to generate a rotation such that  $\begin{bmatrix} c & -s \\ s & c \end{bmatrix}^T \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} \sigma \\ 0 \end{bmatrix}$  is to define  $c := \frac{x_1}{\sqrt{x_1^2 + x_2^2}}$  and  $s := \frac{x_2}{\sqrt{x_1^2 + x_2^2}}$ .



## Stable generation of rotations

A naive method to generate a rotation such that  $\begin{bmatrix} c & -s \\ s & c \end{bmatrix}^T \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} \sigma \\ 0 \end{bmatrix}$  is to define  $c := \frac{x_1}{\sqrt{x_1^2 + x_2^2}}$  and  $s := \frac{x_2}{\sqrt{x_1^2 + x_2^2}}$ . This method may not be numerically stable. A stable method is given by the following.

## Stable generation of rotations

A naive method to generate a rotation such that  $\begin{bmatrix} c & -s \\ s & c \end{bmatrix}^T \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} \sigma \\ 0 \end{bmatrix}$  is to define  $c := \frac{x_1}{\sqrt{x_1^2 + x_2^2}}$  and  $s := \frac{x_2}{\sqrt{x_1^2 + x_2^2}}$ . This method may not be numerically stable. A stable method is given by the following.

```
function [c,s] = Rotation(x1,x2);  
% Input:  x1,x2 scalars  
% Output: c,s such that  $c^2 + s^2 = 1$   
% and  $-s * x1 + c * x2 = 0$ .  
    if x2 == 0 c = 1; s = 0;  
    else  
        if abs(x2)>=abs(x1) k = x1/x2; % computes  $\cot(\theta)$   
        s = 1/sqrt(1+k^2); c = s*k;  
        else  
            t = x2/x1; % computes  $\tan(\theta)$   
            c = 1/sqrt(1+ t^2); s = c*t;  
        end  
    end  
end
```

## Example

Let  $A := \begin{bmatrix} 4 & 5 & 8 \\ 6 & 7 & 9 \\ 3 & 6 & 4 \end{bmatrix}$ . We use rotation to compute QR factorization of  $A$ .

## Example

Let  $A := \begin{bmatrix} 4 & 5 & 8 \\ 6 & 7 & 9 \\ 3 & 6 & 4 \end{bmatrix}$ . We use rotation to compute QR factorization of  $A$ .

The rotation  $G_{21}$  with  $c := 0.5547$  and  $s := 0.8321$  gives

$$G_{21}^T A = \begin{bmatrix} 7.2111 & 8.5979 & 11.9261 \\ 0 & -0.2774 & -1.6641 \\ 3.0000 & 6.0000 & 4.0000 \end{bmatrix}.$$

## Example

Let  $A := \begin{bmatrix} 4 & 5 & 8 \\ 6 & 7 & 9 \\ 3 & 6 & 4 \end{bmatrix}$ . We use rotation to compute QR factorization of  $A$ .

The rotation  $G_{21}$  with  $c := 0.5547$  and  $s := 0.8321$  gives

$$G_{21}^T A = \begin{bmatrix} 7.2111 & 8.5979 & 11.9261 \\ 0 & -0.2774 & -1.6641 \\ 3.0000 & 6.0000 & 4.0000 \end{bmatrix}.$$

The rotation  $G_{31}$  with  $c := 0.9233$  and  $s := 0.3841$  gives

$$G_{31}^T G_{21}^T A = \begin{bmatrix} 7.8102 & 10.2430 & 12.5476 \\ 0 & -0.2774 & -1.6641 \\ 0 & 2.2372 & -0.8878 \end{bmatrix}.$$

## Example

Let  $A := \begin{bmatrix} 4 & 5 & 8 \\ 6 & 7 & 9 \\ 3 & 6 & 4 \end{bmatrix}$ . We use rotation to compute QR factorization of  $A$ .

The rotation  $G_{21}$  with  $c := 0.5547$  and  $s := 0.8321$  gives

$$G_{21}^T A = \begin{bmatrix} 7.2111 & 8.5979 & 11.9261 \\ 0 & -0.2774 & -1.6641 \\ 3.0000 & 6.0000 & 4.0000 \end{bmatrix}.$$

The rotation  $G_{31}$  with  $c := 0.9233$  and  $s := 0.3841$  gives

$$G_{31}^T G_{21}^T A = \begin{bmatrix} 7.8102 & 10.2430 & 12.5476 \\ 0 & -0.2774 & -1.6641 \\ 0 & 2.2372 & -0.8878 \end{bmatrix}.$$

Finally, the rotation  $G_{32}$  with  $c := -0.1230$  and  $s := 0.9924$  gives

$$G_{32}^T G_{31}^T G_{21}^T A = \begin{bmatrix} 7.8102 & 10.2430 & 12.5476 \\ 0 & 2.2543 & -0.6763 \\ 0 & 0 & 1.7607 \end{bmatrix}.$$