

# The PageRank Eigenvalue Problem

Rafikul Alam  
Department of Mathematics  
Indian Institute of Technology Guwahati  
Guwahati - 781039, INDIA

# Outline

- The PageRank Problem
- PageRank as an eigenvalue problem
- Power method for PageRank vector
- Single vector iterations for eigenvalue problems

# The PageRank problem

The task of **extracting information from all the web pages** available on the Internet is performed by **search engines**.

# The PageRank problem

The task of **extracting information from all the web pages** available on the Internet is performed by **search engines**.

**Problem:** If it is found, after a query, that more than a thousand web pages contain information that might be relevant to the user's query, then *in which order should the pages be displayed?*

# The PageRank problem

The task of **extracting information from all the web pages** available on the Internet is performed by **search engines**.

**Problem:** If it is found, after a query, that more than a thousand web pages contain information that might be relevant to the user's query, then *in which order should the pages be displayed?*

Part of the magic behind **Google search engine** is its **PageRank** algorithm, which **ranks all the web pages with public access** independently of user's query.

# The PageRank problem

The task of **extracting information from all the web pages** available on the Internet is performed by **search engines**.

**Problem:** If it is found, after a query, that more than a thousand web pages contain information that might be relevant to the user's query, then *in which order should the pages be displayed?*

Part of the magic behind **Google search engine** is its **PageRank** algorithm, which **ranks all the web pages with public access** independently of user's query.

The PageRank algorithm ranks web pages by solving the **PageRank eigenvalue problem**:

$$\mathbf{G}\mathbf{v} := \underbrace{(\alpha\mathbf{S} + (1 - \alpha)\mathbf{E})}_{\text{Google matrix}} \mathbf{v} = \mathbf{v}.$$

The page with **highest rank** is displayed first. This is amazingly effective!

# Non-negative matrices and vectors

Let  $\mathbf{x} \in \mathbb{R}^n$ . Then  $\mathbf{x}$  is called

- **non-negative** ( $\mathbf{x} \succeq \mathbf{0}$ ) if  $x_j \geq 0$  for  $j = 1 : n$ .
- **positive** ( $\mathbf{x} \succ \mathbf{0}$ ) if  $x_j > 0$  for  $j = 1 : n$ .

# Non-negative matrices and vectors

Let  $\mathbf{x} \in \mathbb{R}^n$ . Then  $\mathbf{x}$  is called

- **non-negative** ( $\mathbf{x} \succeq \mathbf{0}$ ) if  $x_j \geq 0$  for  $j = 1 : n$ .
- **positive** ( $\mathbf{x} \succ \mathbf{0}$ ) if  $x_j > 0$  for  $j = 1 : n$ .

Let  $A \in \mathbb{R}^{m \times n}$ . Then  $A$  is called

- **non-negative** ( $A \succeq \mathbf{0}$ ) if  $A\mathbf{e}_j \succeq \mathbf{0}$  for  $j = 1 : n$ .
- **positive** ( $A \succ \mathbf{0}$ ) if  $A\mathbf{e}_j \succ \mathbf{0}$  for  $j = 1 : n$ .



# Non-negative matrices and vectors

Let  $\mathbf{x} \in \mathbb{R}^n$ . Then  $\mathbf{x}$  is called

- **non-negative** ( $\mathbf{x} \succeq \mathbf{0}$ ) if  $x_j \geq 0$  for  $j = 1 : n$ .
- **positive** ( $\mathbf{x} \succ \mathbf{0}$ ) if  $x_j > 0$  for  $j = 1 : n$ .

Let  $A \in \mathbb{R}^{m \times n}$ . Then  $A$  is called

- **non-negative** ( $A \succeq \mathbf{0}$ ) if  $A\mathbf{e}_j \succeq \mathbf{0}$  for  $j = 1 : n$ .
- **positive** ( $A \succ \mathbf{0}$ ) if  $A\mathbf{e}_j \succ \mathbf{0}$  for  $j = 1 : n$ .

**Fact:** If  $A \succ \mathbf{0}$  and  $\mathbf{x} \succeq \mathbf{0}$  is nonzero then  $A\mathbf{x} \succ \mathbf{0}$ .

# Non-negative matrices and vectors

Let  $\mathbf{x} \in \mathbb{R}^n$ . Then  $\mathbf{x}$  is called

- **non-negative** ( $\mathbf{x} \succeq \mathbf{0}$ ) if  $x_j \geq 0$  for  $j = 1 : n$ .
- **positive** ( $\mathbf{x} \succ \mathbf{0}$ ) if  $x_j > 0$  for  $j = 1 : n$ .

Let  $A \in \mathbb{R}^{m \times n}$ . Then  $A$  is called

- **non-negative** ( $A \succeq \mathbf{0}$ ) if  $A\mathbf{e}_j \succeq \mathbf{0}$  for  $j = 1 : n$ .
- **positive** ( $A \succ \mathbf{0}$ ) if  $A\mathbf{e}_j \succ \mathbf{0}$  for  $j = 1 : n$ .

**Fact:** If  $A \succ \mathbf{0}$  and  $\mathbf{x} \succeq \mathbf{0}$  is nonzero then  $A\mathbf{x} \succ \mathbf{0}$ .

Let  $\mathbf{e} = [1 \ \cdots \ 1]^\top$  and  $A \in \mathbb{R}^{m \times n}$  be non-negative. Then  $A$  is called

- **column stochastic** if  $\mathbf{e}^\top A = \mathbf{e}^\top$ , that is,  $\mathbf{e}^\top A\mathbf{e}_j = 1$  for  $j = 1 : n$ .
- **column sub-stochastic** if  $\mathbf{e}^\top A \preceq \mathbf{e}^\top$ , that is,  $\mathbf{e}^\top A\mathbf{e}_j \leq 1$  for  $j = 1 : n$ .

## Positive eigenpair of a positive matrix

The number  $\rho(A) := \max\{|\lambda| : \lambda \in \Lambda(A)\}$  is called the **spectral radius** of  $A$ .

## Positive eigenpair of a positive matrix

The number  $\rho(A) := \max\{|\lambda| : \lambda \in \Lambda(A)\}$  is called the **spectral radius** of  $A$ . Note that  $\rho(A)$  may or may not be an eigenvalue of  $A$ .

## Positive eigenpair of a positive matrix

The number  $\rho(A) := \max\{|\lambda| : \lambda \in \Lambda(A)\}$  is called the **spectral radius** of  $A$ . Note that  $\rho(A)$  may or may not be an eigenvalue of  $A$ . If  $\lambda$  is called a **dominant eigenvalue** of  $A$  if  $|\lambda| = \rho(A)$ .

## Positive eigenpair of a positive matrix

The number  $\rho(A) := \max\{|\lambda| : \lambda \in \Lambda(A)\}$  is called the **spectral radius** of  $A$ . Note that  $\rho(A)$  may or may not be an eigenvalue of  $A$ . If  $\lambda$  is called a **dominant eigenvalue** of  $A$  if  $|\lambda| = \rho(A)$ .

For  $\mathbf{x} := [x_1 \ \cdots \ x_n]^\top$ , define  $|\mathbf{x}| := [|x_1| \ \cdots \ |x_n|]^\top$ . Similarly, for  $A = [a_{ij}]$ , define  $|A| := [|a_{ij}|]$ . Then  $|A\mathbf{x}| \leq |A| |\mathbf{x}|$ .

## Positive eigenpair of a positive matrix

The number  $\rho(A) := \max\{|\lambda| : \lambda \in \Lambda(A)\}$  is called the **spectral radius** of  $A$ . Note that  $\rho(A)$  may or may not be an eigenvalue of  $A$ . If  $\lambda$  is called a **dominant eigenvalue** of  $A$  if  $|\lambda| = \rho(A)$ .

For  $\mathbf{x} := [x_1 \ \cdots \ x_n]^\top$ , define  $|\mathbf{x}| := [|x_1| \ \cdots \ |x_n|]^\top$ . Similarly, for  $A = [a_{ij}]$ , define  $|A| := [|a_{ij}|]$ . Then  $|A\mathbf{x}| \leq |A| |\mathbf{x}|$ .

**Fact:** If  $A \succ \mathbf{0}$  then the following hold:

- $\rho(A) \in \Lambda(A)$  and  $\rho(A) > 0$
- If  $A\mathbf{v} = \rho(A)\mathbf{v}$  then  $A|\mathbf{v}| = \rho(A)|\mathbf{v}|$  and  $|\mathbf{v}| > \mathbf{0}$ .

## Positive eigenpair of a positive matrix

The number  $\rho(A) := \max\{|\lambda| : \lambda \in \Lambda(A)\}$  is called the **spectral radius** of  $A$ . Note that  $\rho(A)$  may or may not be an eigenvalue of  $A$ . If  $\lambda$  is called a **dominant eigenvalue** of  $A$  if  $|\lambda| = \rho(A)$ .

For  $\mathbf{x} := [x_1 \ \cdots \ x_n]^\top$ , define  $|\mathbf{x}| := [|x_1| \ \cdots \ |x_n|]^\top$ . Similarly, for  $A = [a_{ij}]$ , define  $|A| := [|a_{ij}|]$ . Then  $|A\mathbf{x}| \leq |A| |\mathbf{x}|$ .

**Fact:** If  $A \succ \mathbf{0}$  then the following hold:

- $\rho(A) \in \Lambda(A)$  and  $\rho(A) > 0$
- If  $A\mathbf{v} = \rho(A)\mathbf{v}$  then  $A|\mathbf{v}| = \rho(A)|\mathbf{v}|$  and  $|\mathbf{v}| > \mathbf{0}$ .

**Proof:** If  $A\mathbf{x} = \lambda\mathbf{x}$  and  $|\lambda| = \rho(A)$  then  $\rho(A)|\mathbf{x}| = |A\mathbf{x}| \leq |A||\mathbf{x}|$ .



## Positive eigenpair of a positive matrix

The number  $\rho(A) := \max\{|\lambda| : \lambda \in \Lambda(A)\}$  is called the **spectral radius** of  $A$ . Note that  $\rho(A)$  may or may not be an eigenvalue of  $A$ . If  $\lambda$  is called a **dominant eigenvalue** of  $A$  if  $|\lambda| = \rho(A)$ .

For  $\mathbf{x} := [x_1 \ \cdots \ x_n]^\top$ , define  $|\mathbf{x}| := [|x_1| \ \cdots \ |x_n|]^\top$ . Similarly, for  $A = [a_{ij}]$ , define  $|A| := [|a_{ij}|]$ . Then  $|A\mathbf{x}| \leq |A| |\mathbf{x}|$ .

**Fact:** If  $A \succ \mathbf{0}$  then the following hold:

- $\rho(A) \in \Lambda(A)$  and  $\rho(A) > 0$
- If  $A\mathbf{v} = \rho(A)\mathbf{v}$  then  $A|\mathbf{v}| = \rho(A)|\mathbf{v}|$  and  $|\mathbf{v}| > \mathbf{0}$ .

**Proof:** If  $A\mathbf{x} = \lambda\mathbf{x}$  and  $|\lambda| = \rho(A)$  then  $\rho(A)|\mathbf{x}| = |A\mathbf{x}| \leq |A||\mathbf{x}|$ . Set  $\mathbf{y} := |A||\mathbf{x}| - \rho(A)|\mathbf{x}|$ . Then  $\mathbf{y} \succeq \mathbf{0}$ . Now  $\mathbf{y} \neq \mathbf{0}$  leads to a contradiction.

## Positive eigenpair of a positive matrix

The number  $\rho(A) := \max\{|\lambda| : \lambda \in \Lambda(A)\}$  is called the **spectral radius** of  $A$ . Note that  $\rho(A)$  may or may not be an eigenvalue of  $A$ . If  $\lambda$  is called a **dominant eigenvalue** of  $A$  if  $|\lambda| = \rho(A)$ .

For  $\mathbf{x} := [x_1 \ \cdots \ x_n]^\top$ , define  $|\mathbf{x}| := [|x_1| \ \cdots \ |x_n|]^\top$ . Similarly, for  $A = [a_{ij}]$ , define  $|A| := [|a_{ij}|]$ . Then  $|A\mathbf{x}| \leq |A| |\mathbf{x}|$ .

**Fact:** If  $A \succ \mathbf{0}$  then the following hold:

- $\rho(A) \in \Lambda(A)$  and  $\rho(A) > 0$
- If  $A\mathbf{v} = \rho(A)\mathbf{v}$  then  $A|\mathbf{v}| = \rho(A)|\mathbf{v}|$  and  $|\mathbf{v}| > \mathbf{0}$ .

**Proof:** If  $A\mathbf{x} = \lambda\mathbf{x}$  and  $|\lambda| = \rho(A)$  then  $\rho(A)|\mathbf{x}| = |A\mathbf{x}| \leq |A||\mathbf{x}|$ . Set  $\mathbf{y} := |A||\mathbf{x}| - \rho(A)|\mathbf{x}|$ . Then  $\mathbf{y} \succeq \mathbf{0}$ . Now  $\mathbf{y} \neq \mathbf{0}$  leads to a contradiction.

Hence  $\mathbf{y} = \mathbf{0} \implies |A||\mathbf{x}| = \rho(A)|\mathbf{x}| \implies \rho(A) > 0$  and  $|\mathbf{x}| > \mathbf{0}$ . ■

## Positive eigenpair of a positive matrix

The number  $\rho(A) := \max\{|\lambda| : \lambda \in \Lambda(A)\}$  is called the **spectral radius** of  $A$ . Note that  $\rho(A)$  may or may not be an eigenvalue of  $A$ . ■  $\lambda$  is called a **dominant eigenvalue** of  $A$  if  $|\lambda| = \rho(A)$ .

For  $\mathbf{x} := [x_1 \ \cdots \ x_n]^T$ , define  $|\mathbf{x}| := [|x_1| \ \cdots \ |x_n|]^T$ . Similarly, for  $A = [a_{ij}]$ , define  $|A| := [|a_{ij}|]$ . Then  $|A\mathbf{x}| \leq |A| |\mathbf{x}|$ .

**Fact:** If  $A \succ \mathbf{0}$  then the following hold:

- $\rho(A) \in \Lambda(A)$  and  $\rho(A) > 0$
- If  $A\mathbf{v} = \rho(A)\mathbf{v}$  then  $A|\mathbf{v}| = \rho(A)|\mathbf{v}|$  and  $|\mathbf{v}| > \mathbf{0}$ .

**Proof:** If  $A\mathbf{x} = \lambda\mathbf{x}$  and  $|\lambda| = \rho(A)$  then  $\rho(A)|\mathbf{x}| = |A\mathbf{x}| \leq |A||\mathbf{x}|$ . Set  $\mathbf{y} := |A||\mathbf{x}| - \rho(A)|\mathbf{x}|$ . Then  $\mathbf{y} \succeq \mathbf{0}$ . Now  $\mathbf{y} \neq \mathbf{0}$  leads to a contradiction.

Hence  $\mathbf{y} = \mathbf{0} \implies |A||\mathbf{x}| = \rho(A)|\mathbf{x}| \implies \rho(A) > 0$  and  $|\mathbf{x}| > \mathbf{0}$ . ■

**Fact:** If  $A \succ \mathbf{0}$  then  $\rho(A)$  is a simple eigenvalue of  $A$ . Thus

$$\text{algebraic multiplicity of } \rho(A) = \text{geometric multiplicity of } \rho(A) = 1.$$

## Perron vector and Perron root

Thus  $N(A - \rho(A)I) = \text{span}(\mathbf{v})$  and either  $\mathbf{v} \succ \mathbf{0}$  or  $-\mathbf{v} \succ \mathbf{0}$ .

# Perron vector and Perron root

Thus  $N(A - \rho(A)I) = \text{span}(\mathbf{v})$  and either  $\mathbf{v} \succ \mathbf{0}$  or  $-\mathbf{v} \succ \mathbf{0}$ . Consequently, there is a unique vector  $\mathbf{p} \succ \mathbf{0}$  such that

$$A\mathbf{p} = \rho(A)\mathbf{p} \quad \text{and} \quad \mathbf{e}^\top \mathbf{p} = p_1 + \cdots + p_n = 1.$$

# Perron vector and Perron root

Thus  $N(A - \rho(A)I) = \text{span}(\mathbf{v})$  and either  $\mathbf{v} \succ \mathbf{0}$  or  $-\mathbf{v} \succ \mathbf{0}$ . Consequently, there is a unique vector  $\mathbf{p} \succ \mathbf{0}$  such that

$$A\mathbf{p} = \rho(A)\mathbf{p} \quad \text{and} \quad \mathbf{e}^\top \mathbf{p} = p_1 + \cdots + p_n = 1.$$

**Definition:** The special vector  $\mathbf{p}$  is called the **Perron vector** of  $A \succ \mathbf{0}$  and the associated eigenvalue  $\rho(A)$  is called the **Perron root** of  $A$ .

# Perron vector and Perron root

Thus  $N(A - \rho(A)I) = \text{span}(\mathbf{v})$  and either  $\mathbf{v} \succ \mathbf{0}$  or  $-\mathbf{v} \succ \mathbf{0}$ . Consequently, there is a unique vector  $\mathbf{p} \succ \mathbf{0}$  such that

$$A\mathbf{p} = \rho(A)\mathbf{p} \quad \text{and} \quad \mathbf{e}^\top \mathbf{p} = p_1 + \cdots + p_n = 1.$$

**Definition:** The special vector  $\mathbf{p}$  is called the **Perron vector** of  $A \succ \mathbf{0}$  and the associated eigenvalue  $\rho(A)$  is called the **Perron root** of  $A$ .

**Fact:** If  $A \succ \mathbf{0}$  then there are no nonnegative eigenvectors of  $A$  other than the Perron vector  $\mathbf{p}$  and its positive scalar multiples.

# Perron vector and Perron root

Thus  $N(A - \rho(A)I) = \text{span}(\mathbf{v})$  and either  $\mathbf{v} \succ \mathbf{0}$  or  $-\mathbf{v} \succ \mathbf{0}$ . Consequently, there is a unique vector  $\mathbf{p} \succ \mathbf{0}$  such that

$$A\mathbf{p} = \rho(A)\mathbf{p} \quad \text{and} \quad \mathbf{e}^\top \mathbf{p} = p_1 + \cdots + p_n = 1.$$

**Definition:** The special vector  $\mathbf{p}$  is called the **Perron vector** of  $A \succ \mathbf{0}$  and the associated eigenvalue  $\rho(A)$  is called the **Perron root** of  $A$ .

**Fact:** If  $A \succ \mathbf{0}$  then there are no nonnegative eigenvectors of  $A$  other than the Perron vector  $\mathbf{p}$  and its positive scalar multiples.

**Proof:** Let  $A\mathbf{y} = \lambda\mathbf{y}$  and  $\mathbf{y} \succeq \mathbf{0}$ . Let  $\mathbf{x} \succ \mathbf{0}$  be the Perron vector of  $A^\top$ . Then  $\mathbf{x}^\top \mathbf{y} > 0$ . Hence

$$\rho(A)\mathbf{x}^\top = \mathbf{x}^\top A \implies \rho(A)\mathbf{x}^\top \mathbf{y} = \mathbf{x}^\top A\mathbf{y} = \lambda\mathbf{x}^\top \mathbf{y} \implies \rho(A) = \lambda. \blacksquare$$



# Perron's Theorem

**Theorem:** Let  $A \succ \mathbf{0}$ . Then the following hold:

- $\rho(A) > 0$
- $\rho(A) \in \Lambda(A)$  and  $\rho(A)$  is simple ( $\rho(A)$  is called **Perron root**)

# Perron's Theorem

**Theorem:** Let  $A \succ \mathbf{0}$ . Then the following hold:

- $\rho(A) > 0$
- $\rho(A) \in \Lambda(A)$  and  $\rho(A)$  is simple ( $\rho(A)$  is called **Perron root**)
- There is a unique vector  $\mathbf{p}$  (called **Perron vector**) such that  $A\mathbf{p} = \rho(A)\mathbf{p}$ ,  $\mathbf{p} \succ \mathbf{0}$  and  $\mathbf{e}^\top \mathbf{p} = 1$

# Perron's Theorem

**Theorem:** Let  $A \succ \mathbf{0}$ . Then the following hold:

- $\rho(A) > 0$
- $\rho(A) \in \Lambda(A)$  and  $\rho(A)$  is simple ( $\rho(A)$  is called **Perron root**)
- There is a unique vector  $\mathbf{p}$  (called **Perron vector**) such that  $A\mathbf{p} = \rho(A)\mathbf{p}$ ,  $\mathbf{p} \succ \mathbf{0}$  and  $\mathbf{e}^\top \mathbf{p} = 1$
- There are no nonnegative eigenvectors of  $A$  other than the Perron vector  $\mathbf{p}$  and its positive scalar multiples

# Perron's Theorem

**Theorem:** Let  $A \succ \mathbf{0}$ . Then the following hold:

- $\rho(A) > 0$
- $\rho(A) \in \Lambda(A)$  and  $\rho(A)$  is simple ( $\rho(A)$  is called **Perron root**)
- There is a unique vector  $\mathbf{p}$  (called **Perron vector**) such that  $A\mathbf{p} = \rho(A)\mathbf{p}$ ,  $\mathbf{p} \succ \mathbf{0}$  and  $\mathbf{e}^\top \mathbf{p} = 1$
- There are no nonnegative eigenvectors of  $A$  other than the Perron vector  $\mathbf{p}$  and its positive scalar multiples
- $\rho(A)$  is the only eigenvalue on the spectral circle  $|z| = \rho(A)$

# Perron's Theorem

**Theorem:** Let  $A \succ \mathbf{0}$ . Then the following hold:

- $\rho(A) > 0$
- $\rho(A) \in \Lambda(A)$  and  $\rho(A)$  is simple ( $\rho(A)$  is called **Perron root**)
- There is a unique vector  $\mathbf{p}$  (called **Perron vector**) such that  $A\mathbf{p} = \rho(A)\mathbf{p}$ ,  $\mathbf{p} \succ \mathbf{0}$  and  $\mathbf{e}^\top \mathbf{p} = 1$
- There are no nonnegative eigenvectors of  $A$  other than the Perron vector  $\mathbf{p}$  and its positive scalar multiples
- $\rho(A)$  is the only eigenvalue on the spectral circle  $|z| = \rho(A)$
- **Collatz-Weilandt formula**

$$\rho(A) = \max \left\{ \min_{i, x_i \neq 0} \frac{(A\mathbf{x})_i}{x_i} : \mathbf{x} \succeq \mathbf{0} \text{ and } \mathbf{x} \neq \mathbf{0} \right\}$$

# Google Search Engine

Google search engine does three basic things:

- Crawl the web and locate all web pages with public access.

# Google Search Engine

Google search engine does three basic things:

- **Crawl the web** and locate all web pages with **public access**.
- **Index the data** (create term-document matrix) so that they can be searched efficiently for relevant **key words or phrases**.

# Google Search Engine

Google search engine does three basic things:

- **Crawl the web** and locate all web pages with **public access**.
- **Index the data** (create term-document matrix) so that they can be searched efficiently for relevant **key words or phrases**.
- **Rank each page in the database**, so that when a user does a search and all the pages in the database with the desired information have been found, the **most relevant pages can be presented first**.



# Google Search Engine

Google search engine does three basic things:

- **Crawl the web** and locate all web pages with **public access**.
- **Index the data** (create term-document matrix) so that they can be searched efficiently for relevant **key words or phrases**.
- **Rank each page in the database**, so that when a user does a search and all the pages in the database with the desired information have been found, the **most relevant pages can be presented first**.

**Google's PageRank** algorithm assigns ranks to all the **web pages** and is formulated as a matrix eigenvalue problem.

# World Wide Web as a Graph



Figure : Link from one web page to another web page.

## Web graph:

- Nodes = Web pages
- Edges = links

# Ranking of web pages



The **web** is an example of a **directed graph**. Let all the web pages be ordered as  $P_1, \dots, P_n$ . A link from  $P_i$  to  $P_j$  represents an arrow. Google assigns **rank** to a page based on its **in-links** (incoming links) and **out-links** (outgoing links).

# Ranking of web pages



The **web** is an example of a **directed graph**. Let all the web pages be ordered as  $P_1, \dots, P_n$ . A link from  $P_i$  to  $P_j$  represents an arrow. Google assigns **rank** to a page based on its **in-links** (incoming links) and **out-links** (outgoing links).

**In-links of  $P_j$ :** All the incoming links (pages) to  $P_j$ .

**Out-links of  $P_j$ :** All the outgoing links (pages) from  $P_j$ .

# Ranking of web pages



The **web** is an example of a **directed graph**. Let all the web pages be ordered as  $P_1, \dots, P_n$ . A link from  $P_i$  to  $P_j$  represents an arrow. Google assigns **rank** to a page based on its **in-links** (incoming links) and **out-links** (outgoing links).

**In-links of  $P_j$ :** All the incoming links (pages) to  $P_j$ .

**Out-links of  $P_j$ :** All the outgoing links (pages) from  $P_j$ .

Google assigns a **high rank to a web page** if it has **inlinks** from other pages that have a **high rank**. This **self-referencing** statement can be formulated mathematically as an **eigenvalue problem**.

# Ranking of web pages



The **web** is an example of a **directed graph**. Let all the web pages be ordered as  $P_1, \dots, P_n$ . A link from  $P_i$  to  $P_j$  represents an arrow. Google assigns **rank** to a page based on its **in-links** (incoming links) and **out-links** (outgoing links).

**In-links of  $P_j$ :** All the incoming links (pages) to  $P_j$ .

**Out-links of  $P_j$ :** All the outgoing links (pages) from  $P_j$ .

Google assigns a **high rank to a web page** if it has **inlinks** from other pages that have a **high rank**. This **self-referencing** statement can be formulated mathematically as an **eigenvalue problem**.

However, a **ranking system** based only on the number of **inlinks** is easy to **manipulate**. **Google** overcomes this problem as follows.

# PageRank eigenvalue problem

Let  $x_j \geq 0$  be the rank of page  $P_j$ . Then  $x_j > x_k \implies P_j$  is **more important** than page  $P_k$ .

# PageRank eigenvalue problem

Let  $x_j \geq 0$  be the rank of page  $P_j$ . Then  $x_j > x_k \implies P_j$  is more important than page  $P_k$ . If page  $P_j$  contains  $n_j$  out-links, one of which links to page  $P_k$ , then page  $P_k$ 's score is boosted by  $x_j/n_j$ .



# PageRank eigenvalue problem

Let  $x_j \geq 0$  be the rank of page  $P_j$ . Then  $x_j > x_k \implies P_j$  is more important than page  $P_k$ . If page  $P_j$  contains  $n_j$  out-links, one of which links to page  $P_k$ , then page  $P_k$ 's score is boosted by  $x_j/n_j$ .

Thus each web page gets a total of one vote, weighted by that web page's score, that is evenly divided up among all of its outgoing links.

# PageRank eigenvalue problem

Let  $x_j \geq 0$  be the rank of page  $P_j$ . Then  $x_j > x_k \implies P_j$  is more important than page  $P_k$ . If page  $P_j$  contains  $n_j$  out-links, one of which links to page  $P_k$ , then page  $P_k$ 's score is boosted by  $x_j/n_j$ .

Thus each web page gets a total of one vote, weighted by that web page's score, that is evenly divided up among all of its outgoing links.

Let  $L_k$  be the set of in-links of  $P_k$ . Then the rank  $x_k$  of  $P_k$  is defined by

$$x_k = \sum_{j \in L_k} \frac{x_j}{n_j} \quad (1)$$

where  $n_j$  is the number of outgoing links (outlinks) from page  $P_j$ . Setting  $\mathbf{v} := [x_1, \dots, x_n]^T$ , equation (1) is rewritten as

# PageRank eigenvalue problem

Let  $x_j \geq 0$  be the rank of page  $P_j$ . Then  $x_j > x_k \implies P_j$  is **more important** than page  $P_k$ . If page  $P_j$  contains  $n_j$  **out-links**, one of which links to page  $P_k$ , then **page  $P_k$ 's score is boosted by  $x_j/n_j$** .

Thus each web page gets a **total of one vote**, weighted by that web page's score, that is evenly divided up among all of its **outgoing links**.

Let  $L_k$  be the **set of in-links of  $P_k$** . Then the rank  $x_k$  of  $P_k$  is defined by

$$x_k = \sum_{j \in L_k} \frac{x_j}{n_j} \quad (1)$$

where  $n_j$  is the **number of outgoing links (outlinks)** from page  $P_j$ . Setting  $\mathbf{v} := [x_1, \dots, x_n]^T$ , equation (1) is rewritten as

$$\text{Eigenvalue problem } \mathbf{H}\mathbf{v} = \mathbf{v} ,$$

where  $\mathbf{H}$  is the **hyperlink matrix** and  $\mathbf{v}$  is the **PageRank vector**.

# World Wide Web as a Hyperlink matrix

Let  $n_j$  be the number of outlinks of the web page  $P_j$ . Then

$$\mathbf{H} = \begin{bmatrix} h_{11} & \cdots & h_{1n} \\ \vdots & \cdots & \vdots \\ h_{n1} & \cdots & h_{nn} \end{bmatrix},$$

$h_{ij} = 1/n_j$  if there is a link from  $P_j$  to page  $P_i$  else  $h_{ij} = 0$ .

# World Wide Web as a Hyperlink matrix

Let  $n_j$  be the number of outlinks of the web page  $P_j$ . Then

$$\mathbf{H} = \begin{bmatrix} h_{11} & \cdots & h_{1n} \\ \vdots & \cdots & \vdots \\ h_{n1} & \cdots & h_{nn} \end{bmatrix},$$

$h_{ij} = 1/n_j$  if there is a link from  $P_j$  to page  $P_i$  else  $h_{ij} = 0$ .

Number of nonzero entries in  $\mathbf{H}\mathbf{e}_j$  = number of outlinks of  $P_j$ .

Number of nonzero entries in  $\mathbf{e}_i^T \mathbf{H}$  = number of inlinks of  $P_i$ .

# World Wide Web as a Hyperlink matrix

Let  $n_j$  be the number of outlinks of the web page  $P_j$ . Then

$$\mathbf{H} = \begin{bmatrix} h_{11} & \cdots & h_{1n} \\ \vdots & \cdots & \vdots \\ h_{n1} & \cdots & h_{nn} \end{bmatrix},$$

$h_{ij} = 1/n_j$  if there is a link from  $P_j$  to page  $P_i$  else  $h_{ij} = 0$ .

Number of nonzero entries in  $\mathbf{H}\mathbf{e}_j$  = number of outlinks of  $P_j$ .

Number of nonzero entries in  $\mathbf{e}_i^\top \mathbf{H}$  = number of inlinks of  $P_i$ .

Thus  $\mathbf{H}$  is non-negative:  $\mathbf{H} \succeq 0$  and  $\mathbf{H}\mathbf{e}_k = \mathbf{0}$  if  $P_k$  has no outlink.

On the other hand,  $\mathbf{e}^\top \mathbf{H}\mathbf{e}_k = 1$  if the page  $P_k$  has outlinks.

Hence  $\mathbf{H}$  is column sub-stochastic:  $\mathbf{e}^\top \mathbf{H}\mathbf{e}_j \leq 1$  for  $j = 1 : n$ .

# World Wide Web as a Hyperlink matrix

Let  $n_j$  be the number of outlinks of the web page  $P_j$ . Then

$$\mathbf{H} = \begin{bmatrix} h_{11} & \cdots & h_{1n} \\ \vdots & \cdots & \vdots \\ h_{n1} & \cdots & h_{nn} \end{bmatrix},$$

$h_{ij} = 1/n_j$  if there is a link from  $P_j$  to page  $P_i$  else  $h_{ij} = 0$ .

Number of nonzero entries in  $\mathbf{H}\mathbf{e}_j$  = number of outlinks of  $P_j$ .

Number of nonzero entries in  $\mathbf{e}_i^\top \mathbf{H}$  = number of inlinks of  $P_i$ .

Thus  $\mathbf{H}$  is non-negative:  $\mathbf{H} \succeq 0$  and  $\mathbf{H}\mathbf{e}_k = \mathbf{0}$  if  $P_k$  has no outlink.

On the other hand,  $\mathbf{e}^\top \mathbf{H}\mathbf{e}_k = 1$  if the page  $P_k$  has outlinks.

Hence  $\mathbf{H}$  is column sub-stochastic:  $\mathbf{e}^\top \mathbf{H}\mathbf{e}_j \leq 1$  for  $j = 1 : n$ .

**Dangling node:** If a page  $P_k$  has no outlink ( $\mathbf{H}\mathbf{e}_k = \mathbf{0}$ ) then it is called a dangling node.

# World Wide Web as a Hyperlink matrix

Let  $n_j$  be the number of outlinks of the web page  $P_j$ . Then

$$\mathbf{H} = \begin{bmatrix} h_{11} & \cdots & h_{1n} \\ \vdots & \cdots & \vdots \\ h_{n1} & \cdots & h_{nn} \end{bmatrix},$$

$h_{ij} = 1/n_j$  if there is a link from  $P_j$  to page  $P_i$  else  $h_{ij} = 0$ .

Number of nonzero entries in  $\mathbf{H}\mathbf{e}_j$  = number of outlinks of  $P_j$ .

Number of nonzero entries in  $\mathbf{e}_i^\top \mathbf{H}$  = number of inlinks of  $P_i$ .

Thus  $\mathbf{H}$  is non-negative:  $\mathbf{H} \succeq 0$  and  $\mathbf{H}\mathbf{e}_k = \mathbf{0}$  if  $P_k$  has no outlink.

On the other hand,  $\mathbf{e}^\top \mathbf{H}\mathbf{e}_k = 1$  if the page  $P_k$  has outlinks.


Hence  $\mathbf{H}$  is column sub-stochastic:  $\mathbf{e}^\top \mathbf{H}\mathbf{e}_j \leq 1$  for  $j = 1 : n$ .

**Dangling node:** If a page  $P_k$  has no outlink ( $\mathbf{H}\mathbf{e}_k = \mathbf{0}$ ) then it is called a dangling node. If there are no dangling nodes then  $\mathbf{e}^\top \mathbf{H} = \mathbf{e}^\top$ . Hence  $\mathbf{H}$  is column stochastic.



## Example

Column stochastic hyperlink matrix  $\mathbf{H}$ .



$$\mathbf{H} = \begin{bmatrix} 0 & 0 & 0 & 0 & 1 \\ \frac{1}{2} & 0 & 0 & 0 & 0 \\ 0 & \frac{1}{3} & 0 & 0 & 0 \\ \frac{1}{2} & \frac{1}{3} & 1 & 0 & 0 \\ 0 & \frac{1}{3} & 0 & 1 & 0 \end{bmatrix} \text{ and } \mathbf{e}^\top \mathbf{H} = \mathbf{e}^\top.$$

There is a unique PageRank vector  $\mathbf{v} \succ 0$  such that  $\mathbf{H}\mathbf{v} = \mathbf{v}$  and  $\mathbf{e}^\top \mathbf{v} = 1$ . The entries of  $\mathbf{v}$  are the ranks of web pages.

## Nonunique ranking

If  $\mathbf{H}$  is column stochastic then  $\mathbf{e}^\top \mathbf{H} = \mathbf{e}^\top$  which shows that  $\mathbf{e}$  is a left eigenvector of  $\mathbf{H}$  corresponding the eigenvalue 1.

## Nonunique ranking

If  $\mathbf{H}$  is column stochastic then  $\mathbf{e}^\top \mathbf{H} = \mathbf{e}^\top$  which shows that  $\mathbf{e}$  is a left eigenvector of  $\mathbf{H}$  corresponding the eigenvalue 1.

Now  $\dim N(\mathbf{H} - I) > 1$  leads to non-unique ranking. For ranking, we require  $\dim N(\mathbf{H} - I) = 1$  and a vector  $\mathbf{v} \succ \mathbf{0}$  such that  $\mathbf{H}\mathbf{v} = \mathbf{v}$  and  $\mathbf{e}^\top \mathbf{v} = 1$ .

# Nonunique ranking

If  $\mathbf{H}$  is column stochastic then  $\mathbf{e}^\top \mathbf{H} = \mathbf{e}^\top$  which shows that  $\mathbf{e}$  is a left eigenvector of  $\mathbf{H}$  corresponding the eigenvalue 1.

Now  $\dim N(\mathbf{H} - I) > 1$  leads to non-unique ranking. For ranking, we require  $\dim N(\mathbf{H} - I) = 1$  and a vector  $\mathbf{v} \succ \mathbf{0}$  such that  $\mathbf{H}\mathbf{v} = \mathbf{v}$  and  $\mathbf{e}^\top \mathbf{v} = 1$ .

**Theorem:** Suppose that  $\mathbf{H} \succ \mathbf{0}$  and  $\mathbf{e}^\top \mathbf{H} = \mathbf{e}^\top$ . Then

- 1 is the simple eigenvalue of  $\mathbf{H}$  and there exists  $\mathbf{v} \succ \mathbf{0}$  such that  $\mathbf{H}\mathbf{v} = \mathbf{v}$ . If  $\mathbf{v} \in N(\mathbf{H} - I)$  is nonzero then either  $\mathbf{v} \succ \mathbf{0}$  or  $-\mathbf{v} \succ \mathbf{0}$ .

# Nonunique ranking

If  $\mathbf{H}$  is column stochastic then  $\mathbf{e}^\top \mathbf{H} = \mathbf{e}^\top$  which shows that  $\mathbf{e}$  is a left eigenvector of  $\mathbf{H}$  corresponding the eigenvalue 1.

Now  $\dim N(\mathbf{H} - I) > 1$  leads to non-unique ranking. For ranking, we require  $\dim N(\mathbf{H} - I) = 1$  and a vector  $\mathbf{v} \succ \mathbf{0}$  such that  $\mathbf{H}\mathbf{v} = \mathbf{v}$  and  $\mathbf{e}^\top \mathbf{v} = 1$ .

**Theorem:** Suppose that  $\mathbf{H} \succ \mathbf{0}$  and  $\mathbf{e}^\top \mathbf{H} = \mathbf{e}^\top$ . Then

- 1 is the simple eigenvalue of  $\mathbf{H}$  and there exists  $\mathbf{v} \succ \mathbf{0}$  such that  $\mathbf{H}\mathbf{v} = \mathbf{v}$ . If  $\mathbf{v} \in N(\mathbf{H} - I)$  is nonzero then either  $\mathbf{v} \succ \mathbf{0}$  or  $-\mathbf{v} \succ \mathbf{0}$ .
- 1 is the **dominant** eigenvalue, that is, if  $\lambda_2, \dots, \lambda_n$  are eigenvalues of  $\mathbf{H}$  then  $|\lambda_j| < 1$  for  $j = 2 : n$ .

# Nonunique ranking

If  $\mathbf{H}$  is column stochastic then  $\mathbf{e}^\top \mathbf{H} = \mathbf{e}^\top$  which shows that  $\mathbf{e}$  is a left eigenvector of  $\mathbf{H}$  corresponding the eigenvalue 1.

Now  $\dim N(\mathbf{H} - I) > 1$  leads to non-unique ranking. For ranking, we require  $\dim N(\mathbf{H} - I) = 1$  and a vector  $\mathbf{v} \succ \mathbf{0}$  such that  $\mathbf{H}\mathbf{v} = \mathbf{v}$  and  $\mathbf{e}^\top \mathbf{v} = 1$ .

**Theorem:** Suppose that  $\mathbf{H} \succ \mathbf{0}$  and  $\mathbf{e}^\top \mathbf{H} = \mathbf{e}^\top$ . Then

- 1 is the simple eigenvalue of  $\mathbf{H}$  and there exists  $\mathbf{v} \succ \mathbf{0}$  such that  $\mathbf{H}\mathbf{v} = \mathbf{v}$ . If  $\mathbf{v} \in N(\mathbf{H} - I)$  is nonzero then either  $\mathbf{v} \succ \mathbf{0}$  or  $-\mathbf{v} \succ \mathbf{0}$ .
- 1 is the **dominant** eigenvalue, that is, if  $\lambda_2, \dots, \lambda_n$  are eigenvalues of  $\mathbf{H}$  then  $|\lambda_j| < 1$  for  $j = 2 : n$ .
- any other positive eigenvector of  $\mathbf{H}$  is a scalar multiple of  $\mathbf{v}$ .

## Adjustment of hyperlink matrix

The web is full of dangling nodes (e.g., pdf files, image files, etc.). To fix this problem, Brin and Page replaced zero columns of  $\mathbf{H}$  with  $\frac{1}{n}\mathbf{e}$ . Define

$$\mathbf{S} := \mathbf{H} + \frac{1}{n}\mathbf{e}\mathbf{a}^\top,$$

where  $a_j = 1$  if  $P_j$  is a dangling node (i.e.,  $\mathbf{H}\mathbf{e}_j = \mathbf{0}$ ) and  $a_j = 0$  else.

## Adjustment of hyperlink matrix

The web is full of dangling nodes (e.g., pdf files, image files, etc.). To fix this problem, Brin and Page replaced zero columns of  $\mathbf{H}$  with  $\frac{1}{n}\mathbf{e}$ . Define

$$\mathbf{S} := \mathbf{H} + \frac{1}{n}\mathbf{e}\mathbf{a}^\top,$$

where  $a_j = 1$  if  $P_j$  is a dangling node (i.e.,  $\mathbf{H}\mathbf{e}_j = \mathbf{0}$ ) and  $a_j = 0$  else. Then  $\mathbf{e}^\top \mathbf{S} = \mathbf{e}^\top$ .




## Adjustment of hyperlink matrix

The web is full of dangling nodes (e.g., pdf files, image files, etc.). To fix this problem, Brin and Page replaced zero columns of  $\mathbf{H}$  with  $\frac{1}{n}\mathbf{e}$ . Define

$$\mathbf{S} := \mathbf{H} + \frac{1}{n}\mathbf{e}\mathbf{a}^\top,$$

where  $a_j = 1$  if  $P_j$  is a dangling node (i.e.,  $\mathbf{H}\mathbf{e}_j = \mathbf{0}$ ) and  $a_j = 0$  else. Then  $\mathbf{e}^\top \mathbf{S} = \mathbf{e}^\top$ .

The binary vector  $\mathbf{a}$  is called the **dangling node vector**. Note that  $\mathbf{S}$  is column stochastic and the adjustment amounts to adding artificial links to the dangling nodes.



# The Google matrix

The matrix  $\mathbf{S}$ , however, cannot guarantee existence of a unique PageRank vector. So, Brin and Page made the final adjustment to obtain the **Google matrix**

$$\mathbf{G} := \alpha \mathbf{S} + (1 - \alpha) \mathbf{E}, \quad 0 \leq \alpha \leq 1,$$

where  $\mathbf{E} = \frac{1}{n} \mathbf{e} \mathbf{e}^\top$  is the **teleportation** matrix which enables jumping to any page.

# The Google matrix

The matrix  $\mathbf{S}$ , however, cannot guarantee existence of a unique PageRank vector. So, Brin and Page made the final adjustment to obtain the **Google matrix**

$$\mathbf{G} := \alpha \mathbf{S} + (1 - \alpha) \mathbf{E}, \quad 0 \leq \alpha \leq 1,$$

where  $\mathbf{E} = \frac{1}{n} \mathbf{e} \mathbf{e}^\top$  is the **teleportation** matrix which enables jumping to any page. If  $\alpha = 0.6$  then 60% of the time the surfer follows hyperlink structure of web and 40% of the time teleports to a random new page.

# The Google matrix

The matrix  $\mathbf{S}$ , however, cannot guarantee existence of a unique PageRank vector. So, Brin and Page made the final adjustment to obtain the **Google matrix**

$$\mathbf{G} := \alpha \mathbf{S} + (1 - \alpha) \mathbf{E}, \quad 0 \leq \alpha \leq 1,$$

where  $\mathbf{E} = \frac{1}{n} \mathbf{e} \mathbf{e}^\top$  is the **teleportation** matrix which enables jumping to any page. If  $\alpha = 0.6$  then 60% of the time the surfer follows hyperlink structure of web and 40% of the time teleports to a random new page.

**Fact:** The Google matrix  $\mathbf{G}$  has the following properties.

- $\mathbf{G}$  is **positive** ( $\mathbf{G} \succ \mathbf{0}$ ) and **column stochastic** ( $\mathbf{e}^\top \mathbf{G} = \mathbf{e}^\top$ ).
- There is a unique vector  $\mathbf{v}$  such that  $\mathbf{G} \mathbf{v} = \mathbf{v}$ ,  $\mathbf{v} \succ \mathbf{0}$  and  $\mathbf{e}^\top \mathbf{v} = 1$ .
- $\mathbf{G}$  is the rank-1 update of  $\mathbf{H}$  :

$$\mathbf{G} = \alpha \mathbf{S} + (1 - \alpha) \frac{1}{n} \mathbf{e} \mathbf{e}^\top = \alpha \mathbf{H} + \frac{1}{n} \mathbf{e} (\alpha \mathbf{a} + (1 - \alpha) \mathbf{e})^\top.$$

# The Google matrix

**Theorem:** Let  $\mathbf{G} := \alpha \mathbf{S} + (1 - \alpha) \mathbf{E}$  be the Google matrix.

- If  $\Lambda(\mathbf{S}) = \{1, \lambda_2, \dots, \lambda_n\}$  then  $\Lambda(\mathbf{G}) = \{1, \alpha\lambda_2, \dots, \alpha\lambda_n\}$ .

# The Google matrix

**Theorem:** Let  $\mathbf{G} := \alpha\mathbf{S} + (1 - \alpha)\mathbf{E}$  be the Google matrix.

- If  $\Lambda(\mathbf{S}) = \{1, \lambda_2, \dots, \lambda_n\}$  then  $\Lambda(\mathbf{G}) = \{1, \alpha\lambda_2, \dots, \alpha\lambda_n\}$ .
- $\mathbf{G}\mathbf{v} = \mathbf{v}$  and  $\mathbf{e}^\top \mathbf{v} = 1$  yields

$$\mathbf{v} = \alpha\mathbf{S}\mathbf{v} + (1 - \alpha)\frac{1}{n}\mathbf{e} \implies (\mathbf{I} - \alpha\mathbf{S})\mathbf{v} = (1 - \alpha)\frac{1}{n}\mathbf{e}.$$

# The Google matrix

**Theorem:** Let  $\mathbf{G} := \alpha\mathbf{S} + (1 - \alpha)\mathbf{E}$  be the Google matrix.

- If  $\Lambda(\mathbf{S}) = \{1, \lambda_2, \dots, \lambda_n\}$  then  $\Lambda(\mathbf{G}) = \{1, \alpha\lambda_2, \dots, \alpha\lambda_n\}$ .
- $\mathbf{G}\mathbf{v} = \mathbf{v}$  and  $\mathbf{e}^\top \mathbf{v} = 1$  yields

$$\mathbf{v} = \alpha\mathbf{S}\mathbf{v} + (1 - \alpha)\frac{1}{n}\mathbf{e} \implies (\mathbf{I} - \alpha\mathbf{S})\mathbf{v} = (1 - \alpha)\frac{1}{n}\mathbf{e}.$$

- If  $\mathbf{e}^\top \mathbf{x} = 1$  then  $\mathbf{G}\mathbf{x} = \alpha\mathbf{S}\mathbf{x} + (1 - \alpha)\frac{1}{n}\mathbf{e}$ . Hence

# The Google matrix

**Theorem:** Let  $\mathbf{G} := \alpha\mathbf{S} + (1 - \alpha)\mathbf{E}$  be the Google matrix.

- If  $\Lambda(\mathbf{S}) = \{1, \lambda_2, \dots, \lambda_n\}$  then  $\Lambda(\mathbf{G}) = \{1, \alpha\lambda_2, \dots, \alpha\lambda_n\}$ .
- $\mathbf{G}\mathbf{v} = \mathbf{v}$  and  $\mathbf{e}^\top \mathbf{v} = 1$  yields

$$\mathbf{v} = \alpha\mathbf{S}\mathbf{v} + (1 - \alpha)\frac{1}{n}\mathbf{e} \implies (\mathbf{I} - \alpha\mathbf{S})\mathbf{v} = (1 - \alpha)\frac{1}{n}\mathbf{e}.$$

- If  $\mathbf{e}^\top \mathbf{x} = 1$  then  $\mathbf{G}\mathbf{x} = \alpha\mathbf{S}\mathbf{x} + (1 - \alpha)\frac{1}{n}\mathbf{e}$ . Hence

$$\mathbf{G}\mathbf{x} - \mathbf{G}\mathbf{v} = \alpha\mathbf{S}(\mathbf{x} - \mathbf{v}).$$

Consequently, we have

$$\|\mathbf{G}(\mathbf{x} - \mathbf{v})\|_1 = \alpha\|\mathbf{S}(\mathbf{x} - \mathbf{v})\|_1 \leq \alpha\|\mathbf{x} - \mathbf{v}\|_1,$$

where  $\|\mathbf{x}\|_1 = |x_1| + \dots + |x_n|$ .



# The Power method

**Power method:** Choose  $\mathbf{v}_0 \succeq \mathbf{0}$  such that  $\mathbf{e}^\top \mathbf{v}_0 = 1$ . Define  $\mathbf{u}_j = \alpha \mathbf{S} \mathbf{v}_{j-1} + (1 - \alpha) \frac{1}{n} \mathbf{e}$  and  $\mathbf{v}_j = \mathbf{u}_j / \mathbf{e}^\top \mathbf{u}_j$  for  $j = 1, 2, \dots$ . Then

$$\|\mathbf{v}_j - \mathbf{v}\|_1 \leq \alpha^j \|\mathbf{v} - \mathbf{v}_0\|_1.$$

---

**Algorithm:** The Power Method

**Input:** Vector  $\mathbf{v}_0 \succeq \mathbf{0}$  such that  $\mathbf{e}^\top \mathbf{v}_0 = 1$  and number of iterations  $\ell$

**Output:** PageRank vector  $\mathbf{v}_\ell$

---

for  $j = 1: \ell$

$\mathbf{u}_j := \alpha \mathbf{S} \mathbf{v}_{j-1} + (1 - \alpha) \mathbf{e} / n$       (application of  $\mathbf{G}^j$ )

$\mathbf{v}_j := \mathbf{u}_j / \mathbf{e}^\top \mathbf{u}_j$       (normalization)

end

---

## Further reading

Want to know more about Google search engine?

## Further reading

### Want to know more about Google search engine?

Read the following paper/book.

- K. Bryan and T. Leise, [The \\$25,000,000,000 Eigenvector: The Linear Algebra behind Google](#), SIAM Review, 48(2006), pp.569-581.
- Amy N. Langville and Carl D. Meyer, [Google's PageRank and Beyond: The Science of Search Engine Rankings](#), Princeton University Press, 2006

# The power method

The power method (or power iteration) is one of the oldest and simplest iterative method for computing an eigenpair of  $A$ . It is designed for a matrices  $A$  with one dominant eigenvalue. Starting with an arbitrary nonzero vector  $v_0 \in \mathbb{C}^n$ , the power method computes the sequence

$$v_0, Av_0, \dots, A^j v_0, \dots$$

Setting  $v_j := A^j v_0$ , we have  $v_j = Av_{j-1}$  for  $j = 1, 2, \dots$

# The power method

The power method (or power iteration) is one of the oldest and simplest iterative method for computing an eigenpair of  $A$ . It is designed for a matrices  $A$  with one dominant eigenvalue. Starting with an arbitrary nonzero vector  $v_0 \in \mathbb{C}^n$ , the power method computes the sequence

$$v_0, Av_0, \dots, A^j v_0, \dots$$

Setting  $v_j := A^j v_0$ , we have  $v_j = Av_{j-1}$  for  $j = 1, 2, \dots$

---

**Algorithm:** The Power Method

**Input:** Nonzero vector  $v_0$  and the number of iterations  $\ell$

**Output:** Dominant eigenpair  $(\mu_\ell, v_\ell)$

---

```
for  $j = 1: \ell$   
     $w_j := Av_{j-1}$            (application of  $A^j$ )  
     $v_j := w_j / \|w_j\|_2$      (normalization)  
     $\mu_j := v_j^* Av_j$        (Rayleigh quotient)  
end
```

---

# The inverse iteration

The power method can only be used to compute a dominant eigenpair of  $A$ . It turns out that a simple *shift-and-invert* strategy can be utilized to compute any desired eigenvalue of  $A$  and an associated eigenvector. Assume for the moment that  $A$  is invertible. Then

$$Av = \lambda v \iff A^{-1}v = \frac{1}{\lambda}v.$$

This shows that  $1/\lambda$  is the dominant eigenvalue of  $A^{-1} \iff \lambda$  is the smallest eigenvalue of  $A$ .

# The inverse iteration

The power method can only be used to compute a dominant eigenpair of  $A$ . It turns out that a simple *shift-and-invert* strategy can be utilized to compute any desired eigenvalue of  $A$  and an associated eigenvector. Assume for the moment that  $A$  is invertible. Then

$$Av = \lambda v \iff A^{-1}v = \frac{1}{\lambda}v.$$

This shows that  $1/\lambda$  is the dominant eigenvalue of  $A^{-1} \iff \lambda$  is the smallest eigenvalue of  $A$ .

If  $s$  is not an eigenvalue of  $A$  then

$$Av = \lambda v \iff (A - sI)v = (\lambda - s)v \iff (A - sI)^{-1}v = \frac{1}{\lambda - s}v.$$

Thus with the help of a shift  $s$  that is closer to a simple eigenvalue  $\lambda$  than to the rest of the spectrum of  $A$ , the transformed eigenvalue  $(\lambda - s)^{-1}$  can be made dominant for the transformed matrix  $(A - sI)^{-1}$ . Hence we can apply the power iteration to the shifted matrix  $(A - sI)^{-1}$  and arrive at the shifted inverse iteration.

# The inverse iteration

---

**Algorithm:** Shifted inverse iteration

**Input:** Nonzero vector  $v_0$ , a shift  $s$  and the number of iterations  $\ell$

**Output:** An eigenpair  $(\mu_\ell, v_\ell)$  such that  $\mu_\ell$  closest to  $s$

---

```
for  $j = 1:\ell$   
     $(A - sI)w_j = v_{j-1}$       (application of  $(A - sI)^{-j}$ )  
     $v_j := w_j / \|w_j\|_2$       (normalization)  
     $\mu_j := v_j^* A v_j$          (Rayleigh quotient)  
end
```

---



# The inverse iteration

---

**Algorithm:** Shifted inverse iteration

**Input:** Nonzero vector  $v_0$ , a shift  $s$  and the number of iterations  $\ell$

**Output:** An eigenpair  $(\mu_\ell, v_\ell)$  such that  $\mu_\ell$  closest to  $s$

---

```
for  $j = 1:\ell$   
     $(A - sI)w_j = v_{j-1}$       (application of  $(A - sI)^{-j}$ )  
     $v_j := w_j / \|w_j\|_2$       (normalization)  
     $\mu_j := v_j^* A v_j$          (Rayleigh quotient)  
end
```

---

The computation of  $w_j$  requires solution of  $(A - sI)w_j = v_j$ . Since  $(A - sI)$  is not changed during the iteration, we compute LU factorization  $P(A - sI) = LU$  only once, where  $P$  is a permutation matrix,  $L$  is a unit lower triangular matrix and  $U$  is an upper triangular matrix.

# The inverse iteration

---

**Algorithm:** Shifted inverse iteration

**Input:** Nonzero vector  $v_0$ , a shift  $s$  and the number of iterations  $\ell$

**Output:** An eigenpair  $(\mu_\ell, v_\ell)$  such that  $\mu_\ell$  closest to  $s$

---

```
for  $j = 1: \ell$   
     $(A - sI)w_j = v_{j-1}$       (application of  $(A - sI)^{-j}$ )  
     $v_j := w_j / \|w_j\|_2$       (normalization)  
     $\mu_j := v_j^* A v_j$          (Rayleigh quotient)  
end
```

---

The computation of  $w_j$  requires solution of  $(A - sI)w_j = v_j$ . Since  $(A - sI)$  is not changed during the iteration, we compute LU factorization  $P(A - sI) = LU$  only once, where  $P$  is a permutation matrix,  $L$  is a unit lower triangular matrix and  $U$  is an upper triangular matrix.

This will reduce the cost of computation to  $2n^2$  flops per iteration. This is an enormous saving as solving the system without exploiting LU factorization would cost  $2n^3/3$  flops per iteration. Hence the total cost of performing  $\ell$  iterations of the inverse iteration is  $(2n^3/3 + 4n^2\ell)$  flops.

## Rayleigh quotient iteration (RQI)

The shifted inverse iteration converges rapidly if the shift parameter  $s \in \mathbb{C}$  is close to an eigenvalue of  $A$ . The Rayleigh quotient  $q(x) := x^*Ax/x^*x$  approximates an eigenvalue  $A$ . Hence it makes sense to use Rayleigh quotient as shift in the shifted inverse iteration. The resulting algorithm is called the *Rayleigh quotient iteration* (RQI).

# Rayleigh quotient iteration (RQI)

The shifted inverse iteration converges rapidly if the shift parameter  $s \in \mathbb{C}$  is close to an eigenvalue of  $A$ . The Rayleigh quotient  $q(x) := x^*Ax/x^*x$  approximates an eigenvalue  $\lambda$ . Hence it makes sense to use Rayleigh quotient as shift in the shifted inverse iteration. The resulting algorithm is called the *Rayleigh quotient iteration* (RQI).

---

**Algorithm:** Rayleigh quotient iteration (RQI)

**Input:** Nonzero vector  $v_0$  and the number of iterations  $\ell$

**Output:** Approximate eigenpair  $(\mu_\ell, v_\ell)$

---

```
 $v_0 := v_0 / \|v_0\|_2$       (normalize  $v_0$ )  
 $\mu_0 := v_0^* A v_0$       ( Rayleigh quotient)  
for  $j = 1: \ell$   
     $(A - \mu_{j-1}I)w_j = v_{j-1}$     (application of  $(A - \mu_{j-1}I)^{-j}$ )  
     $v_j := w_j / \|w_j\|_2$       (normalization)  
     $\mu_j := v_j^* A v_j$       (Rayleigh quotient)  
end
```

---

**Remark:** RQI requires a new LU factorization in each iteration.