

MA423 Matrix Computations

Lecture 10: Vector and matrix norms

Rafikul Alam
Department of Mathematics
IIT Guwahati

Outline

- Vector norms
- Matrix norms
- Unitarily invariant norms

Vector norms

Let \mathcal{V} be a vector space over \mathbb{C} . Then a function $\|\cdot\| : \mathcal{V} \rightarrow \mathbb{R}$ is called a **norm on \mathcal{V}** if it satisfies the three fundamental properties:

- (a) **Positive definiteness:** $\|v\| \geq 0$ and $\|v\| = 0 \iff v = 0$.
- (b) **Positively homogeneous:** $\|\alpha v\| = |\alpha| \|v\|$ for $\alpha \in \mathbb{C}$ and $v \in \mathcal{V}$.
- (c) **Triangle inequality:** $\|u + v\| \leq \|u\| + \|v\|$ for $u, v \in \mathcal{V}$.

Vector norms

Let \mathcal{V} be a vector space over \mathbb{C} . Then a function $\|\cdot\| : \mathcal{V} \rightarrow \mathbb{R}$ is called a **norm on \mathcal{V}** if it satisfies the three fundamental properties:

- (a) **Positive definiteness:** $\|v\| \geq 0$ and $\|v\| = 0 \iff v = 0$.
- (b) **Positively homogeneous:** $\|\alpha v\| = |\alpha| \|v\|$ for $\alpha \in \mathbb{C}$ and $v \in \mathcal{V}$.
- (c) **Triangle inequality:** $\|u + v\| \leq \|u\| + \|v\|$ for $u, v \in \mathcal{V}$.

Example: Consider \mathbb{C}^n and the vector norms given by

1-norm: $\|x\|_1 := |x_1| + \cdots + |x_n|.$

2-norm: $\|x\|_2 := \sqrt{|x_1|^2 + \cdots + |x_n|^2}.$

∞ -norm: $\|x\|_\infty := \max_{1 \leq j \leq n} |x_j|.$

Vector norms

Let \mathcal{V} be a vector space over \mathbb{C} . Then a function $\|\cdot\| : \mathcal{V} \rightarrow \mathbb{R}$ is called a **norm on \mathcal{V}** if it satisfies the three fundamental properties:

- (a) **Positive definiteness:** $\|v\| \geq 0$ and $\|v\| = 0 \iff v = 0$.
- (b) **Positively homogeneous:** $\|\alpha v\| = |\alpha| \|v\|$ for $\alpha \in \mathbb{C}$ and $v \in \mathcal{V}$.
- (c) **Triangle inequality:** $\|u + v\| \leq \|u\| + \|v\|$ for $u, v \in \mathcal{V}$.

Example: Consider \mathbb{C}^n and the vector norms given by

1-norm: $\|x\|_1 := |x_1| + \cdots + |x_n|.$

2-norm: $\|x\|_2 := \sqrt{|x_1|^2 + \cdots + |x_n|^2}.$

∞ -norm: $\|x\|_\infty := \max_{1 \leq j \leq n} |x_j|.$

More generally, for $1 \leq p < \infty$, the **Höder p -norm** is given by

$$\|x\|_p := (|x_1|^p + \cdots + |x_n|^p)^{1/p}.$$

Matrix norms

Let $A \in \mathbb{C}^{m \times n}$. Then $A : \mathbb{C}^n \longrightarrow \mathbb{C}^m$, $x \longmapsto Ax$, is a linear map. Suppose \mathbb{C}^n and \mathbb{C}^m are equipped with norms. Then

$$\|A\| := \max_{x \neq 0} \frac{\|Ax\|}{\|x\|} = \max_{\|x\|=1} \|Ax\|$$

defines a norm on $\mathbb{C}^{m \times n}$ and is called an **induced matrix norm** or a **subordinate matrix norm**.

Matrix norms

Let $A \in \mathbb{C}^{m \times n}$. Then $A : \mathbb{C}^n \rightarrow \mathbb{C}^m$, $x \mapsto Ax$, is a linear map. Suppose \mathbb{C}^n and \mathbb{C}^m are equipped with norms. Then

$$\|A\| := \max_{x \neq 0} \frac{\|Ax\|}{\|x\|} = \max_{\|x\|=1} \|Ax\|$$

defines a norm on $\mathbb{C}^{m \times n}$ and is called an **induced matrix norm** or a **subordinate matrix norm**.

For the identity matrix $\|Ix\| = \|x\|$ and hence $\|I\| = 1$. Note that

$$\|Ax\| \leq \|A\| \|x\|$$

for all $x \in \mathbb{C}^n$.

Matrix norms

The matrix norm of A induced by the Hölder p -norm is denoted by $\|A\|_p$. Then $\|A\|_1$, $\|A\|_2$ and $\|A\|_\infty$ are called 1-norm, 2-norm and ∞ -norm of A , respectively. Also $\|A\|_2$ is called the spectral norm of A .

Matrix norms

The matrix norm of A induced by the Hölder p -norm is denoted by $\|A\|_p$. Then $\|A\|_1$, $\|A\|_2$ and $\|A\|_\infty$ are called 1-norm, 2-norm and ∞ -norm of A , respectively. Also $\|A\|_2$ is called the spectral norm of A .

Theorem: Let A be an $m \times n$ matrix. Then

$$\begin{aligned}\|A\|_1 &= \max_{1 \leq j \leq n} \|Ae_j\|_1 = \max_{1 \leq j \leq n} \|A(:,j)\|_1 \\ \|A\|_2 &= \sqrt{\lambda_{\max}(A^*A)} \\ \|A\|_\infty &= \max_{1 \leq i \leq m} \|e_i^\top A\|_1 = \max_{1 \leq i \leq m} \|A(i,:)\|_1,\end{aligned}$$

where $\lambda_{\max}(A^*A)$ denotes that largest eigenvalue of A^*A .

Matrix norms

The matrix norm of A induced by the Hölder p -norm is denoted by $\|A\|_p$. Then $\|A\|_1$, $\|A\|_2$ and $\|A\|_\infty$ are called 1-norm, 2-norm and ∞ -norm of A , respectively. Also $\|A\|_2$ is called the spectral norm of A .

Theorem: Let A be an $m \times n$ matrix. Then

$$\begin{aligned}\|A\|_1 &= \max_{1 \leq j \leq n} \|Ae_j\|_1 = \max_{1 \leq j \leq n} \|A(:,j)\|_1 \\ \|A\|_2 &= \sqrt{\lambda_{\max}(A^*A)} \\ \|A\|_\infty &= \max_{1 \leq i \leq m} \|e_i^\top A\|_1 = \max_{1 \leq i \leq m} \|A(i,:)\|_1,\end{aligned}$$

where $\lambda_{\max}(A^*A)$ denotes that largest eigenvalue of A^*A .

Proof: We have

$$Ax = x_1 Ae_1 + \cdots + x_n Ae_n \Rightarrow \|Ax\|_1 \leq \max_{1 \leq j \leq n} \|Ae_j\|_1 \|x\|_1.$$

Matrix norms

The matrix norm of A induced by the Hölder p -norm is denoted by $\|A\|_p$. Then $\|A\|_1$, $\|A\|_2$ and $\|A\|_\infty$ are called 1-norm, 2-norm and ∞ -norm of A , respectively. Also $\|A\|_2$ is called the spectral norm of A .

Theorem: Let A be an $m \times n$ matrix. Then

$$\begin{aligned}\|A\|_1 &= \max_{1 \leq j \leq n} \|Ae_j\|_1 = \max_{1 \leq j \leq n} \|A(:,j)\|_1 \\ \|A\|_2 &= \sqrt{\lambda_{\max}(A^*A)} \\ \|A\|_\infty &= \max_{1 \leq i \leq m} \|e_i^\top A\|_1 = \max_{1 \leq i \leq m} \|A(i,:)\|_1,\end{aligned}$$

where $\lambda_{\max}(A^*A)$ denotes that largest eigenvalue of A^*A .

Proof: We have

$Ax = x_1 Ae_1 + \cdots + x_n Ae_n \Rightarrow \|Ax\|_1 \leq \max_{1 \leq j \leq n} \|Ae_j\|_1 \|x\|_1$. This yields $\|A\|_1 \leq \max_{1 \leq j \leq n} \|Ae_j\|_1$.

Matrix norms

The matrix norm of A induced by the Hölder p -norm is denoted by $\|A\|_p$. Then $\|A\|_1$, $\|A\|_2$ and $\|A\|_\infty$ are called 1-norm, 2-norm and ∞ -norm of A , respectively. Also $\|A\|_2$ is called the spectral norm of A .

Theorem: Let A be an $m \times n$ matrix. Then


$$\begin{aligned}\|A\|_1 &= \max_{1 \leq j \leq n} \|Ae_j\|_1 = \max_{1 \leq j \leq n} \|A(:,j)\|_1 \\ \|A\|_2 &= \sqrt{\lambda_{\max}(A^*A)} \\ \|A\|_\infty &= \max_{1 \leq i \leq m} \|e_i^\top A\|_1 = \max_{1 \leq i \leq m} \|A(i,:)\|_1,\end{aligned}$$

where $\lambda_{\max}(A^*A)$ denotes that largest eigenvalue of A^*A .

Proof: We have

$Ax = x_1 Ae_1 + \cdots + x_n Ae_n \Rightarrow \|Ax\|_1 \leq \max_{1 \leq j \leq n} \|Ae_j\|_1 \|x\|_1$. This yields $\|A\|_1 \leq \max_{1 \leq j \leq n} \|Ae_j\|_1$. But $\|Ae_j\|_1 \leq \|A\|_1$ for all $j = 1 : n$. Hence we have $\|A\|_1 = \max_{1 \leq j \leq n} \|Ae_j\|_1$. ■

Spectral norm



Frobenius norm

The spectral norm of a matrix is difficult to compute. By contrast, **Frobenius norm** is easy to compute and provides a nice geometric framework for solving problems. Let A be an m -by- n matrix.

Frobenius norm

The spectral norm of a matrix is difficult to compute. By contrast, **Frobenius norm** is easy to compute and provides a nice geometric framework for solving problems. Let A be an m -by- n matrix.

Then the **Frobenius norm** is given by

$$\|A\|_F := \left(\sum_{i=1}^m \sum_{j=1}^n |A(i,j)|^2 \right)^{1/2} = \sqrt{\text{Tr}(A^*A)} = \sqrt{\text{Tr}(AA^*)},$$

where $\text{Tr}(A^*A)$ is the trace of A^*A .

Frobenius norm

The spectral norm of a matrix is difficult to compute. By contrast, **Frobenius norm** is easy to compute and provides a nice geometric framework for solving problems. Let A be an m -by- n matrix.

Then the **Frobenius norm** is given by

$$\|A\|_F := \left(\sum_{i=1}^m \sum_{j=1}^n |A(i,j)|^2 \right)^{1/2} = \sqrt{\text{Tr}(A^*A)} = \sqrt{\text{Tr}(AA^*)},$$

where $\text{Tr}(A^*A)$ is the trace of A^*A .

Note that $\|A\|_F = \left(\sum_{i=1}^m \|e_i^\top A\|_2^2 \right)^{1/2} = \left(\sum_{j=1}^n \|Ae_j\|_2^2 \right)^{1/2}$. Also note that for an n -by- n identity matrix I , we have $\|I\|_F = \sqrt{n}$ which shows that the Frobenius norm is not an induced matrix norm.

As induced matrix norm for Identity matrix is 1.

Example

Let $D := \text{diag}(\lambda_1, \dots, \lambda_n)$ be a diagonal matrix. Then we have

$$\|A\|_2 = \|A\|_1 = \|A\|_\infty = \max(|\lambda_1|, \dots, |\lambda_n|) \text{ and } \|D\|_F = \sqrt{\sum_{j=1}^n |\lambda_j|^2}.$$

Example

Let $D := \text{diag}(\lambda_1, \dots, \lambda_n)$ be a diagonal matrix. Then we have

$$\|A\|_2 = \|A\|_1 = \|A\|_\infty = \max(|\lambda_1|, \dots, |\lambda_n|) \text{ and } \|D\|_F = \sqrt{\sum_{j=1}^n |\lambda_j|^2}.$$

Consider $A := \begin{bmatrix} 1000 & 999 \\ 999 & 998 \end{bmatrix}$. Then $A^{-1} = \begin{bmatrix} -998 & 999 \\ 999 & -1000 \end{bmatrix}$.

Example

Let $D := \text{diag}(\lambda_1, \dots, \lambda_n)$ be a diagonal matrix. Then we have $\|A\|_2 = \|A\|_1 = \|A\|_\infty = \max(|\lambda_1|, \dots, |\lambda_n|)$ and $\|D\|_F = \sqrt{\sum_{j=1}^n |\lambda_j|^2}$.

Consider $A := \begin{bmatrix} 1000 & 999 \\ 999 & 998 \end{bmatrix}$. Then $A^{-1} = \begin{bmatrix} -998 & 999 \\ 999 & -1000 \end{bmatrix}$.

Thus $\|A\|_\infty = \|A\|_1 = \|A^{-1}\|_\infty = \|A^{-1}\|_1 = 1999$.

On the other hand,

$$\|A\|_2 = 1998.000500500375 \text{ and } \|A\|_F = 1998.000500500438.$$

Submultiplicative matrix norm

A matrix norm is said to be **submultiplicative** if

$$\|AB\| \leq \|A\| \|B\|$$

holds for all A and B . An induced matrix norm is submultiplicative.

Indeed, $\|ABx\| \leq \|A\| \|Bx\| \leq \|A\| \|B\| \|x\| \implies \|AB\| \leq \|A\| \|B\|$.

Submultiplicative matrix norm

A matrix norm is said to be **submultiplicative** if

$$\|AB\| \leq \|A\| \|B\|$$

holds for all A and B . An induced matrix norm is submultiplicative.

Indeed, $\|ABx\| \leq \|A\| \|Bx\| \leq \|A\| \|B\| \|x\| \implies \|AB\| \leq \|A\| \|B\|$.

The Frobenius norm is also submultiplicative. Indeed, suppose A is m -by- n and B is n -by- p matrices.

Submultiplicative matrix norm

A matrix norm is said to be **submultiplicative** if

$$\|AB\| \leq \|A\| \|B\|$$

holds for all A and B . An induced matrix norm is submultiplicative.

Indeed, $\|ABx\| \leq \|A\| \|Bx\| \leq \|A\| \|B\| \|x\| \implies \|AB\| \leq \|A\| \|B\|$.

The Frobenius norm is also submultiplicative. Indeed, suppose A is m -by- n and B is n -by- p matrices. Then $(AB)_{ij} = e_i^\top A B e_j$ shows that $|(AB)_{ij}| \leq \|e_i^\top A\|_2 \|B e_j\|_2$ for $i = 1 : m$ and $j = 1 : n$. Consequently,

Submultiplicative matrix norm

A matrix norm is said to be **submultiplicative** if

$$\|AB\| \leq \|A\| \|B\|$$

holds for all A and B . An induced matrix norm is submultiplicative.

Indeed, $\|ABx\| \leq \|A\| \|Bx\| \leq \|A\| \|B\| \|x\| \implies \|AB\| \leq \|A\| \|B\|$.

The Frobenius norm is also submultiplicative. Indeed, suppose A is m -by- n and B is n -by- p matrices. Then $(AB)_{ij} = e_i^\top A B e_j$ shows that $|(AB)_{ij}| \leq \|e_i^\top A\|_2 \|B e_j\|_2$ for $i = 1 : m$ and $j = 1 : n$. Consequently,

$$\begin{aligned}\|AB\|_F^2 &= \sum_{i=1}^m \sum_{j=1}^p |(AB)_{ij}|^2 \leq \sum_{i=1}^m \sum_{j=1}^p \|e_i^\top A\|_2^2 \|B e_j\|_2^2 \\ &= \sum_{i=1}^m \|e_i^\top A\|_2^2 \sum_{j=1}^p \|B e_j\|_2^2 = \|A\|_F^2 \|B\|_F^2.\end{aligned}$$

Unitarily invariant matrix norm

Let $Q \in \mathbb{R}^{n \times n}$. Then Q is said to be **orthogonal** if $Q^T Q = Q Q^T = I$.

Unitarily invariant matrix norm

Let $Q \in \mathbb{R}^{n \times n}$. Then Q is said to be **orthogonal** if $Q^T Q = Q Q^T = I$. If Q is orthogonal then

$$\|Qx\|_2 = \sqrt{\langle Qx, Qx \rangle} = \sqrt{\langle x, Q^T Qx \rangle} = \sqrt{\langle x, x \rangle} = \|x\|_2.$$

Unitarily invariant matrix norm

Let $Q \in \mathbb{R}^{n \times n}$. Then Q is said to be **orthogonal** if $Q^T Q = Q Q^T = I$. If Q is orthogonal then

$$\|Qx\|_2 = \sqrt{\langle Qx, Qx \rangle} = \sqrt{\langle x, Q^T Qx \rangle} = \sqrt{\langle x, x \rangle} = \|x\|_2.$$

Let $Q \in \mathbb{C}^{n \times n}$. Then Q is said to be **unitary** if $Q^* Q = Q Q^* = I$.

Unitarily invariant matrix norm

Let $Q \in \mathbb{R}^{n \times n}$. Then Q is said to be **orthogonal** if $Q^T Q = Q Q^T = I$. If Q is orthogonal then

$$\|Qx\|_2 = \sqrt{\langle Qx, Qx \rangle} = \sqrt{\langle x, Q^T Qx \rangle} = \sqrt{\langle x, x \rangle} = \|x\|_2.$$

Let $Q \in \mathbb{C}^{n \times n}$. Then Q is said to be **unitary** if $Q^* Q = Q Q^* = I$. If Q is unitary then $\|Qx\|_2 = \sqrt{\langle Qx, Qx \rangle} = \sqrt{\langle x, Q^* Qx \rangle} = \sqrt{\langle x, x \rangle} = \|x\|_2.$

Unitarily invariant matrix norm

Let $Q \in \mathbb{R}^{n \times n}$. Then Q is said to be **orthogonal** if $Q^T Q = Q Q^T = I$. If Q is orthogonal then

$$\|Qx\|_2 = \sqrt{\langle Qx, Qx \rangle} = \sqrt{\langle x, Q^T Qx \rangle} = \sqrt{\langle x, x \rangle} = \|x\|_2.$$

Let $Q \in \mathbb{C}^{n \times n}$. Then Q is said to be **unitary** if $Q^* Q = Q Q^* = I$. If Q is unitary then $\|Qx\|_2 = \sqrt{\langle Qx, Qx \rangle} = \sqrt{\langle x, Q^* Qx \rangle} = \sqrt{\langle x, x \rangle} = \|x\|_2$.

It follows that $\|Q\|_2 = 1$ when Q is unitary and that error in x is NOT magnified in the vector Qx .

Unitarily invariant matrix norm

Let $Q \in \mathbb{R}^{n \times n}$. Then Q is said to be **orthogonal** if $Q^T Q = Q Q^T = I$. If Q is orthogonal then

$$\|Qx\|_2 = \sqrt{\langle Qx, Qx \rangle} = \sqrt{\langle x, Q^T Qx \rangle} = \sqrt{\langle x, x \rangle} = \|x\|_2.$$

Let $Q \in \mathbb{C}^{n \times n}$. Then Q is said to be **unitary** if $Q^* Q = Q Q^* = I$. If Q is unitary then $\|Qx\|_2 = \sqrt{\langle Qx, Qx \rangle} = \sqrt{\langle x, Q^* Qx \rangle} = \sqrt{\langle x, x \rangle} = \|x\|_2$.

It follows that $\|Q\|_2 = 1$ when Q is unitary and that error in x is NOT magnified in the vector Qx .

A matrix norm is said to be **unitarily invariant** if $\|QAP\| = \|A\|$ holds for any **unitary matrices** Q and P .

Unitarily invariant matrix norm

Let $Q \in \mathbb{R}^{n \times n}$. Then Q is said to be **orthogonal** if $Q^T Q = Q Q^T = I$. If Q is orthogonal then

$$\|Qx\|_2 = \sqrt{\langle Qx, Qx \rangle} = \sqrt{\langle x, Q^T Qx \rangle} = \sqrt{\langle x, x \rangle} = \|x\|_2.$$

Let $Q \in \mathbb{C}^{n \times n}$. Then Q is said to be **unitary** if $Q^* Q = Q Q^* = I$. If Q is unitary then $\|Qx\|_2 = \sqrt{\langle Qx, Qx \rangle} = \sqrt{\langle x, Q^* Qx \rangle} = \sqrt{\langle x, x \rangle} = \|x\|_2$.

It follows that $\|Q\|_2 = 1$ when Q is unitary and that error in x is NOT magnified in the vector Qx .

A matrix norm is said to be **unitarily invariant** if $\|QAP\| = \|A\|$ holds for any **unitary matrices** Q and P . The spectral norm and the Frobenius norm are unitarily invariant norms.

Unitarily invariant matrix norm

Let $Q \in \mathbb{R}^{n \times n}$. Then Q is said to be **orthogonal** if $Q^T Q = Q Q^T = I$. If Q is orthogonal then

$$\|Qx\|_2 = \sqrt{\langle Qx, Qx \rangle} = \sqrt{\langle x, Q^T Qx \rangle} = \sqrt{\langle x, x \rangle} = \|x\|_2.$$

Let $Q \in \mathbb{C}^{n \times n}$. Then Q is said to be **unitary** if $Q^* Q = Q Q^* = I$. If Q is unitary then $\|Qx\|_2 = \sqrt{\langle Qx, Qx \rangle} = \sqrt{\langle x, Q^* Qx \rangle} = \sqrt{\langle x, x \rangle} = \|x\|_2$.

It follows that $\|Q\|_2 = 1$ when Q is unitary and that error in x is NOT magnified in the vector Qx .

A matrix norm is said to be **unitarily invariant** if $\|QAP\| = \|A\|$ holds for any **unitary matrices** Q and P . The spectral norm and the Frobenius norm are unitarily invariant norms. Indeed,

$$\begin{aligned}\|Qx\|_2 &= \|x\|_2 \Rightarrow \|QAP\|_2 = \|A\|_2. \\ \|QAP\|_F &= \sqrt{\text{Tr}((QAP)^* QAP)} = \sqrt{\text{Tr}(P^* A^* AP)} \\ &= \sqrt{\text{Tr}(A^* A)} = \|A\|_F.\end{aligned}$$