

Householder QR factorization

Rafikul Alam
Department of Mathematics
Indian Institute of Technology Guwahati
Guwahati - 781039, INDIA

Outline

- Reflector
- Householder reflector
- Householder QR factorization

Orthogonal projection

Let $P \in \mathbb{C}^{n \times n}$. Then P is called a **projection (or idempotent)** if $P^2 = P$. If P is a projection then $\mathbb{C}^n = R(P) \oplus N(P)$. Indeed, $x = Px + (I - P)x$.

Orthogonal projection

Let $P \in \mathbb{C}^{n \times n}$. Then P is called a **projection (or idempotent)** if $P^2 = P$. If P is a projection then $\mathbb{C}^n = R(P) \oplus N(P)$. Indeed, $x = Px + (I - P)x$.

A projection P is said to be an **orthogonal projection** if $R(P) \perp N(P)$. If P is an orthogonal projection then


$$\mathbb{C}^n = R(P) \oplus N(P) \text{ and } R(P) \perp N(P).$$

Orthogonal projection

Let $P \in \mathbb{C}^{n \times n}$. Then P is called a **projection (or idempotent)** if $P^2 = P$. If P is a projection then $\mathbb{C}^n = R(P) \oplus N(P)$. Indeed, $x = Px + (I - P)x$.

A projection P is said to be an **orthogonal projection** if $R(P) \perp N(P)$. If P is an orthogonal projection then

$$\mathbb{C}^n = R(P) \oplus N(P) \text{ and } R(P) \perp N(P).$$



Examples

Obviously, $P = 0$ and $P = I$ are trivial projections.

Examples

Obviously, $P = 0$ and $P = I$ are trivial projections. Let $u \in \mathbb{C}^n$ be nonzero. Then $P := \frac{uu^*}{\|u\|_2^2}$ is an orthogonal projection with $R(P) = \text{span}(u)$ and $N(P) = \{u\}^\perp$.

Examples

Obviously, $P = 0$ and $P = I$ are trivial projections. Let $u \in \mathbb{C}^n$ be nonzero. Then $P := \frac{uu^*}{\|u\|_2^2}$ is an orthogonal projection with $R(P) = \text{span}(u)$ and $N(P) = \{u\}^\perp$.

Centering projection: Set $\mathbf{e} := [1 \ \cdots \ 1]^\top \in \mathbb{C}^n$. Then $P := I - \frac{1}{n}\mathbf{e}\mathbf{e}^\top$ is an orthogonal projection. The projection P is also called a **centering matrix** as it projects a vector to a zero mean vector.

Examples

Obviously, $P = 0$ and $P = I$ are trivial projections. Let $u \in \mathbb{C}^n$ be nonzero. Then $P := \frac{uu^*}{\|u\|_2^2}$ is an orthogonal projection with $R(P) = \text{span}(u)$ and $N(P) = \{u\}^\perp$.

Centering projection: Set $\mathbf{e} := [1 \ \cdots \ 1]^\top \in \mathbb{C}^n$. Then $P := I - \frac{1}{n}\mathbf{e}\mathbf{e}^\top$ is an orthogonal projection. The projection P is also called a **centering matrix** as it projects a vector to a zero mean vector.

Indeed, let $\mathbf{x} = [x_1 \ \cdots \ x_n]^\top \in \mathbb{R}^n$ and $\mu := \frac{x_1 + \cdots + x_n}{n} = \frac{\mathbf{e}^\top \mathbf{x}}{n}$. Then

$$P\mathbf{x} = \mathbf{x} - \frac{1}{n}\mathbf{e}\mathbf{e}^\top \mathbf{x} = \mathbf{x} - \mathbf{e}\mu = [x_1 - \mu \ \cdots \ x_n - \mu]^\top.$$

Examples

Obviously, $P = 0$ and $P = I$ are trivial projections. Let $u \in \mathbb{C}^n$ be nonzero. Then $P := \frac{uu^*}{\|u\|_2^2}$ is an orthogonal projection with $R(P) = \text{span}(u)$ and $N(P) = \{u\}^\perp$.

Centering projection: Set $\mathbf{e} := [1 \ \cdots \ 1]^\top \in \mathbb{C}^n$. Then $P := I - \frac{1}{n}\mathbf{e}\mathbf{e}^\top$ is an orthogonal projection. The projection P is also called a **centering matrix** as it projects a vector to a zero mean vector.

Indeed, let $\mathbf{x} = [x_1 \ \cdots \ x_n]^\top \in \mathbb{R}^n$ and $\mu := \frac{x_1 + \cdots + x_n}{n} = \frac{\mathbf{e}^\top \mathbf{x}}{n}$. Then

$$P\mathbf{x} = \mathbf{x} - \frac{1}{n}\mathbf{e}\mathbf{e}^\top \mathbf{x} = \mathbf{x} - \mathbf{e}\mu = [x_1 - \mu \ \cdots \ x_n - \mu]^\top.$$


Note that if $\mathbf{y} = P\mathbf{x}$ then $\mu_y := \frac{y_1 + \cdots + y_n}{n} = \frac{\mathbf{e}^\top \mathbf{y}}{n} = 0$. Hence \mathbf{y} has zero mean.

Reflector

Let P be an orthogonal projection, that is, $P^2 = P$ and $R(P) \perp N(P)$.


Reflector

Let P be an orthogonal projection, that is, $P^2 = P$ and $R(P) \perp N(P)$.




Reflector

Let P be an orthogonal projection, that is, $P^2 = P$ and $R(P) \perp N(P)$.




Reflector

Let P be an orthogonal projection, that is, $P^2 = P$ and $R(P) \perp N(P)$.



Reflector

Let P be an orthogonal projection, that is, $P^2 = P$ and $R(P) \perp N(P)$.



Then $H := I - 2P$ is a reflector. Hx is a reflection of x through $N(P)$.

Householder reflector

Let $u \in \mathbb{R}^n$ be nonzero. Then $P := \frac{uu^\top}{\|u\|_2^2}$ is an orthogonal projection.

Householder reflector

Let $u \in \mathbb{R}^n$ be nonzero. Then $P := \frac{uu^\top}{\|u\|_2^2}$ is an orthogonal projection. Note that $N(P) = \{u\}^\perp$ and $R(P) = \text{span}(u)$.

Householder reflector

Let $u \in \mathbb{R}^n$ be nonzero. Then $P := \frac{uu^T}{\|u\|_2^2}$ is an orthogonal projection. Note that $N(P) = \{u\}^\perp$ and $R(P) = \text{span}(u)$. Set $\alpha := 2/\|u\|_2^2$.

Householder reflector

Let $u \in \mathbb{R}^n$ be nonzero. Then $P := \frac{uu^\top}{\|u\|_2^2}$ is an orthogonal projection. Note that $N(P) = \{u\}^\perp$ and $R(P) = \text{span}(u)$. Set $\alpha := 2/\|u\|_2^2$.

The reflector $H := I - 2P = I - \frac{2uu^\top}{\|u\|_2^2} = I - \alpha uu^\top$ is called a **Householder reflector** and u is called a **Householder vector**.

Householder reflector

Let $u \in \mathbb{R}^n$ be nonzero. Then $P := \frac{uu^\top}{\|u\|_2^2}$ is an orthogonal projection. Note that $N(P) = \{u\}^\perp$ and $R(P) = \text{span}(u)$. Set $\alpha := 2/\|u\|_2^2$.

The reflector $H := I - 2P = I - \frac{2uu^\top}{\|u\|_2^2} = I - \alpha uu^\top$ is called a **Householder reflector** and u is called a **Householder vector**.

Remark: The reflector $H := I - \alpha uu^\top$ is completely determined by the Householder vector u .

Householder reflector

Let $u \in \mathbb{R}^n$ be nonzero. Then $P := \frac{uu^\top}{\|u\|_2^2}$ is an orthogonal projection. Note that $N(P) = \{u\}^\perp$ and $R(P) = \text{span}(u)$. Set $\alpha := 2/\|u\|_2^2$.

The reflector $H := I - 2P = I - \frac{2uu^\top}{\|u\|_2^2} = I - \alpha uu^\top$ is called a **Householder reflector** and u is called a **Householder vector**.

Remark: The reflector $H := I - \alpha uu^\top$ is completely determined by the Householder vector u . Often we write H as $H = (\alpha, u)$.

Householder reflector

Let $u \in \mathbb{R}^n$ be nonzero. Then $P := \frac{uu^\top}{\|u\|_2^2}$ is an orthogonal projection. Note that $N(P) = \{u\}^\perp$ and $R(P) = \text{span}(u)$. Set $\alpha := 2/\|u\|_2^2$.

The reflector $H := I - 2P = I - \frac{2uu^\top}{\|u\|_2^2} = I - \alpha uu^\top$ is called a **Householder reflector** and u is called a **Householder vector**.

Remark: The reflector $H := I - \alpha uu^\top$ is completely determined by the Householder vector u . Often we write H as $H = (\alpha, u)$.

Fact: Let $H := I - \alpha uu^\top$ be a reflector. Then the following hold:

- $Hu = -u$ and $Hx = x$ for $x \perp u$.

Householder reflector

Let $u \in \mathbb{R}^n$ be nonzero. Then $P := \frac{uu^\top}{\|u\|_2^2}$ is an orthogonal projection. Note that $N(P) = \{u\}^\perp$ and $R(P) = \text{span}(u)$. Set $\alpha := 2/\|u\|_2^2$.

The reflector $H := I - 2P = I - \frac{2uu^\top}{\|u\|_2^2} = I - \alpha uu^\top$ is called a **Householder reflector** and u is called a **Householder vector**.

Remark: The reflector $H := I - \alpha uu^\top$ is completely determined by the Householder vector u . Often we write H as $H = (\alpha, u)$.

Fact: Let $H := I - \alpha uu^\top$ be a reflector. Then the following hold:

- $Hu = -u$ and $Hx = x$ for $x \perp u$.
- $H^\top = H$ and $H^2 = I$. Hence H is unitary and $H = H^{-1}$.

Householder reflector


Let $u \in \mathbb{R}^n$ be nonzero. Then $P := \frac{uu^\top}{\|u\|_2^2}$ is an orthogonal projection. Note that $N(P) = \{u\}^\perp$ and $R(P) = \text{span}(u)$. Set $\alpha := 2/\|u\|_2^2$.

The reflector $H := I - 2P = I - \frac{2uu^\top}{\|u\|_2^2} = I - \alpha uu^\top$ is called a **Householder reflector** and u is called a **Householder vector**.

Remark: The reflector $H := I - \alpha uu^\top$ is completely determined by the Householder vector u . Often we write H as $H = (\alpha, u)$.

Fact: Let $H := I - \alpha uu^\top$ be a reflector. Then the following hold:

- $Hu = -u$ and $Hx = x$ for $x \perp u$.
- $H^\top = H$ and $H^2 = I$. Hence H is unitary and $H = H^{-1}$.
- For $x \in \mathbb{R}^n$, we have $Hx = x - (\alpha u^\top x)u$. Hence computation of Hx costs only $4n$ flops.



Householder reflector

Theorem: Let $x, y \in \mathbb{R}^n$ be such that $x \neq y$ and $\|x\|_2 = \|y\|_2$.


Householder reflector

Theorem: Let $x, y \in \mathbb{R}^n$ be such that $x \neq y$ and $\|x\|_2 = \|y\|_2$. Set $u := x - y$ and $\alpha := 2/\|u\|_2^2$. Then $H := I - \alpha uu^T$ is a unique reflector such that $Hx = y$.

Householder reflector

Theorem: Let $x, y \in \mathbb{R}^n$ be such that $x \neq y$ and $\|x\|_2 = \|y\|_2$. Set $u := x - y$ and $\alpha := 2/\|u\|_2^2$. Then $H := I - \alpha uu^\top$ is a unique reflector such that $Hx = y$.


Proof: Note that $(x + y) \perp (x - y)$.



Householder reflector

Theorem: Let $x, y \in \mathbb{R}^n$ be such that $x \neq y$ and $\|x\|_2 = \|y\|_2$. Set $u := x - y$ and $\alpha := 2/\|u\|_2^2$. Then $H := I - \alpha uu^\top$ is a unique reflector such that $Hx = y$.

Proof: Note that $(x + y) \perp (x - y)$.




Note that $u \perp (x + y)$. Hence $Hu = -u$ and $H(x + y) = x + y$.

Householder reflector

Theorem: Let $x, y \in \mathbb{R}^n$ be such that $x \neq y$ and $\|x\|_2 = \|y\|_2$. Set $u := x - y$ and $\alpha := 2/\|u\|_2^2$. Then $H := I - \alpha uu^\top$ is a unique reflector such that $Hx = y$.

Proof: Note that $(x + y) \perp (x - y)$.



Note that $u \perp (x + y)$. Hence $Hu = -u$ and $H(x + y) = x + y$. Now $x = \frac{1}{2}(u + x + y) \Rightarrow Hx = \frac{1}{2}(Hu + H(x + y)) = \frac{1}{2}(-u + x + y) = y$. ■

Creating zeros via Householder reflector

Let $x := [x_1 \ x_2 \ \cdots \ x_n]^\top \in \mathbb{R}^n$. Define $y := [-\sigma \ 0 \ \cdots \ 0]^\top \in \mathbb{R}^n$, where $\sigma := \text{sign}(x_1)\|x\|_2$. Set $u := [x_1 + \sigma \ x_2 \ \cdots \ x_n]^\top$.

Creating zeros via Householder reflector

Let $x := [x_1 \ x_2 \ \cdots \ x_n]^\top \in \mathbb{R}^n$. Define $y := [-\sigma \ 0 \ \cdots \ 0]^\top \in \mathbb{R}^n$, where $\sigma := \text{sign}(x_1)\|x\|_2$. Set $u := [x_1 + \sigma \ x_2 \ \cdots \ x_n]^\top$. Then $H := I - \alpha uu^\top$ is a unique reflector such that $Hx = y$ with $\alpha = 2/\|u\|_2^2 = 1/\sigma(x_1 + \sigma) = 1/\sigma u_1$, where $u_1 := e_1^\top u$.

Creating zeros via Householder reflector

Let $x := [x_1 \ x_2 \ \cdots \ x_n]^\top \in \mathbb{R}^n$. Define $y := [-\sigma \ 0 \ \cdots \ 0]^\top \in \mathbb{R}^n$, where $\sigma := \text{sign}(x_1)\|x\|_2$. Set $u := [x_1 + \sigma \ x_2 \ \cdots \ x_n]^\top$. Then $H := I - \alpha uu^\top$ is a unique reflector such that $Hx = y$ with $\alpha = 2/\|u\|_2^2 = 1/\sigma(x_1 + \sigma) = 1/\sigma u_1$, where $u_1 := e_1^\top u$.

Indeed, $\|u\|_2^2 = (x_1 + \sigma)^2 + x_2^2 + \cdots + x_n^2 = 2\sigma^2 + 2x_1\sigma = 2\sigma(x_1 + \sigma)$. Hence $\alpha = 2/\|u\|_2^2 = 1/\sigma(x_1 + \sigma) = 1/\sigma u_1$.

Creating zeros via Householder reflector

Let $x := [x_1 \ x_2 \ \cdots \ x_n]^\top \in \mathbb{R}^n$. Define $y := [-\sigma \ 0 \ \cdots \ 0]^\top \in \mathbb{R}^n$, where $\sigma := \text{sign}(x_1)\|x\|_2$. Set $u := [x_1 + \sigma \ x_2 \ \cdots \ x_n]^\top$. Then $H := I - \alpha uu^\top$ is a unique reflector such that $Hx = y$ with $\alpha = 2/\|u\|_2^2 = 1/\sigma(x_1 + \sigma) = 1/\sigma u_1$, where $u_1 := e_1^\top u$.

Indeed, $\|u\|_2^2 = (x_1 + \sigma)^2 + x_2^2 + \cdots + x_n^2 = 2\sigma^2 + 2x_1\sigma = 2\sigma(x_1 + \sigma)$. Hence $\alpha = 2/\|u\|_2^2 = 1/\sigma(x_1 + \sigma) = 1/\sigma u_1$. By construction, $Hx = y$.

Creating zeros via Householder reflector

Let $x := [x_1 \ x_2 \ \cdots \ x_n]^\top \in \mathbb{R}^n$. Define $y := [-\sigma \ 0 \ \cdots \ 0]^\top \in \mathbb{R}^n$, where $\sigma := \text{sign}(x_1)\|x\|_2$. Set $u := [x_1 + \sigma \ x_2 \ \cdots \ x_n]^\top$. Then $H := I - \alpha uu^\top$ is a unique reflector such that $Hx = y$ with $\alpha = 2/\|u\|_2^2 = 1/\sigma(x_1 + \sigma) = 1/\sigma u_1$, where $u_1 := e_1^\top u$.

Indeed, $\|u\|_2^2 = (x_1 + \sigma)^2 + x_2^2 + \cdots + x_n^2 = 2\sigma^2 + 2x_1\sigma = 2\sigma(x_1 + \sigma)$. Hence $\alpha = 2/\|u\|_2^2 = 1/\sigma(x_1 + \sigma) = 1/\sigma u_1$. By construction, $Hx = y$.

Remark: The choice of σ ensures that computation of $x_1 + \sigma$ avoids subtraction and hence catastrophic cancellation.

Creating zeros via Householder reflector

Let $x := [x_1 \ x_2 \ \cdots \ x_n]^\top \in \mathbb{R}^n$. Define $y := [-\sigma \ 0 \ \cdots \ 0]^\top \in \mathbb{R}^n$, where $\sigma := \text{sign}(x_1)\|x\|_2$. Set $u := [x_1 + \sigma \ x_2 \ \cdots \ x_n]^\top$. Then $H := I - \alpha uu^\top$ is a unique reflector such that $Hx = y$ with $\alpha = 2/\|u\|_2^2 = 1/\sigma(x_1 + \sigma) = 1/\sigma u_1$, where $u_1 := e_1^\top u$.

Indeed, $\|u\|_2^2 = (x_1 + \sigma)^2 + x_2^2 + \cdots + x_n^2 = 2\sigma^2 + 2x_1\sigma = 2\sigma(x_1 + \sigma)$. Hence $\alpha = 2/\|u\|_2^2 = 1/\sigma(x_1 + \sigma) = 1/\sigma u_1$. By construction, $Hx = y$.

Remark: The choice of σ ensures that computation of $x_1 + \sigma$ avoids subtraction and hence catastrophic cancellation.

Note that we only need to store u and σ for constructing $H := I - \alpha uu^\top$. Hence construction of H requires only $2n$ flops.

Creating zeros via Householder reflector

Let $x := [x_1 \ x_2 \ \cdots \ x_n]^\top \in \mathbb{R}^n$. Define $y := [-\sigma \ 0 \ \cdots \ 0]^\top \in \mathbb{R}^n$, where $\sigma := \text{sign}(x_1)\|x\|_2$. Set $u := [x_1 + \sigma \ x_2 \ \cdots \ x_n]^\top$. Then $H := I - \alpha uu^\top$ is a unique reflector such that $Hx = y$ with $\alpha = 2/\|u\|_2^2 = 1/\sigma(x_1 + \sigma) = 1/\sigma u_1$, where $u_1 := e_1^\top u$.

Indeed, $\|u\|_2^2 = (x_1 + \sigma)^2 + x_2^2 + \cdots + x_n^2 = 2\sigma^2 + 2x_1\sigma = 2\sigma(x_1 + \sigma)$. Hence $\alpha = 2/\|u\|_2^2 = 1/\sigma(x_1 + \sigma) = 1/\sigma u_1$. By construction, $Hx = y$.

Remark: The choice of σ ensures that computation of $x_1 + \sigma$ avoids subtraction and hence catastrophic cancellation.

Note that we only need to store u and σ for constructing $H := I - \alpha uu^\top$. Hence construction of H requires only $2n$ flops.

For a vector $w \in \mathbb{R}^n$, we have $Hw = w - (\alpha u^\top w)u$. Hence computation of Hw requires $4n$ flops instead of $2n^2$ flops.

Creating zeros via Householder reflector

Let $x := [x_1 \ x_2 \ \cdots \ x_n]^\top \in \mathbb{R}^n$. Define $y := [-\sigma \ 0 \ \cdots \ 0]^\top \in \mathbb{R}^n$, where $\sigma := \text{sign}(x_1)\|x\|_2$. Set $u := [x_1 + \sigma \ x_2 \ \cdots \ x_n]^\top$. Then $H := I - \alpha uu^\top$ is a unique reflector such that $Hx = y$ with $\alpha = 2/\|u\|_2^2 = 1/\sigma(x_1 + \sigma) = 1/\sigma u_1$, where $u_1 := e_1^\top u$.

Indeed, $\|u\|_2^2 = (x_1 + \sigma)^2 + x_2^2 + \cdots + x_n^2 = 2\sigma^2 + 2x_1\sigma = 2\sigma(x_1 + \sigma)$. Hence $\alpha = 2/\|u\|_2^2 = 1/\sigma(x_1 + \sigma) = 1/\sigma u_1$. By construction, $Hx = y$.

Remark: The choice of σ ensures that computation of $x_1 + \sigma$ avoids subtraction and hence catastrophic cancellation.

Note that we only need to store u and σ for constructing $H := I - \alpha uu^\top$. Hence construction of H requires only $2n$ flops.

For a vector $w \in \mathbb{R}^n$, we have $Hw = w - (\alpha u^\top w)u$. Hence computation of Hw requires $4n$ flops instead of $2n^2$ flops.

For a matrix $A \in \mathbb{R}^{n \times p}$, we have $HA = A - \alpha uu^\top A = A - \alpha u(A^\top u)^\top$.

Creating zeros via Householder reflector

Let $x := [x_1 \ x_2 \ \cdots \ x_n]^\top \in \mathbb{R}^n$. Define $y := [-\sigma \ 0 \ \cdots \ 0]^\top \in \mathbb{R}^n$, where $\sigma := \text{sign}(x_1)\|x\|_2$. Set $u := [x_1 + \sigma \ x_2 \ \cdots \ x_n]^\top$. Then $H := I - \alpha uu^\top$ is a unique reflector such that $Hx = y$ with $\alpha = 2/\|u\|_2^2 = 1/\sigma(x_1 + \sigma) = 1/\sigma u_1$, where $u_1 := e_1^\top u$.

Indeed, $\|u\|_2^2 = (x_1 + \sigma)^2 + x_2^2 + \cdots + x_n^2 = 2\sigma^2 + 2x_1\sigma = 2\sigma(x_1 + \sigma)$. Hence $\alpha = 2/\|u\|_2^2 = 1/\sigma(x_1 + \sigma) = 1/\sigma u_1$. By construction, $Hx = y$.

Remark: The choice of σ ensures that computation of $x_1 + \sigma$ avoids subtraction and hence catastrophic cancellation.

Note that we only need to store u and σ for constructing $H := I - \alpha uu^\top$. Hence construction of H requires only $2n$ flops.

For a vector $w \in \mathbb{R}^n$, we have $Hw = w - (\alpha u^\top w)u$. Hence computation of Hw requires $4n$ flops instead of $2n^2$ flops.



For a matrix $A \in \mathbb{R}^{n \times p}$, we have $HA = A - \alpha uu^\top A = A - \alpha u(A^\top u)^\top$. Hence the cost of computing HA is $4np$ flops instead of $2n^2p$ flops.

Creating zeros via Householder reflector

Let $x := [x_1 \ x_2 \ \cdots \ x_n]^\top \in \mathbb{R}^n$ and $y := [x_1 \ \cdots x_{i-1} \ -\sigma_i \ 0 \ \cdots \ 0]^\top$, where $\sigma_i := \text{sign}(x_i) \sqrt{x_i^2 + \cdots + x_n^2}$. Then $\|x\|_2 = \|y\|_2$ and $x \neq y$.

Creating zeros via Householder reflector

Let $x := [x_1 \ x_2 \ \cdots \ x_n]^\top \in \mathbb{R}^n$ and $y := [x_1 \ \cdots x_{i-1} \ -\sigma_i \ 0 \ \cdots \ 0]^\top$, where $\sigma_i := \text{sign}(x_i) \sqrt{x_i^2 + \cdots + x_n^2}$. Then $\|x\|_2 = \|y\|_2$ and $x \neq y$.

Set $u := x - y = [0 \ \cdots \ 0 \ x_i + \sigma_i \ x_{i+1} \ \cdots \ x_n]^\top$. Then $H := I - \alpha uu^\top$ is a unique reflector such that $Hx = y$, where $\alpha = 2/\|u\|_2^2 = 1/\sigma_i(x_i + \sigma_i) = 1/\sigma_i u_i$.

Creating zeros via Householder reflector

Let $x := [x_1 \ x_2 \ \cdots \ x_n]^\top \in \mathbb{R}^n$ and $y := [x_1 \ \cdots x_{i-1} \ -\sigma_i \ 0 \ \cdots \ 0]^\top$, where $\sigma_i := \text{sign}(x_i) \sqrt{x_i^2 + \cdots + x_n^2}$. Then $\|x\|_2 = \|y\|_2$ and $x \neq y$.

Set $u := x - y = [0 \ \cdots \ 0 \ x_i + \sigma_i \ x_{i+1} \ \cdots \ x_n]^\top$. Then $H := I - \alpha uu^\top$ is a unique reflector such that $Hx = y$, where $\alpha = 2/\|u\|_2^2 = 1/\sigma_i(x_i + \sigma_i) = 1/\sigma_i u_i$.

Set $\hat{u} := [x_i + \sigma_i \ x_{i+1} \ \cdots \ x_n]^\top$. Then $u = \begin{bmatrix} 0 \\ \hat{u} \end{bmatrix}$ and

$$H = I - \alpha uu^\top = \left[\begin{array}{c|c} I_{i-1} & 0 \\ \hline 0 & I_{n-i+1} - \alpha \hat{u} \hat{u}^\top \end{array} \right] = \left[\begin{array}{c|c} I_{i-1} & 0 \\ \hline 0 & \hat{H} \end{array} \right].$$

Creating zeros via Householder reflector

Let $x := [x_1 \ x_2 \ \cdots \ x_n]^\top \in \mathbb{R}^n$ and $y := [x_1 \ \cdots x_{i-1} \ -\sigma_i \ 0 \ \cdots \ 0]^\top$, where $\sigma_i := \text{sign}(x_i) \sqrt{x_i^2 + \cdots + x_n^2}$. Then $\|x\|_2 = \|y\|_2$ and $x \neq y$.

Set $u := x - y = [0 \ \cdots \ 0 \ x_i + \sigma_i \ x_{i+1} \ \cdots \ x_n]^\top$. Then $H := I - \alpha uu^\top$ is a unique reflector such that $Hx = y$, where $\alpha = 2/\|u\|_2^2 = 1/\sigma_i(x_i + \sigma_i) = 1/\sigma_i u_i$.

Set $\hat{u} := [x_i + \sigma_i \ x_{i+1} \ \cdots \ x_n]^\top$. Then $u = \begin{bmatrix} 0 \\ \hat{u} \end{bmatrix}$ and

$$H = I - \alpha uu^\top = \left[\begin{array}{c|c} I_{i-1} & 0 \\ \hline 0 & I_{n-i+1} - \alpha \hat{u} \hat{u}^\top \end{array} \right] = \left[\begin{array}{c|c} I_{i-1} & 0 \\ \hline 0 & \hat{H} \end{array} \right].$$

This shows that for $w \in \mathbb{R}^n$, the first $i - 1$ components of Hw remain unchanged.

Creating zeros via Householder reflector

Let $x := [x_1 \ x_2 \ \cdots \ x_n]^\top \in \mathbb{R}^n$ and $y := [x_1 \ \cdots x_{i-1} \ -\sigma_i \ 0 \ \cdots \ 0]^\top$, where $\sigma_i := \text{sign}(x_i) \sqrt{x_i^2 + \cdots + x_n^2}$. Then $\|x\|_2 = \|y\|_2$ and $x \neq y$.

Set $u := x - y = [0 \ \cdots \ 0 \ x_i + \sigma_i \ x_{i+1} \ \cdots \ x_n]^\top$. Then $H := I - \alpha uu^\top$ is a unique reflector such that $Hx = y$, where $\alpha = 2/\|u\|_2^2 = 1/\sigma_i(x_i + \sigma_i) = 1/\sigma_i u_i$.

Set $\hat{u} := [x_i + \sigma_i \ x_{i+1} \ \cdots \ x_n]^\top$. Then $u = \begin{bmatrix} 0 \\ \hat{u} \end{bmatrix}$ and

$$H = I - \alpha uu^\top = \left[\begin{array}{c|c} I_{i-1} & 0 \\ \hline 0 & I_{n-i+1} - \alpha \hat{u} \hat{u}^\top \end{array} \right] = \left[\begin{array}{c|c} I_{i-1} & 0 \\ \hline 0 & \hat{H} \end{array} \right].$$

This shows that for $w \in \mathbb{R}^n$, the first $i - 1$ components of Hw remain unchanged. Similarly, for $A \in \mathbb{R}^{n \times n}$, the first $i - 1$ rows of HA and the first $i - 1$ columns of AH remain unchanged.

Householder QR factorization

Construct $H_1 := I - \alpha_1 u_1 u_1^\top$ such that $H_1 A e_1 = [-\sigma_1, 0, \dots, 0]^\top$. Then

Householder QR factorization

Construct $H_1 := I - \alpha_1 u_1 u_1^\top$ such that $H_1 A e_1 = [-\sigma_1, 0, \dots, 0]^\top$. Then

$$H_1 A = \left[\begin{array}{c|ccc} -\sigma_1 & a_{12}^{(1)} & \cdots & a_{1n}^{(1)} \\ \hline 0 & a_{22}^{(1)} & \cdots & a_{2n}^{(1)} \\ \vdots & \vdots & & \vdots \\ 0 & a_{m2}^{(1)} & \cdots & a_{mn}^{(1)} \end{array} \right].$$

Householder QR factorization

Construct $H_1 := I - \alpha_1 u_1 u_1^\top$ such that $H_1 A e_1 = [-\sigma_1, 0, \dots, 0]^\top$. Then

$$H_1 A = \left[\begin{array}{c|ccc} -\sigma_1 & a_{12}^{(1)} & \cdots & a_{1n}^{(1)} \\ \hline 0 & a_{22}^{(1)} & \cdots & a_{2n}^{(1)} \\ \vdots & \vdots & & \vdots \\ 0 & a_{m2}^{(1)} & \cdots & a_{mn}^{(1)} \end{array} \right].$$

Construct $H_2 := I - \alpha_2 u_2 u_2^\top$ such that $H_2 H_1 A e_2 = [a_{12}^{(1)}, -\sigma_2, 0, \dots, 0]^\top$.

Householder QR factorization

Construct $H_1 := I - \alpha_1 u_1 u_1^\top$ such that $H_1 A e_1 = [-\sigma_1, 0, \dots, 0]^\top$. Then

$$H_1 A = \left[\begin{array}{c|ccc} -\sigma_1 & a_{12}^{(1)} & \cdots & a_{1n}^{(1)} \\ \hline 0 & a_{22}^{(1)} & \cdots & a_{2n}^{(1)} \\ \vdots & \vdots & & \vdots \\ 0 & a_{m2}^{(1)} & \cdots & a_{mn}^{(1)} \end{array} \right].$$

Construct $H_2 := I - \alpha_2 u_2 u_2^\top$ such that $H_2 H_1 A e_2 = [a_{12}^{(1)}, -\sigma_2, 0, \dots, 0]^\top$.

$$H_2 H_1 A = \left[\begin{array}{c|c|ccc} -\sigma_1 & a_{12}^{(1)} & a_{13}^{(1)} & \cdots & a_{1n}^{(1)} \\ \hline 0 & -\sigma_2 & a_{23}^{(2)} & \cdots & a_{2n}^{(2)} \\ \hline 0 & 0 & a_{33}^{(2)} & \cdots & a_{3n}^{(2)} \\ \vdots & \vdots & \vdots & & \vdots \\ 0 & 0 & a_{m3}^{(2)} & \cdots & a_{mn}^{(2)} \end{array} \right].$$

Householder QR factorization

Construct $H_1 := I - \alpha_1 u_1 u_1^\top$ such that $H_1 A e_1 = [-\sigma_1, 0, \dots, 0]^\top$. Then

$$H_1 A = \left[\begin{array}{c|ccc} -\sigma_1 & a_{12}^{(1)} & \cdots & a_{1n}^{(1)} \\ \hline 0 & a_{22}^{(1)} & \cdots & a_{2n}^{(1)} \\ \vdots & \vdots & & \vdots \\ 0 & a_{m2}^{(1)} & \cdots & a_{mn}^{(1)} \end{array} \right].$$

Construct $H_2 := I - \alpha_2 u_2 u_2^\top$ such that $H_2 H_1 A e_2 = [a_{12}^{(1)}, -\sigma_2, 0, \dots, 0]^\top$.

$$H_2 H_1 A = \left[\begin{array}{c|c|ccc} -\sigma_1 & a_{12}^{(1)} & a_{13}^{(1)} & \cdots & a_{1n}^{(1)} \\ \hline 0 & -\sigma_2 & a_{23}^{(2)} & \cdots & a_{2n}^{(2)} \\ \hline 0 & 0 & a_{33}^{(2)} & \cdots & a_{3n}^{(2)} \\ \vdots & \vdots & \vdots & & \vdots \\ 0 & 0 & a_{m3}^{(2)} & \cdots & a_{mn}^{(2)} \end{array} \right].$$

Hence we have $H_n \cdots H_2 H_1 A = R \Rightarrow A = H_1 H_2 \cdots H_n R = QR$.

Algorithm (Householder QR)

Input: An $m \times n$ matrix A .

Output: QR factorization $A = QR$.

Algorithm (Householder QR)

Input: An $m \times n$ matrix A .

Output: QR factorization $A = QR$.

For $k = 1 : \min(m - 1, n)$

- Construct $H_k = I - \alpha_k u_k u_k^\top$ such that $(H_k A)(k : m, k) = [-\sigma_k, 0, \dots, 0]^\top$.

Algorithm (Householder QR)

Input: An $m \times n$ matrix A .

Output: QR factorization $A = QR$.

For $k = 1 : \min(m - 1, n)$

- Construct $H_k = I - \alpha_k u_k u_k^\top$ such that $(H_k A)(k : m, k) = [-\sigma_k, 0, \dots, 0]^\top$.
- Compute $A \leftarrow H_k A = A - \alpha u_k (u_k^\top A)$.

end

Algorithm (Householder QR)

Input: An $m \times n$ matrix A .

Output: QR factorization $A = QR$.

For $k = 1 : \min(m - 1, n)$

- Construct $H_k = I - \alpha_k u_k u_k^\top$ such that $(H_k A)(k : m, k) = [-\sigma_k, 0, \dots, 0]^\top$.
- Compute $A \leftarrow H_k A = A - \alpha u_k (u_k^\top A)$.

end

Since $H_k = \begin{bmatrix} I_{k-1} & 0 \\ 0 & \hat{H}_k \end{bmatrix}$, $A \leftarrow H_k A$ performs transformation on $A(k : m, k : n)$, which dominates the computation and requires $4(m - k)(n - k)$ flops.

Algorithm (Householder QR)

Input: An $m \times n$ matrix A .

Output: QR factorization $A = QR$.

For $k = 1 : \min(m - 1, n)$

- Construct $H_k = I - \alpha_k u_k u_k^\top$ such that $(H_k A)(k : m, k) = [-\sigma_k, 0, \dots, 0]^\top$.
- Compute $A \leftarrow H_k A = A - \alpha u_k (u_k^\top A)$.

end

Since $H_k = \begin{bmatrix} I_{k-1} & 0 \\ 0 & \hat{H}_k \end{bmatrix}$, $A \leftarrow H_k A$ performs transformation on $A(k : m, k : n)$, which dominates the computation and requires $4(m - k)(n - k)$ flops. Hence the total cost is

$$\sum_{k=1}^n 4(m - k)(n - k) = 4 \int_0^n (m - x)(n - x) dx = 2mn^2 - 2n^3/3 \text{ flops.}$$

A MATLAB function for Householder QR

```
function [Q, R] = hqr(A)
% [Q R] = hqr(A) produces a unitary matrix Q and
% an triangular matrix R so that  $A = QR$ .
```

A MATLAB function for Householder QR

```
function [Q, R] = hqr(A)
% [Q R] = hqr(A) produces a unitary matrix Q and
% an triangular matrix R so that A= QR.
[m, n] = size(A); Q = eye(m);
```

A MATLAB function for Householder QR

```
function [Q, R] = hqr(A)
% [Q R] = hqr(A) produces a unitary matrix Q and
% an triangular matrix R so that A= QR.

[m, n] = size(A); Q = eye(m);

for k = 1:min(m-1, n)

    % compute Householder vector for k-th step
    u = A(k:m,k); sigma = norm(u);
```


A MATLAB function for Householder QR

```
function [Q, R] = hqr(A)
% [Q R] = hqr(A) produces a unitary matrix Q and
% an triangular matrix R so that A= QR.

[m, n] = size(A); Q = eye(m);

for k = 1:min(m-1, n)

    % compute Householder vector for k-th step
    u = A(k:m,k); sigma = norm(u);

    % Skip if u = 0, i.e., column is already zero.
    if sigma ~= 0
        if u(1) ~= 0, sigma = sign(u(1))*sigma; end
```

A MATLAB function for Householder QR

```
function [Q, R] = hqr(A)
% [Q R] = hqr(A) produces a unitary matrix Q and
% an triangular matrix R so that A= QR.

[m, n] = size(A); Q = eye(m);

for k = 1:min(m-1, n)

    % compute Householder vector for k-th step
    u = A(k:m,k); sigma = norm(u);

    % Skip if u = 0, i.e., column is already zero.
    if sigma ~= 0
        if u(1) ~= 0, sigma = sign(u(1))*sigma; end
        u(1) = u(1) + sigma;
        alpha = 1/(conj(sigma)*u(1));
```

A MATLAB function for Householder QR

```
function [Q, R] = hqr(A)
% [Q R] = hqr(A) produces a unitary matrix Q and
% an triangular matrix R so that A= QR.

[m, n] = size(A); Q = eye(m);
for k = 1:min(m-1, n)

    % compute Householder vector for k-th step
    u = A(k:m,k); sigma = norm(u);
    % Skip if u = 0, i.e., column is already zero.
    if sigma ~= 0
        if u(1) ~= 0, sigma = sign(u(1))*sigma; end
        u(1) = u(1) + sigma;
        alpha = 1/(conj(sigma)*u(1));

        % Update the k-th column.
        A(k:m,k) = 0; A(k,k) = -sigma;
```

A MATLAB function for Householder QR

```
% update A(k:m,k+1:n) and assemble Q  
v = u'*A(k:m, k+1:n);  
w = u'*Q(k:m, k:n);
```

A MATLAB function for Householder QR

```
% update A(k:m,k+1:n) and assemble Q
v = u'*A(k:m, k+1:n);
w = u'*Q(k:m, k:n);

A(k:m, k+1:n) = A(k:m,k+1:n)- (alpha * u) * v;
Q(k:m, k:m) = Q(k:m, k:m)- (alpha * u) * w;
end % end of if

end % end of for loop
```

A MATLAB function for Householder QR

```
% update A(k:m,k+1:n) and assemble Q
v = u'*A(k:m, k+1:n);
w = u'*Q(k:m, k:n);

A(k:m, k+1:n) = A(k:m,k+1:n)- (alpha * u) * v;
Q(k:m, k:m) = Q(k:m, k:m)- (alpha * u) * w;
end % end of if

end % end of for loop
R = A;
```

A MATLAB function for Householder QR

```
% update A(k:m,k+1:n) and assemble Q
v = u'*A(k:m, k+1:n);
w = u'*Q(k:m, k:n);

A(k:m, k+1:n) = A(k:m,k+1:n)- (alpha * u) * v;
Q(k:m, k:m) = Q(k:m, k:m)- (alpha * u) * w;
end % end of if

end % end of for loop
R = A;
```

Remark: The matrix Q is usually not required and hence is never assembled. It is more efficient to store the Householder vectors for computing Qb for any vector b .

A MATLAB function for Householder QR

```
% update A(k:m,k+1:n) and assemble Q
v = u'*A(k:m, k+1:n);
w = u'*Q(k:m, k:n);

A(k:m, k+1:n) = A(k:m,k+1:n)- (alpha * u) * v;
Q(k:m, k:m) = Q(k:m, k:m)- (alpha * u) * w;
end % end of if

end % end of for loop
R = A;
```

Remark: The matrix Q is usually not required and hence is never assembled. It is more efficient to store the Householder vectors for computing Qb for any vector b .

The MATLAB function `hqr` can be modified to store the Householder vectors in the lower triangular part of A and $-\sigma_1, \dots, -\sigma_n$ can be stored in an additional array `sigma`.

A MATLAB function for Householder QR

```
% update A(k:m,k+1:n) and assemble Q
v = u'*A(k:m, k+1:n);
w = u'*Q(k:m, k:n);

A(k:m, k+1:n) = A(k:m,k+1:n)- (alpha * u) * v;
Q(k:m, k:m) = Q(k:m, k:m)- (alpha * u) * w;
end % end of if

end % end of for loop
R = A;
```

Remark: The matrix Q is usually not required and hence is never assembled. It is more efficient to store the Householder vectors for computing Qb for any vector b .

The MATLAB function `hqr` can be modified to store the Householder vectors in the lower triangular part of A and $-\sigma_1, \dots, -\sigma_n$ can be stored in an additional array `sigma`. In such a case,

$$R = \text{diag}(\text{sigma}) + \text{triu}(A, 1).$$

Solution of LSP by Householder QR factorization

Let $A \in \mathbb{R}^{m \times n}$, $m > n$, and $b \in \mathbb{R}^m$. Suppose that $\text{rank}(A) = n$.

Solution of LSP by Householder QR factorization

Let $A \in \mathbb{R}^{m \times n}$, $m > n$, and $b \in \mathbb{R}^m$. Suppose that $\text{rank}(A) = n$. Then the LSP $Ax = b$ can be solved in three steps:

Solution of LSP by Householder QR factorization

Let $A \in \mathbb{R}^{m \times n}$, $m > n$, and $b \in \mathbb{R}^m$. Suppose that $\text{rank}(A) = n$. Then the LSP $Ax = b$ can be solved in three steps:

- Compute QR factorization $[A, b] = QR$.

Solution of LSP by Householder QR factorization

Let $A \in \mathbb{R}^{m \times n}$, $m > n$, and $b \in \mathbb{R}^m$. Suppose that $\text{rank}(A) = n$. Then the LSP $Ax = b$ can be solved in three steps:

- Compute QR factorization $[A, b] = QR$. Cost: $2mn^2 - \frac{2n^3}{3}$ flops.

Solution of LSP by Householder QR factorization

Let $A \in \mathbb{R}^{m \times n}$, $m > n$, and $b \in \mathbb{R}^m$. Suppose that $\text{rank}(A) = n$. Then the LSP $Ax = b$ can be solved in three steps:

- Compute QR factorization $[A, b] = QR$. Cost: $2mn^2 - \frac{2n^3}{3}$ flops.
- Solve the upper triangular system $R(1:n, 1:n)x = R(1:n, n+1)$.

Solution of LSP by Householder QR factorization

Let $A \in \mathbb{R}^{m \times n}$, $m > n$, and $b \in \mathbb{R}^m$. Suppose that $\text{rank}(A) = n$. Then the LSP $Ax = b$ can be solved in three steps:

- Compute QR factorization $[A, b] = QR$. Cost: $2mn^2 - \frac{2n^3}{3}$ flops.
- Solve the upper triangular system $R(1:n, 1:n)x = R(1:n, n+1)$.
Cost: n^2 flops.

Solution of LSP by Householder QR factorization

Let $A \in \mathbb{R}^{m \times n}$, $m > n$, and $b \in \mathbb{R}^m$. Suppose that $\text{rank}(A) = n$. Then the LSP $Ax = b$ can be solved in three steps:

- Compute QR factorization $[A, b] = QR$. Cost: $2mn^2 - \frac{2n^3}{3}$ flops.
- Solve the upper triangular system $R(1:n, 1:n)x = R(1:n, n+1)$.
Cost: n^2 flops.
- Compute residual norm $d = \text{abs}(R(n+1, n+1))$.

Solution of LSP by Householder QR factorization

Let $A \in \mathbb{R}^{m \times n}$, $m > n$, and $b \in \mathbb{R}^m$. Suppose that $\text{rank}(A) = n$. Then the LSP $Ax = b$ can be solved in three steps:

- Compute QR factorization $[A, b] = QR$. Cost: $2mn^2 - \frac{2n^3}{3}$ flops.
- Solve the upper triangular system $R(1:n, 1:n)x = R(1:n, n+1)$.
Cost: n^2 flops.
- Compute residual norm $d = \text{abs}(R(n+1, n+1))$.

Solution of LSP by Householder QR factorization

Let $A \in \mathbb{R}^{m \times n}$, $m > n$, and $b \in \mathbb{R}^m$. Suppose that $\text{rank}(A) = n$. Then the LSP $Ax = b$ can be solved in three steps:

- Compute QR factorization $[A, b] = QR$. Cost: $2mn^2 - \frac{2n^3}{3}$ flops.
- Solve the upper triangular system $R(1:n, 1:n)x = R(1:n, n+1)$.
Cost: n^2 flops.
- Compute residual norm $d = \text{abs}(R(n+1, n+1))$.

The total cost for solving the LSP $Ax = b$ by Householder QR factorization is $2mn^2 - \frac{2n^3}{3}$ flops.