

Congratulations! You passed!

Grade received 80% Latest Submission Grade 80% To pass 80% or higher

Go to next item

1. Suppose m=4 students have taken some class, and the class had a midterm exam and a final exam. You have collected a dataset of their scores on the two exams, which is as follows:

0 / 1 point

midterm exam	$(midterm\ exam)^2$	final exam
89	7921	96
72	5184	74
94	8836	87
69	4761	78

You'd like to use polynomial regression to predict a student's final exam score from their midterm exam score. Concretely, suppose you want to fit a model of the form $h_{\theta}(x) = \theta_0 + \theta_1 x_1 + \theta_2 x_2$, where x_1 is the midterm score and x_2 is (midterm score)². Further, you plan to use both feature scaling (dividing by the "max-min", or range, of a feature) and mean normalization.

What is the normalized feature $x_1^{(3)}$? (Hint: midterm = 94, final = 87 is training example 3.) Please round off your answer to two decimal places and enter in the text box below.



2. You run gradient descent for 15 iterations

1/1 point

with lpha=0.3 and compute

J(heta) after each iteration. You find that the

value of J(heta) decreases quickly then levels

off. Based on this, which of the following conclusions seems

most plausible?

⊘ Correct

Suppose you have m=23 training examples with n=5 features (excluding the additional all-ones feature for the intercept term, which you should add). The normal equation is $\theta=(X^TX)^{-1}X^Ty$. For the given values of m and n, what are the dimensions of θ , X, and y in this equation?

1/1 point

✓ Correct

Suppose you have a dataset with m=1000000 examples and n=200000 features for each example. You want to use multivariate linear regression to fit the parameters θ to our data. Should you prefer gradient descent or the normal equation?

1/1 point

⊘ Correct

5. Which of the following are reasons for using feature scaling?

1/1 point

✓ Correct