


Computing with Signals



DA 623

Jan - May 2024

IIT Guwahati

Instructors: Neeraj Sharma

Lecture-08

Legendre polynomials

$$L_k(t) = \frac{1}{2^k k!} \frac{d^k}{dt^k} (t^2 - 1)^k, \quad k \in \mathbb{N}, \quad \text{are orthogonal on } [-1, 1]$$

$$L_0(t) = 1,$$


$$L_1(t) = t,$$


$$L_2(t) = \frac{1}{2}(3t^2 - 1),$$


$$L_3(t) = \frac{1}{2}(5t^3 - 3t),$$

$$L_4(t) = \frac{1}{8}(35t^4 - 30t^2 + 3),$$


$$L_5(t) = \frac{1}{8}(63t^5 - 70t^3 + 15t).$$











Do periodic signal exist in real-life?




If you want to find the secrets of the
universe, think in terms of energy,
frequency and vibration.


— Nikola Tesla —

AZ QUOTES

Do periodic signal exist in real-life?



EEG signal (correlate of electrical activity in the brain)




Stock market fluctuations

DJIA History 2017-2020




Air pressure associated with spoken speech utterance



... and a lot more!


- Polynomial representation

$$f(x) = a_0 + a_1x + a_2x^2 + a_3x^3 + \dots$$
$$= \sum_{m=0}^{\infty} a_m x^m$$




- Fourier series representation

$$f(x) = \sum_{m=0}^{\infty} A_m \sin \left(\frac{\pi m x}{L} + \phi_n \right)$$




- Polynomial representation

$$\begin{aligned} f(x) &= a_0 + a_1x + a_2x^2 + a_3x^3 + \dots \\ &= \sum_{m=0}^{\infty} a_mx^m \end{aligned}$$



- Fourier series representation

$$f(x) = \sum_{m=0}^{\infty} A_m \sin \left(\frac{\pi mx}{L} + \phi_n \right)$$



Focus of this lecture

- Fourier series representation

$$f(x) = \sum_{m=0}^{\infty} A_m \sin\left(\frac{\pi m x}{L} + \phi_n\right)$$

$\sin(A + B) = \sin(A)\cos(B) + \cos(A)\sin(B)$

$$= \frac{a_0}{2} + \sum_{m=1}^{\infty} a_m \cos\left(\frac{\pi m x}{L}\right) + b_m \sin\left(\frac{\pi m x}{L}\right)$$

A function can be written as sum of scaled cosine() and sine() functions



Jean-Baptiste Joseph Fourier
French Mathematician & Physicist
(1768 - 1830)

THE
9937
ANALYTICAL THEORY OF HEAT

BY

Jean Baptiste JOSEPH FOURIER.



TRANSLATED, WITH NOTES,

BY

ALEXANDER FREEMAN, M.A.,

FELLOW OF ST JOHN'S COLLEGE, CAMBRIDGE.

EDITED FOR THE SYNDICS OF THE UNIVERSITY PRESS.

Cambridge:
AT THE UNIVERSITY PRESS.

LONDON: CAMBRIDGE WAREHOUSE, 17, PATERNOSTER ROW.

CAMBRIDGE: DEIGHTON, BELL, AND CO.

LEIPZIG: F. A. BROCKHAUS.

1878

[All Rights reserved.]



Jean-Baptiste Joseph Fourier
French Mathematician & Physicist
(1768 - 1830)



On the Eiffel Tower, 72 names of French scientists, engineers, and mathematicians are engraved in recognition of their contributions.

Fourier series representation

$$f(x) = \frac{a_0}{2} + \sum_{m=1}^{\infty} \left(a_m \cos\left(\frac{m\pi x}{L}\right) + b_m \sin\left(\frac{m\pi x}{L}\right) \right)$$

- It is a linear summation of $\cos(\cdot)$ and $\sin(\cdot)$, with no cross-terms
- It is sum of many (infinite) terms
- Each of the $\cos(\cdot)$ and $\sin(\cdot)$ term is periodic - $2L/m$
- Parameters of the sum are $\{a_0, a_m, b_m\}$

Fourier series representation


$$f(x) = \frac{a_0}{2} + \sum_{m=1}^{\infty} \left(a_m \cos\left(\frac{m\pi x}{L}\right) + b_m \sin\left(\frac{m\pi x}{L}\right) \right)$$

Let's visualize the **cosine** term

$$f(x) = \frac{a_0}{2} + \sum_{m=1}^{\infty} \left(a_m \cos\left(\frac{m\pi x}{L}\right) + b_m \sin\left(\frac{m\pi x}{L}\right) \right)$$

Example: Let $L = \pi$.


$m=1$




This function has a period of $2L$

$$f(x) = \frac{a_0}{2} + \sum_{m=1}^{\infty} \left(a_m \cos\left(\frac{m\pi x}{L}\right) + b_m \sin\left(\frac{m\pi x}{L}\right) \right)$$

Example: Let $L = \pi$.




This function has a period of $2L$




$$f(x) = \frac{a_0}{2} + \sum_{m=1}^{\infty} \left(a_m \cos\left(\frac{m\pi x}{L}\right) + b_m \sin\left(\frac{m\pi x}{L}\right) \right)$$

Example: Let $L = \pi$.





$$f(x) = \frac{a_0}{2} + \sum_{m=1}^{\infty} \left(a_m \cos\left(\frac{m\pi x}{L}\right) + b_m \sin\left(\frac{m\pi x}{L}\right) \right)$$

Example: Let $L = \pi$.




This function has a period of $2L$



increasing m implies increasing cycles within $2L$

$$\cos\left(\frac{m\pi x}{L}\right)$$




Spans only even functions of period $2L$

$$\cos\left(\frac{m\pi x}{L}\right)$$



$$\sin\left(\frac{m\pi x}{L}\right)$$

Spans only odd functions
of period $2L$



Together they span
~ all functions of
period $2L$

Spans only even functions of period $2L$

Fourier series representation

$$f(x) = \frac{a_0}{2} + \sum_{m=1}^{\infty} \left(a_m \cos\left(\frac{m\pi x}{L}\right) + b_m \sin\left(\frac{m\pi x}{L}\right) \right)$$



- Consider a $2L$ periodic signal $f(x)$
- How do we compute $\{a_0, a_m, b_m\}$ to represent $f(x)$?

Fourier series representation


$$f(x) = \frac{a_0}{2} + \sum_{m=1}^{\infty} \left(a_m \cos\left(\frac{m\pi x}{L}\right) + b_m \sin\left(\frac{m\pi x}{L}\right) \right)$$

- Consider a $2L$ periodic signal $f(x)$
- How do we compute $\{a_0, a_m, b_m\}$ to represent $f(x)$ in the above form?

Let's first review some properties of sine and cosine functions. This will help us.



$$\int_0^{2L} f(x) = 0$$




Integration of $\sin(\cdot)$ function over a period (or its multiples) is 0.

$$f(x) = \frac{a_0}{2} + \sum_{m=1}^{\infty} \left(a_m \cos\left(\frac{m\pi x}{L}\right) + b_m \sin\left(\frac{m\pi x}{L}\right) \right)$$

- Integration of $\sin(\cdot)$ function over a period (or its multiples) is 0.
- The same holds for cosine(\cdot) as well.
- Integrating both sides of the above equation, we get

$$a_0 = \frac{1}{L} \int_{-L}^L f(x) dx$$

Let's see a cross-term, that is multiplication of two sin(.)



$$\int_0^{2L} \sin\left(\frac{\pi X}{L}\right) \sin\left(\frac{2\pi X}{L}\right) = 0$$

Integration of cross-terms over a period (or its multiples) is 0.

Integration of any two cross-terms over a period (or its multiples) is 0.


$$\int_0^{2L} \sin\left(\frac{m_1 \pi X}{L}\right) \sin\left(\frac{m_2 \pi X}{L}\right) = 0, \quad m_1 \neq m_2$$
$$\int_0^{2L} \cos\left(\frac{m_1 \pi X}{L}\right) \cos\left(\frac{m_2 \pi X}{L}\right) = 0,$$
$$\int_0^{2L} \sin\left(\frac{m_1 \pi X}{L}\right) \cos\left(\frac{m_2 \pi X}{L}\right) = 0$$


The $\sin(\cdot)$ and $\cosine(\cdot)$ functions as used in Fourier series are orthogonal functions.

Familiar with orthogonal vectors in Euclidean spaces


Visualize the vectors as composed of values sampled from functions.

Then, orthogonality implies,

$$f^T g = \sum_{k=1}^n f[k]g[k] = 0$$




The $\sin(\cdot)$ and cosine(\cdot) functions as used in Fourier series are orthogonal functions.



$$f^T g = \sum_{k=1}^n f[k]g[k] = 0$$

Related by Riemann integral approximation.

$$\langle f, g \rangle = \int_0^{2L} f(x)g(x)dx = 0$$

$$f(x) = \frac{a_0}{2} + \sum_{m=1}^{\infty} \left(a_m \cos\left(\frac{m\pi x}{L}\right) + b_m \sin\left(\frac{m\pi x}{L}\right) \right)$$

Multiplying both sides by corresponding $\sin()$ or $\cosine()$ term and integrating, we get

$$\begin{aligned} a_m &= \frac{1}{L} \int_{-L}^L f(x) \cos\left(\frac{m\pi x}{L}\right) dx, \text{ and} \\ b_m &= \frac{1}{L} \int_{-L}^L f(x) \sin\left(\frac{m\pi x}{L}\right) dx. \end{aligned}$$

Summary,


$$f(x) = \frac{a_0}{2} + \sum_{m=1}^{\infty} \left(a_m \cos\left(\frac{m\pi x}{L}\right) + b_m \sin\left(\frac{m\pi x}{L}\right) \right)$$

$$a_0 = \frac{1}{L} \int_{-L}^L f(x) dx$$


$$a_m = \frac{1}{L} \int_{-L}^L f(x) \cos\left(\frac{m\pi x}{L}\right) dx, \text{ and}$$

$$b_m = \frac{1}{L} \int_{-L}^L f(x) \sin\left(\frac{m\pi x}{L}\right) dx.$$


Consider the signal,



Consider the signal,




Consider the periodic signal,





Can we express this signal using a Fourier series

Consider the periodic signal,





• ————— $2L$ ————— •

$$f(x) = \frac{a_0}{2} + \sum_{m=1}^M \left(a_m \cos\left(\frac{m\pi x}{L}\right) + b_m \sin\left(\frac{m\pi x}{L}\right) \right)$$




Fourier series approximation,

$$f(x) = \frac{a_0}{2} + \sum_{m=1}^M \left(a_m \cos\left(\frac{m\pi x}{L}\right) + b_m \sin\left(\frac{m\pi x}{L}\right) \right)$$




Another example,





Fourier series approximation,

$$f(x) = \frac{a_0}{2} + \sum_{m=1}^M \left(a_m \cos\left(\frac{m\pi x}{L}\right) + b_m \sin\left(\frac{m\pi x}{L}\right) \right)$$



Fourier series approximation,

$$f(x) = \frac{a_0}{2} + \sum_{m=1}^M \left(a_m \cos\left(\frac{m\pi x}{L}\right) + b_m \sin\left(\frac{m\pi x}{L}\right) \right)$$



Summary, Fourier series approximation,

$$f(x) = \frac{a_0}{2} + \sum_{m=1}^M \left(a_m \cos\left(\frac{m\pi x}{L}\right) + b_m \sin\left(\frac{m\pi x}{L}\right) \right)$$

- Can model (or represent) a periodic signal
- Parameters of the model are $\{a_0, a_m, b_m\}$ and M
- Suitable if signal has oscillatory patterns (or fluctuations)

