

A few special numbers¹

* The recurrence relation for *Fibonacci numbers* is,

$$f_i = \begin{cases} 0 & \text{if } i = 0, \\ 1 & \text{if } i = 1, \\ f_{i-1} + f_{i-2} & \text{if } i \geq 2. \end{cases}$$

The ordinary generating function of this sequence is $G(x) = f_0x^0 + f_1x^1 + \dots$

Since $f_i + f_{i+1} = f_{i+2}$ for $i \geq 2$,

$$G(x) - xG(x) - x^2G(x) = x, \text{ i.e., } G(x) = \frac{x}{1-x-x^2} = \frac{x}{(1-\alpha x)(1-\beta x)} \text{ for } \alpha = \frac{1+\sqrt{5}}{2} \text{ and } \beta = \frac{1-\sqrt{5}}{2}.$$

The $\frac{1+\sqrt{5}}{2}$ is known as the *golden ratio*, denoted with ϕ . Naturally, $\frac{1-\sqrt{5}}{2}$ is known as the *conjugate of the golden ratio*, denoted with $\hat{\phi}$.

$$\text{Specifically, } \frac{x}{(1-\phi x)(1-\hat{\phi} x)} = \frac{1}{\sqrt{5}} \left(\frac{1}{1-\phi x} - \frac{1}{1-\hat{\phi} x} \right) = \frac{1}{\sqrt{5}} ((1 + (\phi x) + (\phi x)^2 + \dots) - (1 + (\hat{\phi} x) + (\hat{\phi} x)^2 + \dots)).$$

$$\text{Therefore, } f_n = [x^n]G(x) = \frac{1}{\sqrt{5}}(\phi^n - \hat{\phi}^n) = \frac{1}{\sqrt{5}} \left(\left(\frac{1+\sqrt{5}}{2} \right)^n - \left(\frac{1-\sqrt{5}}{2} \right)^n \right).$$

* The recurrence relation for *Catalan numbers* is,

$$C_n = \begin{cases} 0 & \text{if } n = 0, \\ 1 & \text{if } n = 1, \\ \sum_{i=1}^{n-1} C_i C_{n-i} & \text{if } n \geq 2. \end{cases}$$

The ordinary generating function of this sequence is $G(x) = C_0x^0 + C_1x^1 + C_2x^2 + \dots$

$$\begin{aligned} \text{Then, } G(x) - x &= C_2x^2 + C_3x^3 + \dots \\ &= \sum_{n \geq 2} C_n x^n \\ &= \sum_{n \geq 2} \sum_{i=1}^{n-1} C_i C_{n-i} x^n \\ &= \sum_{n \geq 2} \sum_{i=1}^{n-1} C_i x^i C_{n-i} x^{n-i} \\ &= \left(\sum_{i \geq 1} C_i x^i \right) \left(\sum_{j \geq 1} C_j x^j \right) \\ &= (G(x))^2. \end{aligned}$$

$$\text{Therefore, } G(x)^2 - G(x) + x = 0, \text{ i.e., } G(x) = \frac{1 \pm \sqrt{1-4x}}{2}.$$

¹Prepared by R. Inkulu, with the help of note taken by Sawinder Kaur (TA) in class.

If $G(x) = \frac{1+\sqrt{1-4x}}{2}$, $G(0) = 1$; however, $C_0 = 0$. Hence, $G(x) = \frac{1}{2} - \frac{1}{2}(1-4x)^{1/2}$.

Using extended binomial theorem, $G(x) = \frac{1}{2} - \frac{1}{2}(\sum_{k \geq 0} \binom{1/2}{k} (-4x)^k)$.

Therefore, $[x^n]G(x) = \frac{-1}{2} \binom{1/2}{n} (-4)^n$.

$$\begin{aligned}
 &\text{But, } \binom{1/2}{n} \\
 &= \frac{(1/2)(1/2-1)(1/2-2)\dots(1/2-(n-1))}{n!} \\
 &= \frac{(1/2)(-1/2)(-3/2)\dots(-(2n-3)/2)}{n!} \\
 &= \frac{1}{2^n} \frac{(-1)^n (1)(3)\dots(2n-3)}{n!} \\
 &= \frac{1}{2^n} (-1)^{n-1} \frac{1}{n!} \frac{(2n-2)!}{(2)(4)\dots(2n-2)} \\
 &= \frac{1}{2^n} (-1)^{n-1} \frac{1}{n!} \frac{(2n-2)!}{2^{n-1}(n-1)!} \\
 &= \frac{2}{4^n} (-1)^{n-1} \frac{(2n-2)!}{n!(n-1)!} \\
 &= \frac{2}{4^n n} (-1)^{n-1} \binom{2n-2}{n-1}.
 \end{aligned}$$

$$\text{Hence, } [x^n]G(x) = \frac{-1}{2} \frac{2}{4^n n} (-1)^{n-1} \binom{2n-2}{n-1} (-1)^n 4^n = \frac{(-1)^{2n}}{n} \binom{2n-2}{n-1} = \frac{1}{n} \binom{2n-2}{n-1}.$$

If the recurrence is defined as,

$$C_n = \begin{cases} 1 & \text{if } n = 0, \\ 2 & \text{if } n = 1, \\ \sum_{i=1}^{n-1} C_i C_{n-i} & \text{if } n \geq 2. \end{cases}$$

substituting $n+1$ for n in the above, $C_n = [x^n]G(x) = \frac{1}{n+1} \binom{2n}{n}$.

* If $f(k)$ is a positive monotonically increasing continuous function (refer to Fig. 1), then

- $\int_{m-1}^n f(x)dx \leq \sum_{k=m}^n f(k) \leq \int_m^{n+1} f(x)dx$
- $f(m) + \int_m^n f(x)dx \leq \sum_{k=m}^n f(k) \leq f(n) + \int_m^n f(x)dx$




Figure 1: Approximating a summation with an integral.

Noting that $\lg(k)$ is a monotonically increasing function,
 $n \ln(n) - n + 1 \leq \sum_{i=1}^n \ln(i) \leq n \ln(n) - n + 1 + \ln(n)$. ——— (1)

Hence, $\frac{n^n}{e^{n-1}} \leq n! \leq \frac{n^{n+1}}{e^{n-1}}$.

* From (1), we know $\lg(n!) - n \ln(n) + n \leq 1 + \ln(n)$.

We claim there exists a positive constant α such that $(\ln(n!) - n \ln n + n - \frac{1}{2} \ln n) \approx \alpha$.

Then, $e^\alpha \approx e^{(\ln(n!) - (n + \frac{1}{2}) \ln n + n)} = \frac{n! e^n}{n^{n+1/2}}$. ——— (2a)

That is, $n! \approx e^\alpha n^{n+1/2} e^{-n}$. ——— (2b)

From Wallis' inequality, when n is asymptotically large, $\frac{(2)(4)(6)\dots(2n)}{(1)(3)(5)\dots(2n-1)\sqrt{2n}} \approx \sqrt{\frac{\pi}{2}}$

$$\Rightarrow \frac{(2^n n!)^2}{(2n)!} \frac{1}{\sqrt{2n}} \approx \sqrt{\frac{\pi}{2}}.$$

Substituting (2b), $\frac{2^{2n} e^{2\alpha} n^{2n+1} e^{-2n}}{e^\alpha (2n)^{2n+1/2} e^{-2n}} \frac{1}{\sqrt{2n}} \approx \sqrt{\frac{\pi}{2}} \Rightarrow e^\alpha \approx \sqrt{2\pi}$.

Substituting (2a), $e^\alpha = \frac{n! e^n}{n^{n+1/2}} \approx \sqrt{2\pi} \Rightarrow n! \approx \sqrt{2\pi n} \left(\frac{n}{e}\right)^n$.

This is known as the *Stirling's approximation of n!*.

* The recurrence relation for *Harmonic numbers* is,

$$H_n = \begin{cases} 1 & \text{if } n = 1, \\ a_{n-1} + \frac{1}{n} & \text{if } n \geq 2. \end{cases}$$

If $f(k)$ is a positive monotonically decreasing continuous function, then

$$\begin{aligned} & \bullet \int_m^{n+1} f(x) dx \leq \sum_{k=m}^n f(k) \leq \int_{m-1}^n f(x) dx \\ & \bullet f(n) + \int_m^n f(x) dx \leq \sum_{k=m}^n f(k) \leq f(m) + \int_m^n f(x) dx \text{ ——— (3)} \end{aligned}$$

Noting that $1/k$ is a monotonically decreasing function, from (3), $\frac{1}{n} + \ln(n) \leq \sum_{k=1}^n \frac{1}{k} \leq 1 + \ln(n)$.

* The *sum of first n Harmonic numbers* is, $\sum_{k=1}^n H_k$

$$\begin{aligned} &= \sum_{k=1}^n \sum_{j=1}^k \frac{1}{j} \\ &= 1 + \left(1 + \frac{1}{2}\right) + \left(1 + \frac{1}{2} + \frac{1}{3}\right) + \dots + \left(1 + \frac{1}{2} + \dots + \frac{1}{n}\right) \\ &= n + \frac{1}{2}(n-1) + \frac{1}{3}(n-2) + \dots + \frac{1}{n}(n-(n-1)) \end{aligned}$$

$$\begin{aligned}
&= \left(\sum_{j=1}^n \frac{n-j+1}{j} \right) = \left(\sum_{j=1}^n \sum_{k=j}^n \frac{1}{j} \right) \\
&= n \left(1 + \frac{1}{2} + \dots + \frac{1}{n} \right) - \left(\frac{1}{2} + \frac{2}{3} + \frac{3}{4} + \dots + \frac{n-1}{n} \right) \\
&= n \left(1 + \frac{1}{2} + \dots + \frac{1}{n} \right) - \left(\frac{1}{2} + \frac{2}{3} + \frac{3}{4} + \dots + \frac{n-1}{n} \right) - H_n + H_n \\
&= (n+1)H_n - n.
\end{aligned}$$

* The r^{th} *Stirling number of second kind* of n is the number of ways to distribute n labeled balls into r unlabeled bins with no bin left empty, denoted by $\left\{ \begin{smallmatrix} n \\ r \end{smallmatrix} \right\}$. The recurrence relation is, $\left\{ \begin{smallmatrix} n+1 \\ r \end{smallmatrix} \right\} = r \left\{ \begin{smallmatrix} n \\ r \end{smallmatrix} \right\} + \left\{ \begin{smallmatrix} n \\ r-1 \end{smallmatrix} \right\}$ for $r > 0$, with initial conditions $\left\{ \begin{smallmatrix} 0 \\ 0 \end{smallmatrix} \right\} = 1, \left\{ \begin{smallmatrix} k \\ 0 \end{smallmatrix} \right\} = \left\{ \begin{smallmatrix} 0 \\ k \end{smallmatrix} \right\} = 0$ for $k > 0$. Further, $\left\{ \begin{smallmatrix} n \\ r \end{smallmatrix} \right\} = \frac{1}{r!} \cdot (\text{number of onto functions from a set with } n \text{ elements to a set with } r \text{ elements}) = \frac{1}{r!} \sum_{i=0}^{r-1} (-1)^i C(r, i) (r-i)^n$.