

## Week 9: Modeling Volatility (ARCH/GARCH)

# Kurtosis

An ARCH(1) specification

$$\varepsilon_t = \sigma_t u_t$$

$$u_t \stackrel{iid}{\sim} N(0, 1)$$

$$\sigma_t^2 = \alpha_0 + \alpha_1 \varepsilon_{t-1}^2.$$

Recall (**Week 8**) unconditional moments of the ARCH(1) model

(**5.**) It can be shown that the kurtosis for  $\varepsilon_t$  is

$$\kappa = 3 \times \frac{(1 - \alpha_1^2)}{(1 - 3\alpha_1^2)} > 3.$$

when  $0 < \alpha_1 < \sqrt{1/3}$  (proof not supplied).

## Moments of the ARCH(1) model

- ▶ (5.) shows that the unconditional distribution of the  $\{\varepsilon_t\}$  from an ARCH model **cannot be a normal distribution... too much kurtosis.**
- ▶ Why is this important?
  1. Returns are not normal.
  2. Tails are too thick; i.e., too much kurtosis.
  3. However, we can use a conditional normal random variable to model returns.
  4. So long as we augment this random variable with ARCH-like characteristics.

# Moments of the ARCH(1) model

This means that this simple model allows us to capture two important stylized facts of returns data:

- (1) Returns have thick tails, i.e., are not (unconditionally) normal;
- (2) Risk (or volatility) is time-varying and autoregressive, i.e., returns display ARCH-like behavior;

## Estimation of an ARCH model

- ▶ MLE chooses  $c$ ,  $\phi_1$ ,  $\alpha_0$  and  $\alpha_1$  to maximise the (log of the) joint probability of the observed data points in our sample (called the log likelihood (LL)).
- ▶ For an ARCH(1) model (with AR(1) mean and normal  $u_t$  the  $LL$  is given by


$$-\frac{T}{2} \ln(2\pi) - \frac{1}{2} \sum_{t=1}^T \ln \sigma_t^2 - \frac{1}{2} \sum_{t=1}^T \frac{(r_t - c - \phi_1 r_{t-1})^2}{\sigma_t^2},$$

with

$$\sigma_t^2 = \alpha_0 + \alpha_1 (r_{t-1} - c - \phi_1 r_{t-2})^2.$$


- ▶ The form of this  $LL$  comes from the AR(1)-ARCH(1) specification and the assumption that  $u_t \stackrel{iid}{\sim} N(0, 1)$ .

ibm.xts is daily 3 Jan 2000 to 3 Jan 2018



```
# log return  
lret <- na.omit(diff(log(ibm.xts)))  
colnames(lret) <- "log return"
```

$\text{lret} = \log \text{ return}$





```

# AR(1,0)-ARCH(1)

library(rugarch)

spec_arch1 <- ugarchspec(
  variance.model = list(
    model = "sGARCH",           # standard GARCH
    garchorder = c(1, 0)        # c(ARCH,GARCH)
  ),
  mean.model = list(
    armaorder = c(1, 0),        # AR(1)
    include.mean = TRUE
  ),
  distribution.model = "norm"   # normal errors
)

# Estimate the AR-GARCH models
fit_arch1 <- ugarchfit(
  spec = spec_arch1,
  data = lret
)

```

```
> fit_arch1@fit$matcoef
```

|        | Estimate     | Std. Error   | t value   | Pr(> t )   |
|--------|--------------|--------------|-----------|------------|
| mu     | 0.0006500238 | 3.252055e-04 | 1.998810  | 0.04562896 |
| ar1    | 0.0199421101 | 1.669515e-02 | 1.194485  | 0.23228834 |
| omega  | 0.0004107126 | 1.179327e-05 | 34.826010 | 0.00000000 |
| alpha1 | 0.3042729939 | 2.621725e-02 | 11.605833 | 0.00000000 |

| Environment                                                                                                                                                                                                                                                            | History                                                                                                | Connections                                                                       | Tutorial |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------|----------|
|    Import Dataset ▾ |  342 MiB ▾            |  |          |
| R ▾                                                                                                                                                                                                                                                                    |  Global Environment ▾ |                                                                                   |          |
| fit_arch1                                                                                                                                                                                                                                                              | Formal class                                                                                           | uGARCHfit                                                                         |          |
| ..@ fit :List of 27                                                                                                                                                                                                                                                    |                                                                                                        |                                                                                   |          |
| .. ..\$ hessian                                                                                                                                                                                                                                                        | : num [1:4, 1:4]                                                                                       | 9472429 647 2623065 -4300 647 ...                                                 |          |
| .. ..\$ cvar                                                                                                                                                                                                                                                           | : num [1:4, 1:4]                                                                                       | 1.06e-07 -4.37e-09 -1.05e-10 3.52e-0                                              |          |
| .. ..\$ var                                                                                                                                                                                                                                                            | : num [1:4529]                                                                                         | 0.000594 0.000782 0.000443 0.0005 0.00                                            |          |
| .. ..\$ sigma                                                                                                                                                                                                                                                          | : num [1:4529]                                                                                         | 0.0244 0.028 0.021 0.0224 0.0414 ...                                              |          |
| .. ..\$ condH                                                                                                                                                                                                                                                          | : num                                                                                                  | 6.81                                                                              |          |
| .. ..\$ z                                                                                                                                                                                                                                                              | : num [1:4529]                                                                                         | 1.434 0.366 -0.814 2.925 -0.26 ...                                                |          |
| .. ..\$ LLH                                                                                                                                                                                                                                                            | : num                                                                                                  | 10639                                                                             |          |
| .. ..\$ log.likelihoods                                                                                                                                                                                                                                                | : num [1:4529]                                                                                         | -1.77 -2.59 -2.61 1.4 -2.23 ...                                                   |          |
| .. ..\$ residuals                                                                                                                                                                                                                                                      | : num [1:4529]                                                                                         | 0.0349 0.0102 -0.0171 0.0654 -0.0108 .                                            |          |
| .. ..\$ coef                                                                                                                                                                                                                                                           | : Named num [1:4]                                                                                      | 0.00065 0.019942 0.000411 0.304273                                                |          |
| .. ..- attr(*, "names")=                                                                                                                                                                                                                                               | chr [1:4]                                                                                              | "mu" "ar1" "omega" "alpha1"                                                       |          |
| .. ..\$ robust.cvar                                                                                                                                                                                                                                                    | : num [1:4, 1:4]                                                                                       | 1.28e-07 4.20e-06 -2.32e-09 8.99e-06                                              |          |
| .. ..\$ A                                                                                                                                                                                                                                                              | : num [1:4, 1:4]                                                                                       | 2091.506 0.143 579.171 -0.949 0.143                                               |          |
| .. ..\$ B                                                                                                                                                                                                                                                              | : num [1:4, 1:4]                                                                                       | 2385.5 23.4 -8312.5 16.8 23.4 ...                                                 |          |
| .. ..\$ scores                                                                                                                                                                                                                                                         | : num [1:4529, 1:4]                                                                                    | -59.45 -24.6 35.6 -205.36 -4.46 .                                                 |          |
| .. ..- attr(*, "dimnames")=                                                                                                                                                                                                                                            | List of 2                                                                                              |                                                                                   |          |
| .. .. . . \$                                                                                                                                                                                                                                                           | : NULL                                                                                                 |                                                                                   |          |
| .. .. . . \$                                                                                                                                                                                                                                                           | : chr [1:4]                                                                                            | "mu" "ar1" "omega" "alpha1"                                                       |          |

continued

```
.. ..$ A          : num [1:4, 1:4] 2091.506 0.143 579.171 -0.949 0.14
.. ..$ B          : num [1:4, 1:4] 2385.5 23.4 -8312.5 16.8 23.4 ...
.. ..$ scores      : num [1:4529, 1:4] -59.45 -24.6 35.6 -205.36 -4.46
.. .. ..- attr(*, "dimnames")=List of 2
.. .. ..$ : NULL
.. .. ..$ : chr [1:4] "mu" "ar1" "omega" "alpha1"
.. ..$ se.coef      : num [1:4] 3.25e-04 1.67e-02 1.18e-05 2.62e-02
.. ..$ tval         : Named num [1:4] 2 1.19 34.83 11.61
.. .. ..- attr(*, "names")= chr [1:4] "mu" "ar1" "omega" "alpha1"
.. ..$ matcoef      : num [1:4, 1:4] 0.00065 0.019942 0.000411 0.304273
.. .. ..- attr(*, "dimnames")=List of 2
.. .. ..$ : chr [1:4] "mu" "ar1" "omega" "alpha1"
.. .. ..$ : chr [1:4] " Estimate" " Std. Error" " t value" "Pr(>|t|)"
```

```
> # residuals
> res_arch1 <- residuals(fit_arch1)
> colnames(res_arch1) <- "ehat ARCH"
> tail(res_arch1)
```

|            | ehat ARCH    |
|------------|--------------|
| 2017-12-26 | 0.007137154  |
| 2017-12-27 | 0.008427157  |
| 2017-12-28 | 0.012418597  |
| 2017-12-29 | -0.009345413 |
| 2018-01-02 | 0.032257987  |
| 2018-01-03 | 0.002700624  |

```
> sigma2hat_arch1 <- sigma(fit_arch1)^2
> tail(sigma2hat_arch1)
      [,1]
2017-12-26 0.0004124374
2017-12-27 0.0004262119
2017-12-28 0.0004323211
2017-12-29 0.0004576380
2018-01-02 0.0004372868
2018-01-03 0.0007273323
```

```
> tail(ibm.xts)
              IBM
2017-12-26 45.35
2017-12-27 45.77
2017-12-28 46.38
2017-12-29 45.99
2018-01-02 47.52
2018-01-03 47.71
```

```
> tail(lret)
              log return
2017-12-26 0.007747643
2017-12-27 0.009218722
2017-12-28 0.013239499
2017-12-29 -0.008444328
2018-01-02 0.032726650
2018-01-03 0.003990324
```

## Mean-Adjusted AR models

The AR(1) model is written in mean-adjusted form as

$$y_t - \hat{\mu} = \hat{\phi}(y_{t-1} - \hat{\mu}) + \hat{e}_t,$$

where

$$\hat{\mu} = 0.0006500238$$

and

$$\hat{\phi} = 0.0199421101.$$



## Mean-Adjusted AR Models

To rewrite the model in the usual intercept form,

$$y_t = c + \phi y_{t-1} + e_t,$$

we expand the mean-adjusted AR(1) model as follows:

$$y_t = \mu + \phi y_{t-1} - \phi \mu + e_t.$$

Therefore,

$$y_t = \mu(1 - \phi) + \phi y_{t-1} + e_t.$$

Hence,

$$c = \mu(1 - \phi).$$

## Mean-Adjusted AR Models

Using the estimated values,

$$\hat{c} = 0.0006500238(1 - 0.0199421101),$$

so

$$\hat{c} \approx 0.000637061.$$

Thus, the AR(1) model in the usual intercept form is

$$y_t = 0.00064 + 0.0199421101y_{t-1} + \hat{e}_t.$$

The unconditional mean is

$$\frac{\hat{c}}{1 - \hat{\phi}} = \frac{0.000637061}{1 - 0.0199421101} = 0.0006500238.$$

## Forecasting: an AR(1)-ARCH(1) model

Fitted model for IBM returns:

$$r_t = 0.00064 + 0.01994r_{t-1} + \hat{\varepsilon}_t,$$

with

$$\hat{\varepsilon}_t = \hat{\sigma}_t \hat{u}_t, \quad \hat{\sigma}_t^2 = 0.00041 + 0.30427 \hat{\varepsilon}_{t-1}^2.$$

LL (Log likelihood) = 10638.59

Daily IBM log returns: 4 Jan 2000 to 3 Jan 2018 (4529 obs)

Let's forecast 4 Jan 2018

$\mathcal{F}_{4529}$ : We have:

$$r_{4529} = 0.004, \quad \hat{\varepsilon}_{4529} = 0.0027, \quad \hat{\sigma}_{4529}^2 = 0.00073, \quad \text{and } P_{4529} = \$47.71.$$

## Forecasting with an AR(1)-ARCH(1) model

Forecasts:

Conditional mean:

$$\begin{aligned}\hat{E}[r_{4530}|\mathcal{F}_{4529}] &= 0.00064 + 0.01994 \times (0.004) \\ &= 0.0007\end{aligned}$$

Conditional variance:

$$\begin{aligned}\hat{V}[r_{4530}|\mathcal{F}_{4529}] &= \hat{V}[\varepsilon_{4530}|\mathcal{F}_{4529}] \\ &= 0.00041 + 0.30427 \times (0.0027)^2 \\ &= 0.00041\end{aligned}$$

$\Rightarrow$  Volatility:

$$\hat{\sigma}_{4530}[r_{4530}|F_{4529}] = \sqrt{0.0004} = 0.02$$

## Forecasting using R

```
# h-step ahead forecast
h <- 1
fc_arch1 <- ugarchforecast(fit_arch1, n.ahead = h)

# Mean forecasts
mu_fc_arch1 <- as.numeric(fitted(fc_arch1))
mu_fc_arch1

# volatility forecasts
sigma_fc_arch1 <- as.numeric(sigma(fc_arch1))
sigma_fc_arch1
# Variance forecasts
var_fc_arch1 <- sigma_fc_arch1^2
var_fc_arch1
```

## Forecasting with an AR(1)-ARCH(1) model

If we assume that  $u_t \sim N(0, 1)$  so that  $\varepsilon_t | \mathcal{F}_{t-1} \sim N(0, \sigma_t^2)$  then the conditional distribution of returns will also be normally distributed.

We can compute the 95% prediction interval for the one-step-ahead forecast of  $r_t$ .

A 95% prediction interval for  $r_{4530}$  would be

$$0.0007166364 \pm 1.96 \times 0.02032072$$

giving the interval

$$[-0.0391, 0.0405]$$

## Forecasting with an AR(1)-ARCH(1) model

If we assume that  $u_t \sim N(0, 1)$  so that  $\varepsilon_t | \mathcal{F}_{t-1} \sim N(0, \sigma_t^2)$  then the conditional distribution of returns will also be normally distributed.

Recall  $P_{4529} = \$47.71$

We can form prediction intervals for the **next price**

A 95% prediction interval for  $P_{4530}$  is

$$[\$47.71e^{-0.0391}, \$47.71e^{0.0405}] = [\$45.88, \$49.68]$$

# Diagnostics for an ARCH model

- ▶ We can use the ARCH LM test to determine if serial correlation in  $\hat{u}_t^2$  remains.
  - ▶ if there appears to be serial correlation remaining in the squared residuals,
  - ▶  $\Rightarrow$  we can improve the volatility model by including more ARCH terms.



## ARCH test

```
> res_s.arch1 <- residuals(fit_arch1, standardize = TRUE)
> FintS::ArchTest(res_s.arch1, lags = 4)
```

ARCH LM-test; Null hypothesis: no ARCH effects

```
data: res_s.arch1
Chi-squared = 303.02, df = 4, p-value < 2.2e-16
```

## ARCH(q)

1. In many cases we can improve our volatility model by working with an ARCH(q) model.
2. ARCH(q) takes  $\varepsilon_t = u_t \sigma_t$ ,  $u_t \sim i.i.d.N(0, 1)$  and

$$\sigma_t^2 = \alpha_0 + \alpha_1 \varepsilon_{t-1}^2 + \dots + \alpha_q \varepsilon_{t-q}^2.$$

- ▶ In many cases, especially with **higher frequency data**, we need  $q$  to be very large;
  1. Otherwise  $\hat{u}_t^2$  will be auto-correlated.
  2. Suggests that we could improve our modeling of conditional variance.
  3. This can be solved using GARCH models (generalized ARCH).
  
- ▶ The  $\hat{u}_t$  do not appear to have come from a normally distributed  $u_t$ .
  1. Can be fixed by changing the assumptions about  $u_t$ .
  2. E.g., to get more kurtosis, we could assume that the  $u_t$  have a t-distribution.
  3. This involves maximising a different likelihood.

## Student-t Distribution

```
> fit_arch1_t@fit$matcoef
```

|        | Estimate      | Std. Error   | t value   | Pr(> t )    |
|--------|---------------|--------------|-----------|-------------|
| mu     | 0.0007977183  | 2.906414e-04 | 2.744682  | 0.006056951 |
| ar1    | -0.0201519356 | 1.689249e-02 | -1.192952 | 0.232888013 |
| omega  | 0.0004235392  | 1.834514e-05 | 23.087267 | 0.000000000 |
| alpha1 | 0.2881657780  | 3.291013e-02 | 8.756143  | 0.000000000 |
| shape  | 5.0133763383  | 3.739183e-01 | 13.407680 | 0.000000000 |

## Student-t Distribution

In 'rugarch' with *distribution.model* = "*std*", the parameter *shape* is the degrees of freedom  $\nu$  of the Student-t distribution for the model errors.

In the R output: *shape* = 5.0134  $\approx$  5

What that means:

Smaller  $\nu \rightarrow$  heavier tails  $\rightarrow$  more probability of large shocks/outliers than Normal.

As  $\nu \rightarrow \infty$  Student-t approaches the Normal.

So  $\nu \approx 5$  is heavy-tailed (typical for financial returns).

# GARCH models

ARCH models have difficulty modelling the autocorrelation in the squared residuals.

GARCH models generalise ARCH models to include lags of  $\sigma_t^2$  in the conditional variance equation as well as lags of  $\varepsilon_t^2$ , and this often alleviates this difficulty.

A GARCH(p,q) model is given by

$$\sigma_t^2 = \alpha_0 + \alpha_1 \varepsilon_{t-1}^2 + \dots + \alpha_q \varepsilon_{t-q}^2 + \beta_1 \sigma_{t-1}^2 + \dots + \beta_p \sigma_{t-p}^2.$$

# GARCH models

A GARCH(p,q) model

$$\sigma_t^2 = \alpha_0 + \alpha_1 \varepsilon_{t-1}^2 + \dots + \alpha_q \varepsilon_{t-q}^2 + \beta_1 \sigma_{t-1}^2 + \dots + \beta_p \sigma_{t-p}^2.$$

We assume that

- (i) all  $\alpha_i > 0$  and all  $\beta_j > 0$  to ensure that  $\sigma_t^2 > 0$ , and
- (ii) that  $\sum_{i=1}^q \alpha_i + \sum_{j=1}^p \beta_j < 1$  to ensure that  $\mathbb{V}(\varepsilon_t) > 0$ .

The  $\alpha_i$  and  $\beta_j$  determine how past news affect current volatility and this persistence is often measured by  $\gamma = \sum_{i=1}^q \alpha_i + \sum_{j=1}^p \beta_j$ .

If  $\alpha_0$  is the only non-zero parameter then we have constant volatility.

## Conditional moments of GARCH(p,q) errors

Recall that  $\varepsilon_t = \sigma_t u_t$  with  $u_t \sim N(0, 1)$ , so the standardised error,  $\frac{\varepsilon_t}{\sigma_t} \sim N(0, 1)$  and the conditional distribution  $\varepsilon_t | \mathcal{F}_{t-1} \sim N(0, \sigma_t^2)$ .

Thus:

1.  $E(\varepsilon_t | \mathcal{F}_{t-1}) = 0$
2.  $\mathbb{V}(\varepsilon_t | \mathcal{F}_{t-1}) = \sigma_t^2 = \alpha_0 + \sum_{i=1}^q \alpha_i \varepsilon_{t-i}^2 + \sum_{j=1}^p \beta_j \sigma_{t-j}^2$
3.  $Cov(\varepsilon_t, \varepsilon_{t-k} | \mathcal{F}_{t-1}) = 0$  for  $k \geq 1$
4. The skewness of  $\varepsilon_t | \mathcal{F}_{t-1}$  is zero; and
5. The kurtosis of  $\varepsilon_t | \mathcal{F}_{t-1}$  is 3.

All but point 3 follow directly from the fact that  $\varepsilon_t | \mathcal{F}_{t-1} \sim N(0, \sigma_t^2)$ .



## Conditional moments of GARCH(p,q) errors

The proof for point 3 relies on the fact that  $\varepsilon_{t-k}$  is known at time  $t-1$  so

$$\text{Cov}(\varepsilon_t, \varepsilon_{t-k} | \mathcal{F}_{t-1}) = E(\varepsilon_t \varepsilon_{t-k} | \mathcal{F}_{t-1}) = \varepsilon_{t-k} E(\varepsilon_t | \mathcal{F}_{t-1}) = 0.$$

## Unconditional moments of the GARCH(p, q) errors

$$(1) E(\varepsilon_t) = 0$$

$$(2) \mathbb{V}(\varepsilon_t) = \alpha_0 / (1 - \sum_{i=1}^q \alpha_i - \sum_{j=1}^p \beta_j)$$

$$(3) \text{Cov}(\varepsilon_t, \varepsilon_{t-k}) = 0 \text{ for all } k \geq 1$$

$$(4) E(\varepsilon_t^3) = 0$$

(5) It can be shown that kurtosis  $\kappa > 3$  (proof not supplied).

## Unconditional moments of the GARCH(p, q) errors

$$(1) \ E(\varepsilon_t) = 0$$

**Proof.**

$E(\varepsilon_t) = E[E(\varepsilon_t|\mathcal{F}_{t-1})]$  by the L.I.E., and since  $\varepsilon_t|\mathcal{F}_{t-1} \sim N(0, \sigma_t^2)$  it follows that  $E(\varepsilon_t|\mathcal{F}_{t-1}) = 0$ .

Therefore  $E(\varepsilon_t) = E(0) = 0$ .



$$(2) \mathbb{V}(\varepsilon_t) = \alpha_0 / (1 - \sum_{i=1}^q \alpha_i - \sum_{j=1}^p \beta_j)$$

**Proof.**

We first note that  $E(\varepsilon_t) = 0$  implies that  $\mathbb{V}(\varepsilon_t) = E(\varepsilon_t^2)$ , and then by the L.I.E.

$$E(\varepsilon_t^2) = E[E(\varepsilon_t^2 | \mathcal{F}_{t-1})] = E(\sigma_t^2).$$

Next we note that since  $\varepsilon_t | \mathcal{F}_{t-1} \sim N(0, \sigma_t^2)$  and  $E(\varepsilon_t | \mathcal{F}_{t-1}) = 0$ , it follows that  $\sigma_t^2 = \mathbb{V}(\varepsilon_t | \mathcal{F}_{t-1}) = E(\varepsilon_t^2 | \mathcal{F}_{t-1})$ . Hence

$$\begin{aligned} \mathbb{V}(\varepsilon_t) &= E(\varepsilon_t^2) \\ &= E(\sigma_t^2) \\ &= \alpha_0 + \sum_{i=1}^q \alpha_i E(\varepsilon_{t-i}^2) + \sum_{j=1}^p \beta_j E(\sigma_{t-j}^2). \end{aligned}$$

□

**Proof.**

But  $E(\varepsilon_{t-i}^2) = E(\sigma_{t-j}^2) = \mathbb{V}(\varepsilon_t)$  since  $\varepsilon_t$  is stationary, so

$$\mathbb{V}(\varepsilon_t) = \alpha_0 + \mathbb{V}(\varepsilon_t) \left[ \sum_{i=1}^q \alpha_i + \sum_{j=1}^p \beta_j \right].$$

Solving this, we get

$$\mathbb{V}(\varepsilon_t) = \alpha_0 / \left( 1 - \sum_{i=1}^q \alpha_i - \sum_{j=1}^p \beta_j \right).$$

□

## Unconditional moments of the GARCH(p,q) errors (cont)

(3)  $Cov(\varepsilon_t, \varepsilon_{t-k}) = 0$  for all  $k \geq 1$

**Proof.**

Note

$$Cov(\varepsilon_t, \varepsilon_{t-k}) = E[(\varepsilon_t - E(\varepsilon_t))(\varepsilon_{t-k} - E(\varepsilon_{t-k}))] = E[(\varepsilon_t)(\varepsilon_{t-k})],$$

since  $E(\varepsilon_t) = E(\varepsilon_{t-k}) = 0$ . Further, the L.I.E. implies that

$$E[(\varepsilon_t)(\varepsilon_{t-k})] = E[E\{\varepsilon_t \varepsilon_{t-k}\} | \mathcal{F}_{t-1}].$$

Next we note that  $\varepsilon_{t-k} | \mathcal{F}_{t-1}$  is known for  $k \geq 1$ , so

$$E[E\{\varepsilon_t \varepsilon_{t-k}\} | \mathcal{F}_{t-1}] = E[(\varepsilon_{t-k})E\{\varepsilon_t | \mathcal{F}_{t-1}\}] = 0,$$

(since  $E\{\varepsilon_t | \mathcal{F}_{t-1}\} = 0$ .)

□

## Unconditional moments of the GARCH(p,q) errors (cont)

$$(4) \ E(\varepsilon_t^3) = 0$$

**Proof.**

We firstly note that, by the L.I.E.,

$$E(\varepsilon_t^3) = E[E(\varepsilon_t^3|\mathcal{F}_{t-1})].$$

Next under the assumption that  $u_t \sim i.i.d.N(0, 1)$ , we have

$$\varepsilon_t|\mathcal{F}_{t-1} \sim N(0, \sigma_t^2) \text{ and } E(\varepsilon_t^3|\mathcal{F}_{t-1}) = 0.$$

Then,  $E(\varepsilon_t^3) = E[E(\varepsilon_t^3|\mathcal{F}_{t-1})] = E[0] = 0 \implies$  unconditional distribution is symmetric. □

## Unconditional moments of the GARCH(p,q) errors (cont)

(5) It can be shown that kurtosis  $\kappa > 3$  (proof not supplied).

- ▶ Clearly this should be true.
- ▶ If  $\beta = 0$ , we get back  $ARCH(1)$ .
- ▶  $ARCH(1)$  has  $\kappa > 3$ .
- ▶ Therefore, for  $0 \leq \beta \leq 1$ ,  $GARCH(1,1)$  has  $\kappa > 3$ .



# GARCH(p,q) Models

- ▶ What conditions are needed to ensure  $\sigma_t^2 > 0$ ?

1.  $\alpha_0 > 0$ ;
2.  $\alpha_i \geq 0$ ;
3.  $\beta_j \geq 0$ ;

- ▶ For GARCH(1,1)model:

$$\epsilon_t = \sigma_t u_t \quad \sigma_t^2 = \alpha_0 + \alpha_1 \epsilon_{t-1}^2 + \beta_1 \sigma_{t-1}^2,$$

this amounts to

1.  $\alpha_0 > 0$ ;
2.  $\alpha_1 \geq 0$ ;
3.  $\beta_1 \geq 0$ ;

# Estimation of a GARCH model

- ▶ OLS does not provide optimal estimators in this setting so we use MLE (maximum likelihood estimation), as for ARCH models.
- ▶ How to estimate GARCH models in R?

```

# ARMA(1,0)-GARCH(1,1)

spec_garch11 <- ugarchspec(
  variance.model = list(
    model = "sGARCH",           # standard GARCH
    garchorder = c(1, 1)       # c(ARCH,GARCH)
  ),
  mean.model = list(
    armaorder = c(1, 0),       # AR(1)
    include.mean = TRUE
  ),
  distribution.model = "norm"
)

# Estimate the ARMA-GARCH models
fit_garch11 <- ugarchfit(
  spec = spec_garch11,
  data = lret
)

```


```
> fit_garch11@fit$matcoef
```

|        | Estimate      | Std. Error   | t value    | Pr(> t )   |
|--------|---------------|--------------|------------|------------|
| mu     | 5.420103e-04  | 2.746391e-04 | 1.973537   | 0.04843443 |
| ar1    | -2.770662e-02 | 1.553571e-02 | -1.783414  | 0.07451887 |
| omega  | 5.565428e-06  | 2.361384e-06 | 2.356851   | 0.01843066 |
| alpha1 | 6.528361e-02  | 7.364966e-03 | 8.864074   | 0.00000000 |
| beta1  | 9.243447e-01  | 9.173868e-03 | 100.758445 | 0.00000000 |

## Estimated (Fitted) Conditional Variance

```
sigma2hat_garch11 <- sigma(fit_garch11)^2  
tail(sigma2hat_garch11)  
  
plot(sigma2hat_garch11)  
  
plot(sigma(fit_garch11)*100)
```


## Estimated (fitted) Conditional Variance



# Estimated (Fitted) Conditional Volatility

$\text{sigma}_{\text{hat\_garch11}} * 100$

2000-01-04 / 2018-01-03



## Comments on our GARCH(1,1) model

$\hat{\alpha}_1 \ll \hat{\beta}_1$ . This is typical of GARCH(1,1) models.

Persistence  $\hat{\gamma} = \hat{\alpha}_1 + \hat{\beta}_1 = 0.06528 + 0.92434 = 0.98962$

$$\hat{V}(\varepsilon_t) = \frac{\hat{\alpha}_0}{1 - \hat{\alpha}_1 - \hat{\beta}_1} = \frac{5.565 \times 10^{-6}}{1 - 0.98962} = \frac{0.000005565}{0.01038} = 0.000537.$$



## Long run variance

The implied long run variance is 0.000537 and thus the implied long run volatility is

$$\sqrt{\frac{5.565 \times 10^{-6}}{1 - 0.98962}} = \sqrt{0.000537} = 0.02317.$$

So the unconditional volatility is about 2.32%.

## Long-Run Volatility

Although financial markets may experience excessive volatility every now and again, volatility appears to eventually settle down to a long run level. The long run level is the unconditional variance.

For the GARCH(1,1) model the long-run variance is

$$\mathbb{V}(\varepsilon_t) = \frac{\alpha_0}{1 - \alpha_1 - \beta_1}.$$

In this case, the volatility is always pulled toward this long run level; that is volatility ‘mean reverts’ to its long-run level.

```
> # ehat
> res_garch11 <- residuals(fit_garch11)
> colnames(res_garch11) <- "ehat GARCH"
> tail(res_garch11)
```

|            | ehat GARCH   |
|------------|--------------|
| 2017-12-26 | 0.007153697  |
| 2017-12-27 | 0.008876356  |
| 2017-12-28 | 0.012937891  |
| 2017-12-29 | -0.008634534 |
| 2018-01-02 | 0.031935659  |
| 2018-01-03 | 0.004340041  |

```

> # variance sigma^2
> sigma2hat.garch11 <- sigma(fit_garch11)^2
> colnames(sigma2hat.garch11) <- "sigma2hat GARCH"
> tail(sigma2hat.garch11)
      sigma2hat GARCH
2017-12-26    0.0002023347
2017-12-27    0.0001959333
2017-12-28    0.0001918190
2017-12-29    0.0001938001
2018-01-02    0.0001895707
2018-01-03    0.0002473760

```

## Forecasting with an AR(1)-GARCH(1,1) model

Estimated model:

$$r_t = 0.000557 - 0.0277r_{t-1} + \hat{\varepsilon}_t, \quad \hat{\varepsilon}_t = \hat{\sigma}_t \hat{u}_t,$$

$$\hat{\sigma}_t^2 = .0000056 + 0.06528\hat{\varepsilon}_{t-1}^2 + 0.92434\hat{\sigma}_{t-1}^2.$$

$$\text{LL (Log likelihood)} = 11045.76$$

Daily IBM log returns: 4 Jan 2000 to 3 Jan 2018 (4529 obs)

$$\mathcal{F}_{4529}: r_{4529} = 0.004, \quad \hat{\varepsilon}_{4529} = 0.00434, \quad \hat{\sigma}_{4529}^2 = 0.000247$$

## Forecasting with an AR(1)-GARCH(1,1) model

- ▶ One-step ahead forecast of  $r_t$ :
- ▶  $E[r_{4530}|\mathcal{F}_{4529}] = 0.000557 - 0.0277 \times (0.004) = 0.0004$
- ▶  $\mathbb{V}[r_{4530}|\mathcal{F}_{4529}] = \mathbb{V}[\varepsilon_{4530}|\mathcal{F}_{4529}] = 0.0000056 + 0.06528 \times (0.00434)^2 + 0.92434 \times (0.000247) = 0.000235$
- ▶ so  $\sqrt{\mathbb{V}[r_{4530}|\mathcal{F}_{4529}]} = \sqrt{0.000235} = 0.0153$

## Forecasting using R

```
> h <- 1
> fc_garch11 <- ugarchforecast(fit_garch11, n.ahead = h)
>
> # Mean forecasts
> mu_fc_garch11 <- as.numeric(fitted(fc_garch11))
> mu_fc_garch11
[1] 0.0004464693
>
> # volatility forecasts
> sigma_fc_garch11 <- as.numeric(sigma(fc_garch11))
> sigma_fc_garch11
[1] 0.01534457
> # variance forecasts
> var_fc_garch11 <- sigma_fc_garch11^2
> var_fc_garch11
[1] 0.0002354558
```

Our model  $\Rightarrow E[r_{4530}|\mathcal{F}_{4529}] = 0.0004$  &  $\sigma[r_{4530}|\mathcal{F}_{4529}] = 0.0153$

If we then assume that  $u_t \sim N(0, 1)$  so that  $\varepsilon_t|\mathcal{F}_{t-1} \sim N(0, \sigma_t^2)$  then the conditional distribution of returns will also be normally distributed.

Compute the 95% prediction interval for the one-step-ahead forecast of  $r_t$  as

$$0.0004464693 \pm 1.96 \times 0.01534457 \implies [-0.0296, 0.0305]$$



## ARCH vs GARCH prediction intervals

One-step-ahead prediction interval for ARCH(1) is

$$[-0.03911125, 0.04054452]$$

One-step-ahead prediction interval for GARCH(1,1) is

$$[-0.0296, 0.0305]$$

So GARCH(1,1) has the narrower one-step ahead prediction interval than ARCH(1).

# Comparing ARCH/GARCH Models

- ▶ Likelihood Ratio test.
- ▶ ARCH( $q$ ) models are nested in GARCH( $p,q$ ) models:
  1.  $\beta_1 = \dots = \beta_p = 0 \implies ARCH(q)$ .
  2. Can jointly test for significance of  $\beta_j$ s
- ▶ If the ARCH specification in question is nested in the GARCH model, can use:

$$LR = 2 * \left( LL(\hat{\theta}_G) - LL(\hat{\theta}_A) \right)$$

1.  $LL(\hat{\theta}_G)$ : likelihood from GARCH model
2.  $LL(\hat{\theta}_A)$ : likelihood from ARCH model
3. Asymptotically  $\chi_p^2$  under the null that the models are the same.

name@fit\$LLH

```
> # log likelihood for ARCH(1)
> LogL_arch <- fit_arch1@fit$LLH
> cat("LogLikelihood : ARCH", LogL_arch, "\n")
LogLikelihood : ARCH 10638.59
>
>
> # log likelihood for GARCH(1,1)
> LogL_garch <- fit_garch11@fit$LLH
> cat("LogLikelihood : GARCH", LogL_garch, "\n")
LogLikelihood : GARCH 11045.76
```

## Comparing ARCH/GARCH Models Example

- ▶ ARCH(1) versus GARCH(1,1)
- ▶ IBM daily returns
- ▶  $LL(\hat{\theta}_A) = 10638.59$
- ▶  $LL(\hat{\theta}_G) = 11045.76$
- ▶  $LR = 2 * (11045.76 - 10638.59) = 814.34$
- ▶  $LR \rightarrow \chi_1^2$  under the null that ARCH(1)=GARCH(1,1); i.e.,  $\beta_1 = 0$
- ▶ Reject this null.

| fit_garch11                             | Formal class                    | uGARCHfit                       |
|-----------------------------------------|---------------------------------|---------------------------------|
| ..@ fit :List of 27                     |                                 |                                 |
| .. ..\$ hessian                         | : num [1:5, 1:5]                | 13260104 185 -3088128 -310      |
| .. ..\$ cvar                            | : num [1:5, 1:5]                | 7.54e-08 1.72e-09 5.74e-12      |
| .. ..\$ var                             | : num [1:4529]                  | 0.000591 0.000633 0.0006 0.0    |
| .. ..\$ sigma                           | : num [1:4529]                  | 0.0243 0.0251 0.0245 0.024 0    |
| .. ..\$ condH                           | : num                           | 8.64                            |
| .. ..\$ z                               | : num [1:4529]                  | 1.441 0.478 -0.674 2.692 -0.    |
| .. ..\$ LLH                             | : num                           | 11046                           |
| .. ..\$ log.likelihoods                 | : num [1:4529]                  | -1.759 -2.65 -2.564 0.814 -2    |
| .. ..\$ residuals                       | : num [1:4529]                  | 0.03505 0.01202 -0.0165 0.06    |
| .. ..\$ coef                            | : Named num [1:5]               | 5.42e-04 -2.77e-02 5.57e-       |
| .. .. ..- attr(*, "names")= chr [1:5]   | "mu" "ar1" "omega" "alpha1"     |                                 |
| .. ..\$ robust.cvar                     | : num [1:5, 1:5]                | 7.90e-08 -4.09e-07 -9.35e-      |
| .. ..\$ A                               | : num [1:5, 1:5]                | 2927.8217 0.0409 -681.8565      |
| .. ..\$ B                               | : num [1:5, 1:5]                | 3.08e+03 -4.51 -2.87e+05 -      |
| .. ..\$ scores                          | : num [1:4529, 1:5]             | -59.73 -22.02 25.82 -10         |
| .. .. ..- attr(*, "dimnames")=List of 2 |                                 |                                 |
| .. .. ..\$ : NULL                       |                                 |                                 |
| .. .. ..\$ : chr [1:5]                  | "mu" "ar1" "omega" "alpha1" ... |                                 |
| .. ..\$ se.coef                         | : num [1:5]                     | 2.75e-04 1.55e-02 2.36e-06 7.36 |
| .. ..\$ tval                            | : Named num [1:5]               | 1.97 -1.78 2.36 8.86 100.       |
| .. .. ..- attr(*, "names")= chr [1:5]   | "mu" "ar1" "omega" "alpha1"     |                                 |
| .. ..\$ matcoef                         | : num [1:5, 1:4]                | 5.42e-04 -2.77e-02 5.57e-0      |
| .. .. ..- attr(*, "dimnames")=List of 2 |                                 |                                 |
| .. .. ..\$ : chr [1:5]                  | "mu" "ar1" "omega" "alpha1" ... |                                 |