

Choosing the Correct Statistical Test

| Number of <u>Dependent*</u> Variables | Number of <u>Independent**</u> Variables | Type of Dependent Variable(s) | Type of Independent Variable(s) | Measure | Test(s) |
|---------------------------------------|---|-------------------------------|---------------------------------|-------------|---|
| 1 | 0 (1 population) | continuous normal | not applicable (none) | mean | one-sample t-test |
| | | continuous non-normal | | median | one-sample median |
| | | categorical | | proportions | Chi Square goodness-of-fit, binomial test |
| | 1 (2 independent populations) | normal | 2 categories | mean | 2 independent sample t-test |
| | | non-normal | | medians | Mann Whitney, Wilcoxon rank sum test |
| | | categorical | | proportions | Chi square test Fisher's Exact test |
| | 0 (1 population measured twice) <i>or</i> 1 (2 matched populations) | normal | not applicable/ categorical | means | paired t-test |
| | | non-normal | | medians | Wilcoxon signed ranks test |
| | | categorical | | proportions | McNemar, Chi-square test |
| | 1 (3 or more populations) | normal | categorical | means | one-way ANOVA |
| | | non-normal | | medians | Kruskal Wallis |
| | | categorical | | proportions | Chi square test |
| | 2 or more (e.g., 2-way ANOVA) | normal | categorical | means | Factorial ANOVA |
| | | non-normal | | medians | Friedman test |
| | | categorical | | proportions | log-linear, logistic regression |
| | 0 (1 population measured 3 or more times) | normal | not applicable | means | Repeated measures ANOVA |
| | | normal | continuous | | correlation simple linear |

| | | | | | |
|--|------------------------|-------------|-------------------------------------|---|---|
| | 1 | | | regression | |
| | | non-normal | | non-parametric | |
| | | categorical | categorical or continuous | correlation logistic regression | |
| | | | continuous | discriminant analysis | |
| | 2 or more | normal | continuous | multiple linear regression | |
| | | non-normal | | | |
| | | categorical | | logistic regression | |
| | | normal | mixed categorical and continuous | Analysis of Covariance General Linear Models (regression) | |
| | | non-normal | | | |
| | | categorical | | logistic regression | |
| | 2 | 2 or more | normal | categorical | MANOVA |
| | 2 or more | 2 or more | normal | continuous | multivariate multiple linear regression |
| | 2 sets of 2 or more | 0 | normal | not applicable | canonical correlation |
| | 2 or more | 0 | normal | not applicable | factor analysis |

* outcome

** predictor



Page editor is [Julia Hartman](#).



Last modified December 1, 2000.

The information in this site is current through August 15, 2000.

Although the authors of this Web site have made every reasonable effort to be factually accurate, no responsibility is assumed for editorial or clerical errors or error occasioned by honest mistake. All information contained on this Web site is subject to change by the appropriate officials of The University of Alabama without prior notice. Material on this Web site does not serve as a contract between The University of Alabama and any other party.