

Date of Birth
1986-04-08

Sex
Female

Physician
Dr. Troy Sosa

Institution
Cline Inc

Tumor specimen:
source Breast
CollectedDate 2023-09-07
ReceivedDate 2023-09-08
TumorPercentage 93%

Normal specimen:
source Blood
CollectedDate 2023-09-12
ReceivedDate 2023-09-12

GENOMIC VARIANTS

Somatic - Potentially Actionable		variant allele fraction
B2M	c.2T>G p.M1R Stopgain-LOF	17.61% <div></div>
Somatic - Biologically Relevant		
HBB	c.364G>C p.E7K Spliceregionvariant-LOF	27.4% <div></div>
NOTCH1	c.7398del p.E122K Spliceregionvariant-GOF	23.28% <div></div>
CCND3	c.766_776del11 p.E27K Spliceregionvariant-GOF	24.74% <div></div>
HDAC2	c.3113A>G p.R256fs*64 Frameshift-LOF	5.19% <div></div>
ERCC2	c.3113A>G p.E7V Stopgain-LOF	9.61% <div></div>

Germline - Pathogenic

No Germline - Pathogenic variants were found in the limited set of genes on which we report.

Pertinent Negatives

EZH2 **PTCH1** **SRSF2** **STAT5B**

IMMUNOTHERAPY MARKERS

Tumor Mutational Burden

29 m/Mb 46%

Microsatellite Instability Status

Stable Equivocal **High**

FDA-APPROVED THERAPIES, Current Diagnosis

KRAS G12C Inhibitors	Sotorasib	NCCN, Consensus, Non-Small Cell Lung Cancer MSK OncoKB, Level 1 KRASp.G12C G12C-GOF
----------------------	------------------	---

FDA-APPROVED THERAPIES, Other Indications

KRAS G12C Inhibitors	Sotorasib	NCCN, Consensus, Non-Small Cell Lung Cancer MSK OncoKB, Level 1 KRASp.G12C G12C-GOF
----------------------	------------------	---

ADDITIONAL INDICATORS

Unfavorable Prognosis	NCCN, Consensus, Non-Small Cell Lung Cancer KRASp.G12C Gain-of-function
-----------------------	--

CLINICAL TRIALS

A Study of VS-6766 v. VS-6766 + Defactinib in Recurrent G12V, Other KRAS and BRAF Non-Small Cell Lung Cancer	Phase 2 City, state - x mi KRAS mutation
A Phase 1/2 Study of MRTX849 in Patients With Cancer Having a KRAS G12C Mutation KRYSTAL-1	Phase 1/2 City, state - x mi KRAS mutation STK11 mutation
First-in-human Study of DRP-104 (Sirpiglenastat) as Single Agent and in Combination With Atezolizumab in Patients With Advanced Solid Tumors. (NCT04471415)	Phase 1/2 City, state - x mi NFE2L2 mutation STK11 mutation

VARIANTS OF UNKNOWN SIGNIFICANCE

Somatic	Mutation effect	Variant allele fraction
EZH2	c.1936T>A p.Y646H Spliceregionvariant-LOF NM_001011645	3.67% <div></div>
MAPK1	c.964G>A p.E322K Frameshift-LOF NM_001011645	8.93% <div></div>
SDHD	c.34G>A p.G12S Frameshift-LOF NM_001011645	10.67% <div></div>
BTK	c.3113A>G p.C481X Nonsense-GOF NM_001011645	7.56% <div></div>
EZH2	c.1937A>G p.Y646C Nonsense-GOF NM_001011645	6.35% <div></div>
NTRK1	c.1792C>T p.G607V Nonsense-GOF NM_001011645	4.79% <div></div>
Germline	Mutation effect	Condition
CTNNB1	c.134C>T p.D32N Missensevariant(exon2)-GOF NM_001011645	care

LOW COVERAGE REGIONS

U2AF1 IDH2

SOMATIC VARIANT DETAILS - POTENTIALLY ACTIONABLE

B2M

c.2T>G p.M1R Stopgain-LOF

VAF: 17.61%

STK11 (LKB1) encodes an enzyme in the serine/threonine kinase family that is responsible for maintaining energy metabolism and cellular polarization through the activation of AMP-activated protein kinase and other members of the AMPK family. The enzyme also acts as a tumor suppressor by regulating cell growth. Loss of function mutations, copy number loss, epigenetic variation, and underexpression of STK11 are associated with cancer progression.

SOMATIC VARIANT DETAILS - BIOLOGICALLY RELEVANT

HBB

c.364G>C p.E7K Spliceregionvariant-LOF

VAF: 27.4%

NFE2L2 acts as a transcription factor for proteins that contain an antioxidant response element (ARE) within their promoter sequence. Genes that contain ARE are involved in injury and inflammation response. Activating mutations and overexpression of NFE2L2 are associated with cancer progression.

NOTCH1

c.7398del p.E122K Spliceregionvariant-GOF

VAF: 23.28%

STK11 (LKB1) encodes an enzyme in the serine/threonine kinase family that is responsible for maintaining energy metabolism and cellular polarization through the activation of AMP-activated protein kinase and other members of the AMPK family. The enzyme also acts as a tumor suppressor by regulating cell growth. Loss of function mutations, copy number loss, epigenetic variation, and underexpression of STK11 are associated with cancer progression.

CCND3

c.766_776del11 p.E27K Spliceregionvariant-GOF

VAF: 24.74%

NFE2L2 acts as a transcription factor for proteins that contain an antioxidant response element (ARE) within their promoter sequence. Genes that contain ARE are involved in injury and inflammation response. Activating mutations and overexpression of NFE2L2 are associated with cancer progression.

HDAC2

c.3113A>G p.R256fs*64 Frameshift-LOF

VAF: 5.19%

PTEN encodes a phosphatase that acts as a tumor suppressor by negatively regulating the PI3K-AKT-mTOR pathway. Loss of function mutations, copy number loss, and underexpression of PTEN are associated with cancer progression.

ERCC2

c.3113A>G p.E7V Stopgain-LOF

VAF: 9.61%

PTEN encodes a phosphatase that acts as a tumor suppressor by negatively regulating the PI3K-AKT-mTOR pathway. Loss of function mutations, copy number loss, and underexpression of PTEN are associated with cancer progression.

CLINICAL HISTORY

Diagnosed on
2023-08-31