

National Immunization Schedule

Age	Vaccines given
Birth	Bacillus Calmette Guerin (BCG), Oral Polio Vaccine (OPV)-0 dose, Hepatitis B birth dose
6 Weeks	OPV-1, Pentavalent-1, Rotavirus Vaccine (RVV)-1, Fractional dose of Inactivated Polio Vaccine (fIPV)-1, Pneumococcal Conjugate Vaccine (PCV) -1*
10 weeks	OPV-2, Pentavalent-2, RVV-2
14 weeks	OPV-3, Pentavalent-3, fIPV-2, RVV-3, PCV-2*
9-12 months	Measles & Rubella (MR)-1, JE-1** , PCV-Booster*
16-24 months	MR-2, JE-2** , Diphtheria, Pertussis & Tetanus (DPT)-Booster-1, OPV – Booster
5-6 years	DPT-Booster-2
10 years	Tetanus & adult Diphtheria (Td)
16 years	Td
Pregnant Mother	Td-1, Td-2 or Td-Booster***

* **PCV** in selected states/districts: Bihar, Himachal Pradesh, Madhya Pradesh, Uttar Pradesh (selected districts) and Rajasthan; in Haryana as state initiative

** JE in endemic districts only

*** One dose if previously vaccinated within 3 years

National Immunization Schedule (NIS) for Infants, Children and Pregnant Women (Vaccine-wise)

Vaccine	When to give	Dose	Route	Site
For Pregnant Women				
Tetanus & adult Diphtheria (Td)-1	Early in pregnancy	0.5 ml	Intra-muscular	Upper Arm
Td-2	4 weeks after Td-1	0.5 ml	Intra-muscular	Upper Arm
Td- Booster	If received 2 TT/Td doses in a pregnancy within the last 3 years*	0.5 ml	Intra-muscular	Upper Arm
For Infants				
Bacillus Calmette Guerin (BCG)	At birth or as early as possible till one year of age	0.1ml (0.05ml until 1 month age)	Intra-dermal	Left Upper Arm
Hepatitis B - Birth dose	At birth or as early as possible within 24 hours	0.5 ml	Intra-muscular	Antero-lateral side of mid-thigh
Oral Polio Vaccine (OPV)-0	At birth or as early as possible within the first 15 days	2 drops	Oral	Oral
OPV 1, 2 & 3	At 6 weeks, 10 weeks & 14 weeks (OPV can be given till 5 years of age)	2 drops	Oral	Oral
Pentavalent 1, 2 & 3	At 6 weeks, 10 weeks & 14 weeks (can be given till one year of age)	0.5 ml	Intra-muscular	Antero-lateral side of mid-thigh
Pneumococcal Conjugate Vaccine(PCV)	Two primary doses at 6 and 14 weeks followed by Booster dose at 9-12 months	0.5 ml	Intra-muscular	Antero-lateral side of mid-thigh
Rotavirus (RVV)	At 6 weeks, 10 weeks & 14 weeks (can be given till one year of age)	5 drops (liquid vaccine) 2.5 ml (lyophilized vaccine)	Oral	Oral
Inactivated Polio Vaccine (IPV)	Two fractional dose at 6 and 14 weeks of age	0.1 ml	Intra dermal two fractional dose	Intra-dermal: Right upper arm
Measles Rubella (MR) 1st dose	9 completed months-12 months.	0.5 ml	Sub-cutaneous	Right upper Arm

Vaccine	When to give	Dose	Route	Site
Japanese Encephalitis (JE) - 1	9 completed months-12 months.	0.5 ml	Sub-cutaneous (Live attenuated vaccine) Intramuscular(Killed vaccine)	Left upper Arm (Live attenuated vaccine) Anterolateral aspect of mid thigh (Killed vaccine)
Vitamin A (1st dose)	At 9 completed months with measles-Rubella	1 ml (1 lakh IU)	Oral	Oral
For Children				
Diphtheria, Pertussis & Tetanus (DPT) booster-1	16-24 months	0.5 ml	Intra-muscular	Antero-lateral side of mid-thigh
MR 2nd dose	16-24 months	0.5 ml	Sub-cutaneous	Right upper Arm
OPV Booster	16-24 months	2 drops	Oral	Oral
JE-2	16-24 months	0.5 ml	Sub-cutaneous (Live attenuated vaccine) Intramuscular(Killed vaccine)	Left upper Arm (Live attenuated vaccine) Anterolateral aspect of mid thigh (Killed vaccine)
Vitamin A (2nd to 9th dose)	16-18 months. Then one dose every 6 months up to the age of 5 years.	2 ml (2 lakh IU)	Oral	Oral
DPT Booster-2	5-6 years	0.5 ml.	Intra-muscular	Upper Arm
Td	10 years & 16 years	0.5 ml	Intra-muscular	Upper Arm

*One dose if previously vaccinated within 3 years

Note:

- JE Vaccine is introduced in select endemic districts after the campaign.
- The 2nd to 9th doses of Vitamin A can be administered to children 1-5 years old during biannual rounds, in collaboration with ICDS.
- PCV in selected states/districts: Bihar, Himachal Pradesh, Madhya Pradesh, Rajasthan & Uttar Pradesh (selected districts), and in Haryana as state initiative