



Probabilistic Synapse Detection in Array Tomography

Anish K. Simhal^{1*}, Cecilia Aguerrebere¹, Forrest Collman² Joshua T. Vogelstein³, Kristina D. Micheva⁴, Richard J. Weinberg⁵, Stephen J. Smith², Guillermo Sapiro^{1, 6}

¹Electrical and Computer Engineering, Duke University ²Synapse Biology, Allen Institute for Brain Sciences ³Department of Biomedical Engineering, Johns Hopkins University ⁴Molecular and Cellular Physiology, Stanford University School of Medicine ⁵Department of Cell Biology and Physiology, University of North Carolina ⁶Department of Biomedical Engineering, Department of Computer Science, Department of Mathematics, Duke University



Challenge

- Unsupervised automatic detection of specific synaptic subtypes with confidence values using array tomography data




Figure 1: Left: PSD-95 (red) and Synapsin (green) data overlaid on EM data. Center: PSD-95 IF image. Right: Synapsin IF image. Synaptic cleft marked in blue.

Background

- What is a Synapse?




Figure 2: Molecular architecture of an excitatory PSD-95 expressing synapse depicted in a simplified cartoon form

- Array Tomography (AT) Pipeline





Figure 3: Array Tomography Methodology [6]

Action

- Overview of Proposed Method



- Step 1: Foreground Probability

$$p_B(x, y, z) = \frac{1}{\sigma_B \sqrt{2\pi}} \int_{v(x,y,z)}^{\infty} e^{-\frac{(t-\mu_B)^2}{2\sigma_B^2}} dt. \quad (1)$$

$$p_F(x, y, z) = 1 - p_B(x, y, z). \quad (2)$$




Figure 4: (Left) Logarithm of the IF raw data. (Right) Foreground probability map. Slices of the PSD-95 antibody of size $5.268 \times 5.827 \mu m$.

- Step 2: 2D Puncta Probability.

$$p_P(x, y, z) = \prod_{i=x-W}^{x+W} \prod_{j=y-W}^{y+W} p_F(i, j, z), \quad (3)$$

W is the expected size of a punctum.




Figure 5: (Left) Probability map. (Right) Probability map of each pixel belonging to a 2D punctum.

- Step 3: 3D Puncta Probability

$$f(x, y, z) = \exp \left\{ - \sum_{j=j_{start}}^{j=j_{end}} [p_P(x, y, z) - p_P(x, y, z + j)]^2 \right\} \quad (4)$$

$$p_{3DP}(x, y, z) = p_P(x, y, z) f(x, y, z), \quad (5)$$




Figure 6: **Top:** Three consecutive slices of the 2D puncta probability. **Bottom:** Factor image given by Eq. (4) (left) and the corresponding 3D puncta probability (right) of the center slice in the top row. Each image is a cutout of size 2261×2501 pixels or $5.268 \times 5.827 \mu m$.

- Step 4: Presynaptic and Postsynaptic Puncta Adjacency.




Figure 7: **First row:** cutout of PSD-95 Punctum with center pixel highlighted. **Second row:** synapsin cutouts with the search grid overlaid.

Resolution

- Result:




Figure 8: Output probability map, each pixel's intensity value represents to the probability of it belonging to a synapse.

Dataset	Excitatory				Inhibitory			
	EM	IF	EM	IF	EM	IF	EM	IF
KDM-SYN-120905	0.88	0.91	0.90	0.93	-	-	-	-
KDM-SYN-140115	0.92	0.94	0.93	0.95	0.82	0.81	0.91	0.91

Figure 9: Precision and recall values on conjugate array tomography data




Figure 10: Synptogram showing IF data for an EM identified synapse. Each 'block' is $1.221 \mu m \times 1.233 \mu m$.

Acknowledgments

This work was supported by the National Institutes of Health (NIH-TRA 1R01NS092474), the Allen Institute for Brain Sciences (AIBS), the U.S. Office of Naval Research (ONR), the U.S. Army Research Office (ARO), the National Science Foundation (NSF), and the U.S. National Geospatial Intelligence Agency (NGA).

References

- Busse B, Smith S. Automated analysis of a diverse synapse population. PLoS Comput Biol. 2013 Mar 28;9(3):e1002976.
- Collman F, Buchanan J, Phend KD, Micheva KD, Weinberg RJ, Smith SJ. Mapping synapses by conjugate light-electron array tomography. The Journal of Neuroscience. 2015 Apr 8;35(14):5792-807.
- Harris KM, Weinberg RJ. Ultrastructure of synapses in the mammalian brain. Cold Spring Harbor Perspectives in Biology. 2012 May 1;4(5):a005587.
- Micheva KD, Smith SJ. Array tomography: a new tool for imaging the molecular architecture and ultrastructure of neural circuits. Neuron. 2007 Jul 5;55(1):25-36.
- Micheva KD, Busse B, Weiler NC, O'Rourke N, Smith SJ. Single-synapse analysis of a diverse synapse population: proteomic imaging methods and markers. Neuron. 2010 Nov 18;68(4):639-53.
- Weiler NC, Collman F, Vogelstein JT, Burns R, Smith SJ. Synaptic molecular imaging in spared and deprived columns of mouse barrel cortex with array tomography. Scientific Data. 2014 Dec 23;1.