

Akshay Sharma (Corresponding Author)
Department of Computer Science and Engineering
SAGE University, Indore 452020, Madhya Pradesh, India
Email: akssha74@gmail.com

September 6, 2026

Prof. Peng Cui (Editor-in-Chief)
Prof. Dunlian Qiu (Executive Editor-in-Chief)
Journal of Mountain Science
Institute of Mountain Hazards and Environment, Chinese Academy of Sciences (CAS)
Email: jms@imde.ac.cn

Subject: Submission of Original Research Article — “Topographic Radar Geometry-Guided Cross-Attention for Alpine Avalanche Debris Mapping in Sentinel-1 SAR: Cross-System Generalization Across the Alps, Pamirs, and Arctic”

Dear Prof. Cui, Prof. Qiu, and Editorial Board,

We are pleased to submit our original research article entitled “Topographic Radar Geometry-Guided Cross-Attention for Alpine Avalanche Debris Mapping in Sentinel-1 SAR: Cross-System Generalization Across the Alps, Pamirs, and Arctic” for consideration for publication in the Journal of Mountain Science.

Background and Motivation:

Snow avalanches represent one of the most destructive natural hazards in alpine and high-mountain regions worldwide, posing acute threats to settlements, transport arteries, and human safety across the European Alps, Central Asian Pamirs, and Arctic mountain belts. Rapid, all-weather mapping of avalanche debris is vital for emergency rescue operations and hazard mitigation. While spaceborne C-band Synthetic Aperture Radar (Sentinel-1 SAR) provides cloud-penetrating surveillance day and night, SAR change detection in rugged alpine topography suffers from severe geometric distortions: local incidence angle (LIA) variation induces intense ground-range foreshortening on radar-facing foreslopes, while slopes facing away from the radar (backslopes) suffer radiometric stretching and radar shadow. Existing deep learning change detection backbones treat SAR channels as flat 2D arrays, ignoring radar-topographic physics and generating spurious false alarms on illuminated ridges while missing shadow-adjacent debris.

Key Scientific and Methodological Contributions:

1. Geometric Formulation of 2D Convolutional Inadequacy (Proposition 2): We formulate the geometric limitations of shift-invariant 2D spatial linear convolutions operating purely on SAR intensity space across sloped mountain facets under non-stationary illumination $\frac{\cos \psi}{\sin \theta_{\text{inc}}}$. We show that projecting terrain into a continuous 4D aspect-look manifold ($\mathcal{M} = [\sin \alpha, \cos \alpha, \cos \psi, \theta_{\text{align}}]^T$) eliminates circular branch-cut discontinuities and provides continuous coordinates to condition feature representations without angular branch cuts.
2. Multi-Scale Radar-Topographic Cross-Attention (RTCAB): We design a lightweight cross-attention module wherein multi-scale SAR change features interrogate adaptively-pooled

regional topographic geometry keys and values. Component-removal and Seed-42 stratum analyses quantify empirical sensitivity without claiming a causal internal mechanism.

3. Heuristic Geomorphic Slope Prior (Geo-Loss): We audit rather than idealize the prior: its mask overlaps 18.47% of training positives and lowers held-out positive recall relative to No-GeoLoss, despite a favorable aggregate Tromsø precision/F1 trade-off. We therefore present it as a documented precision–recall heuristic, not a physical-admissibility mechanism.
4. Empirical Rigour and Cross-System Zero-Shot Evaluation: Evaluated on AvalCD across four mountain systems (7 events; 993 polygons), TopoRadar-Net records $63.79\% \pm 1.88\%$ pooled macro F1 across 3 seeds. Against Attention U-Net, the 339-block pooled differences for Seeds 42, 123, and 456 are $+3.26\%$, -2.99% , and $+4.16\%$ (descriptive mean $+1.48\% \pm 3.18\%$), while the paired three-seed macro-F1 difference is $+0.38\%$ ($p = 0.869$). Only the three-seed comparison with SiamUNet-diff is significant ($+8.92\%$, $p = 0.021$).
5. Multi-Seed Topographic Stratification: Exploratory Seed-42 backslope differences ($+5.85\%$ in Tromsø; $+5.09\%$ in Pamir) do not persist as stable three-seed effects: mean differences versus Attention U-Net are $-0.34\% \pm 6.28\%$ and $+1.20\% \pm 7.73\%$, respectively. This confirmatory stability check transparently delimits directional-conditioning claims.
6. Calibrated Operational Proxy with Failure Boundaries: Validation-only Platt calibration, connected components, and a frozen OSM major-road snapshot yield 83.0% precision and 62.5% recall for Pamir corridor alerts, with all cached TopoRadar scene runs below 5 s, but all seeds miss the single Tromsø corridor avalanche. This directly measures and limits operational readiness rather than asserting dispatch benefit.

Journal Fit and Policy Compliance:

- Format Compliance: In accordance with the Editorial Office instructions for double-blind external review, the main manuscript is submitted as an editable Word (.docx) document with all author and institutional identifying information removed; the 4 figures and 8 tables are embedded within the single main document (file size well under 10 MB). Author names, affiliations, and e-mail addresses are provided separately in the online submission system.
- Originality: This manuscript represents original, previously unpublished research and is not under consideration for publication elsewhere.
- Open Science and Reproducibility: The AvalCD benchmark is openly accessible on Zenodo (DOI: [10.5281/zenodo.15863589](https://doi.org/10.5281/zenodo.15863589)). All code and replication suites are public at <https://github.com/akssha74/TopoRadar-Net-AvalCD>. Publication-bound checkpoints, the formatted anonymous Word manuscript, and all artifacts are distributed in versioned post-freeze Release v1.0.21: <https://github.com/akssha74/TopoRadar-Net-AvalCD/releases/tag/v1.0.21>.

Suggested Independent Reviewers:

1. Dr. Markus Eckerstorfer, Senior Research Scientist, NORCE Norwegian Research Centre, Tromsø, Norway. Email: maec@norceresearch.no
Expertise: Sentinel-1 SAR change detection for snow avalanches, Arctic cryospheric hazards.

2. Dr. Achille Gatti, Department of Electronics, Information and Bioengineering, Politecnico di Milano, Milan, Italy. Email: achille.gatti@polimi.it
Expertise: AvalCD benchmark lead author, deep learning change detection on SAR imagery.
3. Dr. Silvan Leinss, Senior Researcher, Institute of Environmental Engineering, ETH Zurich, Switzerland. Email: leinss@ifu.baug.ethz.ch
Expertise: Microwave radar backscatter characteristics of snow avalanches, SAR polarimetry.

Conflict of Interest Statement:

The authors declare that they have no financial or non-financial conflicts of interest with any person, institution, or company in relation to the work reported in this manuscript. The suggested reviewers listed above are independent researchers with no personal, professional, financial, or collaborative relationship with the authors; none of them has co-authored publications with the authors, shared institutional affiliation, or participated in this study. This work received no external grant funding, and the authors have no competing commercial interests.

Opposed Reviewers (Reviewer Exclusions):

The authors do not request the exclusion of any specific individual from the pool of potential reviewers. Should the Editorial Office nevertheless wish to avoid any perception of conflict, we have no objection to the standard exclusion of reviewers sharing the authors' current institutional affiliation.

Author Agreement:

All authors have read and approved the final version of the manuscript and agree to its submission to the Journal of Mountain Science. This manuscript represents original, previously unpublished work that is not under consideration for publication elsewhere, in whole or in part.

Thank you very much for your time and consideration of our manuscript.

Sincerely,

Akshay Sharma (on behalf of all authors)
Department of Computer Science and Engineering
SAGE University, Indore 452020, India
Email: akssha74@gmail.com