

WORKSHEET

STATISTICS WORKSHEET-1

Q1 to Q9 have only one correct answer. Choose the correct option to answer your question.

1. Bernoulli random variables take (only) the values 1 and 0.

a) True

b) False

2. Which of the following theorem states that the distribution of averages of iid variables, properly normalized, becomes that of a standard normal as the sample size increases?

a) Central Limit Theorem

b) Central Mean Theorem

c) Centroid Limit Theorem

d) All of the mentioned

3. Which of the following is incorrect with respect to use of Poisson distribution?

a) Modeling event/time data

b) Modeling bounded count data

c) Modeling contingency tables

d) All of the mentioned

4. Point out the correct statement.

a) The exponent of a normally distributed random variables follows what is called the log- normal distribution

b) Sums of normally distributed random variables are again normally distributed even if the variables are dependent

c) The square of a standard normal random variable follows what is called chi-squared distribution

d) All of the mentioned

5. _____ random variables are used to model rates.

a) Empirical

b) Binomial

c) Poisson

d) All of the mentioned

6. 10. Usually replacing the standard error by its estimated value does change the CLT.

a) True

b) False

7. 1. Which of the following testing is concerned with making decisions using data?

a) Probability

b) Hypothesis

c) Causal

d) None of the mentioned

8. 4. Normalized data are centered at _____ and have units equal to standard deviations of the original data.

a) 0

b) 5

c) 1

d) 10

9. Which of the following statement is incorrect with respect to outliers?

a) Outliers can have varying degrees of influence

b) Outliers can be the result of spurious or real processes

c) Outliers cannot conform to the regression relationship

d) None of the mentioned

WORKSHEET

Q10 and Q15 are subjective answer type questions, Answer them in your own words briefly.

10. What do you understand by the term Normal Distribution?

***normal distribution is an arrangement of a data set in which most values cluster in the middle of the range and the rest taper off symmetrically toward either extreme.**

11. How do you handle missing data? What imputation techniques do you recommend?

12. What is A/B testing?

**** A/B Testing, also known as split testing, refers to a randomized experimentation process wherein two or more versions of a variable (web page, page element, etc.) are shown to different segments of website visitors at the same time to determine which version leaves the maximum impact and drive business metrics.***

13. Is mean imputation of missing data acceptable practice?

****The process of replacing null values in a data collection with the data's mean is known as mean imputation.***

Mean imputation is typically considered terrible practice since it ignores feature correlation. Consider the following scenario: we have a table with age and fitness scores, and an eight-year-old has a missing fitness score. If we average the fitness scores of people between the ages of 15 and 80, the eighty-year-old will appear to have a significantly greater fitness level than he actually does. Second, mean imputation decreases the variance of our data while increasing bias. As a result of the reduced variance, the model is less accurate and the confidence interval is narrower.

14. What is linear regression in statistics?

**** Linear regression is a kind of statistical analysis that attempts to show a relationship between two variables. Linear regression looks at various data points and plots a trend line. Linear regression can create a predictive model on apparently random data, showing trends in data, such as in cancer diagnoses or in stock prices.***

15. What are the various branches of statistics?

**** There are three real branches of statistics: data collection, descriptive statistics and inferential statistics.***