Emerging patterns of tyrosine sulfation and O-glycosylation cross-talk and co-localization

Search within citing articles

Sulfoconjugation of protein peptides and glycoproteins in physiology and diseases

P Xu, X Cai, X Guan, W Xie - Pharmacology & therapeutics, 2023 - Elsevier

Protein sulfoconjugation, or sulfation, represents a critical post-translational modification (PTM) process that involves the attachment of sulfate groups to various positions of …

☆ Save 99 Cite Cited by 15 Related articles All 6 versions

GalNT2-mediated O-glycosylation affects pancreas development and function in mice

B Mercanoglu, SA Waschkowski, E Neuburg… - Scientific Reports, 2024 - nature.com

GALNT2, also known as polypeptide N-acetylgalactosaminyltransferase 2, is an enzyme that catalyzes the initial step of O-linked glycosylation, a crucial posttranslational modification …

☆ Save 99 Cite Cited by 7 Related articles All 11 versions

Development and biological applications of sulfur–triazole exchange (SuTEx) chemistry

AL Borne, JW Brulet, K Yuan, KL Hsu - RSC chemical biology, 2021 - pubs.rsc.org

Sulfur electrophiles constitute an important class of covalent small molecules that have found widespread applications in synthetic chemistry and chemical biology. Various …

☆ Save 99 Cite Cited by 41 Related articles All 10 versions

Uncommon posttranslational modifications in proteomics: ADP-ribosylation, tyrosine nitration, and tyrosine sulfation

A Bashyal, JS Brodbelt - Mass spectrometry reviews, 2024 - Wiley Online Library

Posttranslational modifications (PTMs) are covalent modifications of proteins that modulate the structure and functions of proteins and regulate biological processes. The development …

☆ Save 99 Cite Cited by 23 Related articles All 10 versions

The C-terminal peptide of CCL21 drastically augments CCL21 activity through the dendritic cell lymph node homing receptor CCR7 by interaction with the receptor N …

AS Jørgensen, EP Brandum, JM Mikkelsen… - Cellular and Molecular …, 2021 - Springer

The endogenous chemokines CCL19 and CCL21 signal via their common receptor CCR7. CCL21 is the main lymph node homing chemokine, but a weak chemo-attractant compared …

☆ Save 99 Cite Cited by 31 Related articles All 15 versions

“Glyco-sulfo barcodes” regulate chemokine receptor function

L Verhallen, JJ Lackman, R Wendt… - Cellular and Molecular …, 2023 - Springer

Chemokine ligands and receptors regulate the directional migration of leukocytes. Post-translational modifications of chemokine receptors including O-glycosylation and tyrosine …

☆ Save 99 Cite Cited by 21 Related articles All 11 versions

Fluorosulfate as a latent sulfate in peptides and proteins

C Liu, X Liu, M Zhou, C Xia, Y Lyu, Q Peng… - Journal of the …, 2023 - ACS Publications

Sulfation widely exists in the eukaryotic proteome. However, understanding the biological functions of sulfation in peptides and proteins has been hampered by the lack of methods to …

☆ Save 99 Cite Cited by 16 Related articles All 10 versions

Cosmc controls B cell homing

J Zeng, M Eljalby, RP Aryal, S Lehoux… - Nature …, 2020 - nature.com

The molecular mechanisms regulating lymphocyte homing into lymph nodes are only partly understood. Here, we report that B cell-specific deletion of the X-linked gene, Cosmc, and …

☆ Save 99 Cite Cited by 37 Related articles All 10 versions

Overexpression of Tn antigen induces chronic pancreatitis in mice

B Mercanoglu, N Schraps, AD Giannou, E Neuburg… - Scientific Reports, 2025 - nature.com

Altered O-glycosylation is a key contributor to various pathophysiological processes. Notably, the expression of the Tn antigen is primarily attributed to dysfunction of the …

☆ Save 99 Cite Cited by 7 Related articles All 10 versions

Electron capture vs transfer dissociation for site determination of tryptic peptide tyrosine sulfation: direct detection of Fibrinogen Sulfation sites and Identification of …

MM Youssef, CW Szot, J Folz, LM Collier… - Journal of proteome …, 2024 - ACS Publications

Tyrosine sulfation, an understudied but crucial post-translational modification, cannot be directly detected in conventional nanoflow liquid chromatography-tandem mass …

☆ Save 99 Cite Cited by 8 Related articles All 7 versions