bioRxiv preprint doi: https://doi.org/10.1101/2023.04.28.538759; this version posted April 29, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Stability and functional evolution of a novel family of hyperstable apple snail egg proteins

Pasquevich, M. Y.[a,b]; Dreon, M. S.[a,b], Diupotex-Chong M. E.[c] & Heras, H.[a,d]

[a] Instituto de Investigaciones Bioquímicas de La Plata ¨Prof. Dr. Rodolfo R. Brenner¨ (INIBIOLP), Universidad Nacional de La Plata (UNLP) —CONICET CCT-La Plata, La Plata, Argentina.

[b] Cátedra de Bioquímica y Biología Molecular, Facultad de Ciencias Médicas, UNLP, Argentina.

[c] Instituto de Ciencias del Mar y Limnología, Universidad Nacional Autónoma de México, Ciudad de México, México.

[d] Cátedra de Química Biológica, Facultad de Ciencias Naturales y Museo, UNLP, Argentina.

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.28.538759; this version posted April 29, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

ABSTRACT

The relationship between protein stability and function evolution in nature has not been explored in detail. We investigate this fundamental question by characterizing the Perivitellin-1 (PV1) orthologs within *Pomacea* snails. These highly stable egg carotenoproteins, crucial to the snail's reproductive success, have acquired clade-related protective functions. To gain insight into the molecular evolution of reproductive proteins along a genus we studied *P. patula* PV1 (PpaPV1) from Flagellata eggs, the most basal clade of *Pomacea*. PpaPV1 stands as the most stable PV1, with a much longer unfolding half-life and resistance to detergent unfolding, indicating a loss of kinetic stability in the derived clades. In fact, PpaPV1 is among the most hyperstable proteins described in nature. Its molecular weight, size, and high resistance to proteolysis are similar to and shared with all PV1s. Its spectral characteristics and associated pale egg coloration as well as a mild lectin activity indicate these protein traits appeared early in the evolution of the genus. However, lectin binding is less strong, and glycan specificity is narrower than those PV1s of its sister Bridgesii clade.

Our results provide experimental evidence of the relationship between protein stability and evolvability, showing that along their rapid evolution, PV1s tradeoff between structural and kinetic stability loss and the acquisition of diverse defensive traits: a strong lectin activity (Bridgesii clade), or an aposematic warning coloration (Canaliculata clade). This study offers insight into the molecular evolution of a novel family of invertebrate reproductive proteins that accompanied the fast diversification of the genus and may have contributed to Canaliculata snails invasiveness.

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.28.538759; this version posted April 29, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

INTRODUCTION

Protein stability affects functional aspects and evolutionarily, and this has established a tight balance between increased functionality through accumulation of mutations and the ability to maintain an adequate level of stability (Gershenson et al., 2014). There are, however, not many examples from nature regarding these tradeoff, as opposed to directed evolution in laboratory (Zheng et al., 2020) or theoretical models (Agozzino & Dill, 2018; Tokuriki & Tawfik, 2009a; Zeldovich et al., 2007).

Apple snails (Pomacea spp.) are an emerging model for evolutionary studies due to their high diversity, ancient history, and wide geographical distribution (Hayes et al., 2009; Sun et al., 2019). *Pomacea* spp. are amphibious snails that have evolved an unusual reproductive strategy, laying eggs outside the water (Hayes et al., 2009). This transition to terrestrial egg laying went along with the acquisition of remarkable molecular and biochemical changes, particularly their reproductive egg proteins (Ip et al., 2019). Multiple lines of evidence on adaptive evolution in the egg proteins contribute to our understanding of how aquatic gastropod ancestors invaded terrestrial habitats (Ip et al., 2019).

The traditional view that proteins possess absolute functional specificity and a single, fixed structure conflicts with their marked ability to adapt and evolve new functions and structures (Tokuriki & Tawfik, 2009b). In some animal genera, orthologous proteins play similar roles but undergo major functional adaptations. This is the case of *Pomacea* snail eggs' multifunctional carotenoproteins which evolved different functions in different lineages while retaining others unchanged (Brola et al., 2020). The pathway of paralogous gene evolution leading to diversification of the original function is unknown for *Pomacea* proteins. Particularly, one family of *Pomacea* reproductive egg carotenoprotein, called Perivitellin-1 (PV1) have orthologous in *Pomacea* and the basal genera (Ip et al., 2019). Remarkably, no sequence similarity with proteins of other organisms outside the ampullariid family was found suggesting that PV1s have aroused by duplication of orphan genes. PV1s orthologous have been studied only in the most derived clades. They share several structural features: are colored, high molecular weight oligomers, very glycosylated, and composed by combinations of subunits with similar amino acid sequences (Brola et al. 2020). Besides, the association of PV1s with carotenoids seems to be exclusive of *Pomacea* genus, at least those from the most derived Bridgesii and Canaliculata clades (Dreon et al., 2004a; Ituarte et al., 2008; Pasquevich et al., 2014, Brola et al., 2020). Members of this novel family of invertebrate egg reproductive proteins also

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.28.538759; this version posted April 29, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

display high thermal and pH structural stability, as well as a kinetic stability (Pasquevich et al., 2017, Brola et al., 2020).

PV1s are massively accumulated in eggs (Giglio et al., 2016) playing a role as a storage protein and a major source of nutrients during embryo development (Heras et al., 1998). Moreover, PV1s carry and protect antioxidant carotenoids from the harsh environmental conditions of development (Dreon et al., 2004a; Ituarte et al., 2008; Pasquevich et al., 2014). Remarkably, they are a poor amino acids source to predators because of its low digestibility which renders it an antinutritive protein. Noteworthy, besides embryo nutrition and the antinutritive (non-digestive) role, PV1s have clade-related functions according to their phylogenetic position: those PV1s from the Canaliculata clade, PcOvo and PmPV1, provide a bright reddish coloration associated to a warning signal, an ecological function that goes along with the presence of a toxic perivitelin, PV2 only found in the Canaliculata clade (Giglio et al., 2020; Heras et al., 2008). On the contrary, the members of the Bridgesii clade lay more pale eggs of pinkish color (presumably non-warning signal) eggs have PV1s like PsSC from *P. scalaris* (Ituarte et al., 2008, 2010, 2012) and PdPV1 from *P. diffusa* (Brola et al., 2020) that possess a strong lectin activity, *i. e.* capacity to recognize and bind glycans, and adversely affect gut morphophysiology of predators, a role absent in PV1s from the Canaliculata clade. These disparate functions among orthologous goes along with their remarkably high stability and provide a unique and unexplored model to understand the evolution of hyperstable proteins, an aspect poorly studied experimentally in biological models.

We began by studying the structure, stability, and functional features of a PV1s from a species of Flagellata the most basal clade of the genus, which allowed a phylogenetical analysis of the evolutionary structural and functional trends of these fascinating hyperstable proteins. We found that the molecular evolution of these novel family of reproductive proteins accompanied the diversification of the genus and may have facilitated some apple snails to become notorious invasive pests.

MATERIAL AND METHODS

Sample collection and PV1s purification

Pomacea patula eggs were collected in the Catemaco Lake, Veracruz, Mexico and kept in the laboratory at -70 ºC until processed. The perivitellin fluid (PVF) was obtained as previously described (Pasquevich et al., 2014). In short, eggs were homogenized on ice in Tris/HCl 20 mM buffer 1:3 w/v and sequentially centrifuged at 4 ºC for 20 min. at

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.28.538759; this version posted April 29, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

10.000 xg and 50 min at 100.000 xg. The obtained supernatant contains the soluble egg fraction.

To compare and trace the evolution of PV1 carotenoproteins from *Pomacea*, PV1 from *P. patula catemaquensis* (hereafter PpaPV1) was purified from egg clutches as previously described for others apple snail eggs carotenoproteins (see below). Total protein was quantified following the method described by Bradford using bovine serum albumin (BSA, Sigma cat. 7906) as standard. Purity was checked by polyacrylamide gel electrophoresis (PAGE). Another *Pomacea* spp. carotenoproteins (*i. e.* PcOvo, PmPV1 and PsSC), used in some assays, were purified as previously described (Dreon et al., 2004b, Ituarte et al., 2008, Pasquevich et al., 2014).

Structure

Oligomer and subunits electrophoretic behavior

Gel electrophoresis was used to characterize and compare the oligomer and subunits of PpaPV1 with other *Pomacea* PV1s.

Native PAGE with Laemmli buffer (pH 8.8) without SDS was performed in 4–20% gradient polyacrylamide gels in a miniVE Electrophoresis System (GE Healthcare, Life Science). High molecular weight standards (Pharmacia) were run in the same gels. Subunits were separated by SDS-PAGE in 4–20% gradient polyacrylamide gels containing 0.1% SDS; samples were denatured at 95 °C, with dithiotreitol and β-mercaptoethanol treatment (Laemmli, 1970). Low molecular weight standards (Pharmacia) were used, and gels were stained with Coomassie Brilliant Blue R-250 (Sigma Chemicals). In both gels PcOvo PmPV1 and PsSC were run for comparison of PV1s from other clades.

Immunoblotting

Cross-reactivity of PpaPV1 with PV1s from other clades were assayed with anti-PcOvo and anti-PsSC sera. PV1s (7 µg) and molecular weight marker (Dual Color, Bio-Rad) were transferred from SDS-PAGE 16% gels onto nitrocellulose membranes (Amersham) in a Mini Transblot Cell (Bio RadLaboratories, Inc.), using 25 mM Tris–HCl, 192 mM glycine, 20% (v/v) methanol, pH 8.3 buffer. After blocking for 2 h at room temperature with 3% (w/v) nonfat dry milk in PBS-Tween, the membranes were incubated overnight at 4 °C with polyclonal antibodies against PcOvo (Dreon et al., 2003) in 1:10.000 dilution, or polyclonal antibodies against PsSC (Ituarte et al., 2008) in a 1:12.000 dilution, in 3%

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.28.538759; this version posted April 29, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

(w/v) nonfat dry milk in PBS-Tween. Specific antigens were detected after incubating 2 h at room temperature goat anti-rabbit IgG horseradish peroxidase conjugate (BioRad Laboratories, Inc.). Immunoreactivity was visualized by electro-chemiluminiscence in a Chemi-Doc MP Imaging System (Bio Rad).

Size Exclusion Chromatography

Size exclusion chromatography (SEC) allows to estimate the molecular weight of proteins comparing the chromatographic retention times with those of standard proteins of precise weight (Barth & Boyes, 1990). Molecular weight of PV1s by SEC were analyzed with a SEC-Superdex 200 10/300 GL column (Amersham) in an isocratic size exclusion HPLC (1260 Infinity, Agilent technologies) with UV detection in isocratic mode. Mobil phase contained 137 mM. NaCl, 2.7 mM KCl, 2.7 mM KCl mM, 10 mM, Na_2HPO_4, 1.8 mM KH_2PO_4 (PBS). The flow rate was 0.5 mL/min and the detector was set at 280 nm. The elution volume (Ve) of PV1s and 500 µL of standard proteins in PBS (5 mg/mL Thyroglobulin (MW 669000), 2.8 mg/mL Ferritin (MW 440000), 4 mg/mL Aldolase (MW 159000) and Ovoalbumin (MW 45000)) were determined by measuring the volume of the eluent from point of injection to the center of the elution peak. Blue Dextran 2000, 1 mg/mL was used to calculate the column void volume (V_0). The calibration curve was performed fitting a curve in a plot of $K_{av} = (V_e-V_0)/(V_c-V_0)$, where V_c is the geometric volume of the column (24 mL), versus the log molecular weight of each standard. Standard curve was fitted to a line and PV1s MW calculate extrapolating from the standard curve.

Dynamic Light Scattering

The dynamic light scattering (DLS) of a nanoparticles sample in solution revealed the particle size distribution in real-time (Falke & Betzel, 2019). DLS is particularly sensitive to large aggregates, common in some PV1s (Ituarte et al., 2008, Brola et al., 2020). Thus, DLS analysis were performed in line right after SEC-eluted PV1s. PV1s sizes were monitored in PBS at 25°C, using a Malvern Zetasizer nano-zs instrument. The scatter light signals were collected at 173-degree scattering angle (backscatter) and three measurements of an automatic number of runs each were conducted per sample. Protein parameters were analyzed with the Zetasizer Software v 7.13. Data used to size measurement meets quality criteria. Intensity size distributions was used to size

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.28.538759; this version posted April 29, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

calculation. Volume size distribution was used to check the main peak contribution to the analysis.

N-terminal sequence

Subunits of purified PpaPV1 separated by electrophoresis were transferred to PVDF membranes and sequenced by Edman degradation at the Laboratorio Nacional de Investigación y Servicios en Péptidos y Proteínas (LANAIS-PRO, Universidad de Buenos Aires—CONICET). The system used was an Applied Biosystems 477a Protein/Peptide Sequencer interfaced with an HPLC 120 for one-line phenylthiohydantoin amino acid analysis. N-terminal sequences were compared with other *Pomacea* sequences using the multiple sequence alignment program CLUSTAL 2.1 (Larkin et al., 2007).

Spectrophotometric analysis

Absorption spectra of egg carotenoproteins are valuable taxonomic characters in *Pomacea* spp. (Pasquevich & Heras, 2020). PV1s absorb light in the visible region of the spectrum because of the presence of carotenoids pigments (Heras et al., 2007). Absorption spectra of PVF and purified carotenoproteins were recorded between 350 nm to 650 nm in an Agilent 8453 UV/Vis diode array spectrophotometer (Agilent Technologies, Waldbronn, Germany).

STRUCTURAL AND KINETIC STABILITY

Effect of pH and temperature on structural stability. To study the effect of pH on PpaPV1 structural stability, the protein was incubated overnight in different buffers ranging from pH 2.0 to 12.0 following a previously used method (Pasquevich et al., 2017). Samples were analysed by absorbance and fluorescence spectroscopy. Absorbance spectra were recorded between 300–650 nm in an Agilent 8453 UV/Vis diode array spectrophotometer (Agilent Technologies, Waldbronn, Germany) taking advantage of the fact that PV1s absorb in the visible range allowing to follow the protein-carotenoid interaction by its spectrum in this range. Fluorescence emission was recorded as described in the *Chemical denaturation* section (see below). Two independent samples were measured, and the corresponding buffer blank was subtracted. The effect of temperature on PpaPV1 at pH=7.4 was also measured by absorption and fluorescence spectroscopy in the range 25–85 °C. The effect of extreme thermal conditions was analyzed by boiling

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.28.538759; this version posted April 29, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

PpaPV1 for 50 min evaluating the oligomer integrity using native (non-denaturing) gel electrophoresis, as previously done (Pasquevich et al. 2017).

Chemical denaturation. The intrinsic fluorescence emission of PpaPV1 and PsSC tryphtophans was used to follow the PV1s denaturation induced by guanidine hydrochloride (GdnHCl) (Sigma). Chemical denaturation was performed by incubating overnight 50 µg/mL of PV1s in the presence of increasing concentrations (0–6.5 M) of GdnHCl buffered with 0.1 M phosphate buffer at pH 7.4 at 8°C.

Protein intrinsic fluorescence spectra were recorded on an Fluorolog 3 Spectrofluorometer coupled with a Lauda Alpha RA 8 thermostatic bath. Fluorescence spectra were recorded in emission scanning mode at 25 °C. Tryptophan emission was excited at 295 nm (6 nm slit) and recorded between 310 and 450 nm (3 nm slit). The corresponding buffer blank was subtracted. Two independent samples were measured. Spectra were characterized by their center of mass (CM) and the populations associated to the unfolded fraction (fu) were calculated from the CM as for PmPV1 in Pasquevich *et al*. (2017). The equilibrium reached in each GdnHCl concentration allowed the calculation of an equilibrium constant K=fu/(1-fu) and the Gibb´s free energy for the unfolding reaction in terms of this mole fractions ($\Delta G^0 = -RT \ln K$) were calculated. The dependence of ΔG^0 on GdnHCl concentration can be approximated by the linear equation $\Delta G^0 = \Delta G^0 H_2O - m[\text{GdnHCl}]$, where the free energy of unfolding in the absence of denaturant ($\Delta G^0 H_2O$) represents the conformational stability of the protein. The GdnHCl concentration in which half of the protein is unfolded (C_m) was estimated as a function of denaturant concentration from the linear extrapolation method.

Resistance to sodium dodecyl sulphate

Resistance to sodium dodecyl sulphate (SDS)-induced denaturation serves to identify proteins whose native conformations are kinetically trapped in a specific conformation because of an unusually high-unfolding barrier that results in very slow unfolding rates. The resistance to SDS was assayed following the Manning and Colon procedure previously used with other *Pomacea* spp. carotenoproteins (Pasquevich et al 2017, Brola et al. 2020). Briefly, PcOvo, PmPV1, PsSC and PpaPV1 were incubated in Laemmli sample buffer (pH=6.8) containing 1% SDS and either boiled for 10 min or unheated prior to its analysis by 4–20% SDS-PAGE. The gels were then stained with Coomassie blue.

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.28.538759; this version posted April 29, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Unfolding Kinetics induced by GdnHCl. Unfolding of proteins in increased concentrations of GdnHCl allow to calculate the rate of unfolding (half-life) in the absence of denaturant under native conditions (Manning & Colón, 2004a). Fluorolog-3 (Horiba-Jobin Yvon) fluorometer was used to measure PpaPV1, PsSC and PmPV1 kinetics of unfolding. Protein solutions in 100 mM phosphate buffer, pH 7.4 (PB) were treated with GdnHCl solution in the same buffer. For the fluorescence kinetic experiment, increasing concentrations of GdnHCl in PB in a 10 mm pathlength cuvette were incubated. PV1s (final concentration of 50 µg/mL) were mixed by pipetting up and down with 1 mL pipette. Data collections were started after the chamber was closed. The shutters open automatically. Time zero was manually determined as 10 sec after the protein were added to the denaturant. The excitation/emission were 299/360 nm. The relaxation time were fit to an exponential equation. The unfolding constants were obtained for each GdnHCl concentration. The rate constants as a function of GdnHCl were extrapolated to native conditions to obtain an estimate of the rate constant (k) in absence of denaturant. Half-life were calculated as ln 2/k.

Resistance to Proteolysis: proteinase K assay.

Protein structural rigidity makes proteins resistant to proteolysis. PV1s rigidity were assay by Proteinase K treatment, performed following Kim *et al.* (2004) using the concentrations modified by Frassa *et al.* (2010) and previously performed to PmPV1 (Pasquevich et al., 2017). PpaPV1 (1 mg/mL) was incubated with proteinase K (1, 10 and 100 µg/mL) in 50 mM Tris/HCl buffer (pH 8.0) containing 10 mM CaCl2 at 37 °C for 30 min. Digestion was ended by boiling samples in SDS sample buffer, and products were analysed by SDS-PAGE as above.

Functions of PpaPV1

Hemagglutinating activity

PV1s of the Bridgesii clade has a strong lectin activity and ability to agglutinate rabbit erythrocytes (Ituarte et al., 2012, Brola 2020), but PV1s of canaliculata clade lack this capacity (Pasquevich et al., 2017). We tested this capacity in *P. patula* PpaPV1 using the same methodology. In short, PpaPV1 hemagglutinating activity was tested by hemagglutination of red blood cells (RBC) from rabbit obtained in facilities of University of La Plata. Erythrocytes were prepared as stated in Ituarte et al. (2012) with minimal

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.28.538759; this version posted April 29, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

modifications. Two-fold serial dilutions of PpaPV1 in PBS (25 µl) were incubated with an equal volume of 2% (v/v) erythrocytes in PBS in U-shaped microtiter plates (Greiner Bio-One) at 37°C for 2 h. The initial PVF protein concentration was 6.7 mg/mL and PpaPV1 concentration was 1.3 mg/mL. Results are expressed as the lowest protein concentration showing visible hemagglutinating activity by the naked eye.

Specificity for glycan-binding

Glycan binding specificity of PpaPV1 was determined at the Core H of the Consortium for Functional Glycomics (http://www.functionalglycomics.org, Emory University, Atlanta, GA, USA). To detect primary binding of PpaPV1 to glycans, the protein was fluorescently labeled using the Alexa Fluor 488 Protein Labeling kit (Invitrogen, Life Technologies-Molecular Probes) according to the manufacturer's instructions. Protein concentration and the degree of labeling were determined spectrophotometrically. Fluorescently labeled PpaPV1 was assayed on a glycan array which comprised 585 glycan targets (version 5.4). The highest and lowest points from a set of six replicates were removed and the four remaining values were averaged. PpaPV1 glycan microarray data were compared with PsSC (Ituarte et al., 2018) and PdPV1 (Brola et al., 2020) microarrays using the Glycan Array Dashboard (glycotoolkit.com/GLAD/) (Mehta & Cummings, 2019).

***In vitro* intestinal digestion and high proteolysis assays**

The simulated gastroduodenal digestion of PpaPV1 was analyzed by sequentially incubating the protein for 2 h with pepsin (gastric) and 2 h with trypsin (intestinal) at 37 °C, using the method described by Moreno *et al.* (2005), with some modifications as described in Pasquevich et al. 2017. Briefly, PpaPV1 in double-distilled water was dissolved in simulated gastric fluid (SGF) (0.15 M NaCl, pH 2.5) to a final concentration of 0.5 µg/µL. Digestion commenced by adding porcine pepsin (Sigma, cat. P6887) at an enzyme:substrate ratio of 1:20 (w/w). Gastric digestion was conducted at 37 °C with shaking for 120 min. Aliquots of 5 µg protein were taken at 0, 60 and 120 min for SDS-PAGE. The reaction was stopped by increasing the pH with 150 mM Tris/Cl buffer pH 8.5. Samples were immediately boiled for 5 min in SDS electrophoresis buffer with β-mercaptoethanol (4%) and analysed as described above.

For *in vitro* duodenal digestion, the gastric digest was used as starting material. The digest was adjusted to 8.5 and sodium taurocholate (Sigma) added. The simulated

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.28.538759; this version posted April 29, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

duodenal digestion was conducted at 37 °C with shaking using bovine pancreas trypsin (Sigma cat. T9935) at an enzyme:substrate ratio of 1:2.8 (w/w). Aliquots were taken at 0, 60 and 120 min for SDS-PAGE analysis. BSA was used as positive (with enzyme) and negative (without enzyme) control in both gastric and duodenal digestion.

RESULTS

STRUCTURAL FEATURES

Native PpaPV1 is 265.9 kDa which is rather similar to that of all other PV1s (**Table 1**). DLS analysis showed a single main peak of distribution of intensity and volume in the 12.2-14.0 diameter size range for PV1s from the 3 clades (**Fig. 1A, Fig S1**). PV1s from basal clades also showed a minor aggregation peak (**Fig. S1B**).

Electrophoretic analysis revealed that under native conditions the mobility of PpaPV1 differ from others PV1s of the derived clades: PsSC from the Bridgesii clade, PcOvo and PmPV1 from the Canaliculata clade (**Fig. 1B**). Under denaturing conditions (SDS-PAGE) PpaPV1 showed several subunits between 25 and 35 KDa, the same molecular weight range as reported for the other PV1 of the genus (**Fig. 1B**).

Anti-serum against PsSC carotenoprotein (Bridgesii clade) cross reacts with PpaPV1 (Flagellata), but not with PmPV1 and PcOvo (Canaliculata clade) **(Fig. 1C)** while PcOvo anti-serum recognized Canaliculata and Bridgesii clades PV1s (*i.e.* PcOvo, PmPV1 and PsSC) as previously described (Dreon et al., 2003, Pasquevich et al., 2017, Brola et al., 2020) but not subunits of PpaPV1 (**Fig. 1C**).

PpaPV1 was separated into 6 subunits numbered according to their electrophoretic mobility. The seven N terminal sequences obtained were grouped into two nearly identical sequences (**Fig. S2-A**). Sequence similarity analysis with PcOvo, PmPV1, and PsSC. PpaPV1_3a revealed 70.6% similarity with all three PV1s (**Fig. S2-B**), while PpaPV1_6 presents 70.6% similarity with PsSC (Bridgesii clade) and 55.6% with PmPV1 and PcOvo (Canaliculata clade) reflecting a greater evolutionary distance with the latter (**Fig. S2-C**). PpaPV1_3b Nt region has no sequence similarity with any reported *Pomacea* spp. perivitellin.

P. patula PpaPV1 and PVF absorbs in a wide range of the visible spectra (350-650 nm) (**Fig. 2**) with a maximum at 380 nm an absorption maximum shared with Bridgesii clade (*P. scalaris*) but that differ from Canaliculata clade PV1s. Besides, the overall absorption

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.28.538759; this version posted April 29, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

intensity between 450-600 nm increased according to the phylogenetic position from Flagellata towards Canaliculata clades (**Fig. 2**).

STRUCTURAL AND KINETIC STABILITY

Structural stability against pH, temperature and chemical chaotropes

PpaPV1 remained stable in a wide range of pH. A slight alteration in the fine structure of the UV-visible spectrum (**Fig. S3A**) and an increase in fluorescence emission intensity (**Fig. S3B**) was only observed at pH=2.0. The absorption and emission spectra of PpaPV1 remained virtually unchanged even at temperatures of 80-85 °C (**Fig. S3 C-D**). Also, the electrophoretic behavior after boiling PpaPV1 for 60 min was unchanged, as reported for others *Pomacea* carotenoproteins (**Fig. S4**).

The chemical stability of PV1s showed an overall increase in fluorescence intensity and a systematic red shift of the spectra when increasing GdnHCl concentrations (**Fig. S5**). **Figure 3A** shows that the GndHCl unfolding transition of PpaPV1 reaches a plateau and experimental data fits a two-state model. The GdnHCl concentration required to reach the midpoint of the transition between both states (Cm) was lower in PmPV1 and PsSC than in PpaPV1 (the GdnHCl concentration to obtain fifty percent unfolded) (Figure 3B, Table 2). The disassembling/unfolding process followed by changes in the standard free energy $\Delta G°H_2O$, was higher in PpaPV1 than in PmPV1 (Figure 3B, Table 2) shows this parameter in different *Pomacea* spp. PV1s.

Resistance to sodium dodecyl sulphate induced denaturation

Proteins with a high energetic barrier between the folded and unfolded states are very resistant to unfolding and are considered kinetically stable. Comparison of the migration on polyacrylamide gels of identical boiled and unboiled PV1s previously incubated with SDS indicated PpaPV1 was resistant to SDS-induced denaturation. However, PV1s from Bridgesii and Canaliculata clades, display a partial loss and therefore some oligomers in the unheated samples disaggregates in its subunits (**Fig. 4A**).

Unfolding rate and resistance to proteolysis: kinetic stability of PV1s

Subjecting carotenoprotein to increasing GdnHCl concentration and measuring the time they took to unfold allowed the calculation of the half-life of proteins in native conditions without the presence of the chaotrope. PV1s of the Flagellata, Bridgesii and Canaliculata clades were assayed, namely PpaPV1, PsSC and PmPV1. The rate at which PpaPV1

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.28.538759; this version posted April 29, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

carotenoprotein unfolds is markedly lower than that of its derived clade counterparts. Consequently, the half-life of PpaPV1 was several orders greater than PsSC and PmPV1 (**Table 3**).

Limited proteolysis of PpaPV1 with Proteinase K, a fungal protease with broad specificity showed no evidence of PpaPV1 hydrolysis, while BSA (control) was completely digested (**Fig. S6**).

FUNCTIONAL CHARACTERISTICS

Lectin activity

The carbohydrate binding capacity of PpaPV1 evaluated by hemagglutination assays showed activity in a dose dependent manner. The hemagglutinating activity was up to 0.16 µg/µL (**Figure 5A**). The specificity and relative affinity of PpaPV1 towards oligosaccharides structures evaluated by a high-throughput glycan array assay. Albeit with low affinity, PpaPV1 showed a binding pattern to glycans related to Blood A group containing a specific motif (GalNAca1-3(Fuca1-2)Galb1-4GlcNAc) (Figure 5B). The specificity toward oligosaccharides are shown in Table 4. Among them, the specificity of PpaPV1 with Blood group A type 2 antigens, as well as the lack of specificity toward sialic acid antigens, are remarkable.

In vitro simulated gastrointestinal digestion

PpaPV1 resists hydrolysis when exposed sequentially to 2 h of gastric and duodenal phases (Fig. 7S) while BSA (control) was readily degraded. PpaPV1 maintained its electrophoretic behaviour for up to 120 min.

DISCUSSION

The complexity of the heteroligomerization increases in the most derived species.

PV1s heteroligomers are composed by combinations of related subunits, probably paralogues arised by duplication and speciation from an orphan gene (Sun et al. 2012). Analysis of PV1 from the most basal clade of the genus (Hayes et al. 2009) allowed for the first time to look into the evolution of structure, stability and functional features of proteins within a single genus. Figure 6 synthesizes our current knowledge on this. The first conclusion is that while mass and size of PV1 particles remained similar along evolution and their subunits have roughly the same molecular weight regardless of the

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.28.538759; this version posted April 29, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

phylogenetic position, marked structural changes are present. Particles seem to have acquired different post-translational modifications and surface structural changes were further evidenced by the lack of cross-reactivity of Flagelatta and Bridgesii PV1 with antibodies directed against Canalicula PV1s. As expected from the orphan gene origin of this novel family of proteins, no similarities with sequences reported outside ampullarids were observed in PpaPV1. Besides, while oligomers from derived clades combine 6 different paralogue subunits, PpaPV1 combine only 3 different subunits, one with no similarities with any other PV1s, further suggesting that subunits aroused by gene duplication (Sun et al. 2012, Pasquevich et al. 2017, Ip et al. 2019, Brola et al. 2020) and some subunits were lost along evolution. In addition, the Flagellata and Bridgesii clades carotenoproteins share common spectroscopic features that change markedly in PV1s from the most derived species.

Evolutionary significance of the stability loss of apple snail eggs carotenoproteins

The term *protein stability* indicates the ability to retain the native conformation when subjected to physical or chemical manipulation. PpaPV1 is structurally highly stable, rather difficult to unfold (denature) by near-boiling temperatures, extreme pHs and high concentrations of a denaturing chemical, a common feature shared with Bridgesii and Canaliculata carotenoproteins (Ituarte et al., 2012, Dreon et al., 2007, Dreon et al., 2007, Pasquevich et al., 2017, Brola et al. 2020). However, PpaPV1 displayed a much higher resistance to chemical denaturation that the other orthologs.

Our results strengthen the notion that long half-lives proteins (kinetically stables) withstand SDS detergent retaining a rigid folded core that only unfolds when boiled (Manning and Colon 2004). In our study PpaPV1 shows a much longer half-life and a greater resistance to denaturation by SDS than other PV1s. Moreover, PpaPV1 is among the most stable proteins so far reported while PmPV1 lies in the transition between the most stable proteins and those that are not (Table 2). PpaPV1, as other *Pomacea* spp. carotenoproteins are also exceptionally resistant to proteolysis, suggesting that a common mechanism may account for their resistance to SDS and proteolytic cleavage (Manning & Colón, 2004). More broadly, kinetically stable proteins are typically extracellular and several protecting from oxidative stress have evolved as kinetically stable (Colón et al., 2017). In this regard, PV1s have both features as they are located in the fluid surrounding the embryos and its pigments provide antioxidant protection to eggs (Dreon et al., 2004a).

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.28.538759; this version posted April 29, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Along the evolution of PV1s we noticed a partial loss of stability, but only to a certain point as the removal of carotenoids does not affect the stability of PcOvo (Dreon et al., 2007). This supports the idea that PcOvo stability and probably other members of the PV1 family favours its physiological role in the storage, transport, and protection of carotenoids during snail embryogenesis.

The hyperstability of PpaPV1 basal carotenoprotein could have allowed tolerance to mutations, *i.e.*, acquiring new functions (functional evolution) without losing the native structure, in agreement with Bloom *et al.* (2006), that proposed that the high stability allow the evolvability of proteins.

That high kinetic stability of these orthologs defense proteins is a vital property to their protective role and is related with the snail fitness (reproductive success).

Roles of PpaPV1 and evolutionary functions of PV1s

Terrestrial egg deposition in *Pomacea* was a key adaptation to avoid aquatic predation and/or parasitism (Sun et al., 2019). This evolutionary driver modulated sanil egg defenses and under this selective pressure, PV1 orthologous, while retaining some roles, underwent major functional adaptations. Thus, PV1s kept their ancestral traits as storage proteins to nurture embryo but not digestible by predators, but gradually lost its stability and gained new roles (**Figure 6**) a known tradeoff between the evolution of new-function and protein stability (Tokuriki & Tawfik, 2009b). The unfolding speed of PV1 in the basal clade is dramatically lower than that of the most derived clade homologue and we can hypothesize that this loss of stability allowed in Canaliculata proteins structural changes favoring a better stabilization of more polar carotenoids and the ability to accommodate larger quantities of this pigment. This agrees with other studies that indicate the loss of stability could have contributed to the gain of new functions (Bloom et al. 2006). This PV1 feature came at the price of losing the lectin capacity as it is discussed below.

To the best of our knowledge, PV1 evolution is one of the few examples taken from nature where the tradeoff between stability and evolvability of a protein is reported. One interesting aspect is the evolution of a lectin capacity that is lost in the most derived clade. This is at first intriguing considering its significant role in the defenses against predation (Brola et al., 2020) but may be explained by the evolutionary novelty in canaliculate clade of a dual enterotoxic/neurotoxin lectin (PV2) combining two ancient immune proteins (Giglio et al., 2020). On the other hand, the Bridgesii clade not only retained Flagelatta lectin capacity, but PpaPV1 moderate lectin activity gives way to PV1s with higher

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.28.538759; this version posted April 29, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

affinity to glycans and a broader specificity indicative of at least two high-affinity recognition sites in *Bridgesii* clade (Ituarte et al., 2018). Remarkably, PsSC and PdPV1 binding recognition pattern includes Galβ1-3GalNAc and a common sialic acid in vertebrate gangliosides (Brola et al., 2020; Ituarte et al., 2018) that are virtually absent in the glycans patterns recognized by PpaPV1. The inability of PpaPV1 to recognize gangliosides, and the nearly limited group A type II antigen recognition patterns, suggest that a single recognition site would be present in this ancient PV1. It can be speculated that the partial loss of PV1s stability in Bridgesii may have favored the evolution of their improved glycan recognition and binding strength capabilities.

CONCLUSION

This study shows the evolution of the structural and functional features of a novel family of invertebrate egg proteins originated from orphan genes. The basal clade contains PpaPV1, one of the most kinetically stable proteins known to date. More broadly, the study provides one of the few examples taken from nature, as opposed to directed evolution in laboratory, showing that during orthologue evolution there is a tradeoff between a loss of structural and kinetic stability and a simultaneous acquisition of new defensive traits. The extent to which these mechanisms are evolutionary steps or alternative trajectories in the evolution of selective expression of defensive strategies in the eggs is an open question. This work also increases the knowledge of ampullarids biology referred to the evolution of the complex defense system of *Pomacea* apple snail eggs.

Considering that *Pomacea* have split from its sister genus just about 28 mya (Sun et al., 2019), the study also highlights how a rapid evolution of structure-function features of reproductive proteins accompanied the spread and diversification of *Pomacea* across freshwater habitats. Predator-induced protein evolution may have contributed to the evolved defence strategies that accompanied the diversification of the genus and may have contributed to canaliculata snails worldwide invasiveness.

.

ACKNOWLEDGMENTS

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.28.538759; this version posted April 29, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

This work was supported by grants from Agencia Nacional de Promoción Científica y Técnica (PICT 2017-3142 to MYP and PICT 2017-1815 to HH) and Universidad Nacional de La Plata, Argentina. MYP and HH are members of CONICET, Argentina. MSD is member of CICBA, Argentina. We thank L. Bauzá for her help in PpaPV1 purification and Dr. L. Falomir-Lockard for help in DLS analysis. We acknowledge the participation of the Protein-*Glycan* Interaction Resource of the CFG and the National Center for Functional Glycomics (NCFG) at Beth Israel Deaconess Medical Center, Harvard Medical School **(***supporting grant R24 GM137763***).**

REFERENCES

Agozzino, L., & Dill, K. A. (2018). Protein evolution speed depends on its stability and abundance and on chaperone concentrations. *Proceedings of the National Academy of Sciences of the United States of America*, *115*(37). https://doi.org/10.1073/pnas.1810194115

Barth, H. G., & Boyes, B. E. (1990). Size Exclusion Chromatography. *Analytical Chemistry*, *62*(12), 268–303. https://doi.org/10.1021/ac00211a020

Bloom, J. D., Labthavikul, S. T., Otey, C. R., & Arnold, F. H. (2006). Protein stability promotes evolvability. *Proceedings of the National Academy of Sciences of the United States of America*, *103*(15), 5869–5874. https://doi.org/10.1073/pnas.0510098103

Brola, T. R., Dreon, M. S., Qiu, J. W., & Heras, H. (2020). A highly stable, non-digestible lectin from Pomacea diffusa unveils clade-related protection systems in apple snail eggs. *Journal of Experimental Biology*, *223*(19). https://doi.org/10.1242/jeb.231878

Colón, W., Church, J., Sen, J., Thibeault, J., Trasatti, H., & Xia, K. (2017). Biological Roles of Protein Kinetic Stability. *Biochemistry*, *56*(47), 6179–6186. https://doi.org/10.1021/acs.biochem.7b00942

Dreon, M., Lavarias, S., Garin, C. F., Heras, H., & Pollero, R. J. (2002). Synthesis, distribution, and levels of an egg lipoprotein from the apple snail Pomacea canaliculata (mollusca: Gastropoda). *Journal of Experimental Zoology*, *292*(3). https://doi.org/10.1002/jez.10043

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.28.538759; this version posted April 29, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Dreon, M. S., Ceolín, M., & Heras, H. (2007). Astaxanthin binding and structural stability of the apple snail carotenoprotein ovorubin. *Archives of Biochemistry and Biophysics*, *460*(1). https://doi.org/10.1016/j.abb.2006.12.033

Dreon, M. S., Heras, H., & Pollero, R. J. (2003). Metabolism of ovorubin, the major egg lipoprotein from the apple snail. *Molecular and Cellular Biochemistry*, *243*(1–2), 9–14. https://doi.org/10.1023/A:1021616610241

Dreon, M. S., Schinella, G., Heras, H., & Pollero, R. J. (2004a). Antioxidant defense system in the apple snail eggs, the role of ovorubin. *Archives of Biochemistry and Biophysics*, *422*(1). https://doi.org/10.1016/j.abb.2003.11.018

Dreon, M. S., Heras, H., & Pollero, R. J. (2004b). Characterization of the major egg glycolipoproteins from the perivitellin fluid of the apple snail Pomacea canaliculata. *Molecular Reproduction and Development*, *68*(3). https://doi.org/10.1002/mrd.20078

Falke, S., & Betzel, C. (2019). Dynamic Light Scattering (DLS). Principles, Perspectives, Applications to Biological Samples. In Pereira, Alice, S., P. Tavares, & P. Limão-Vieira (Eds.), *Radiation in Bioanalysis: Spectroscopic Techniques and Theoretical Methods* (pp. 173–193). Springer International Publishing. https://doi.org/10.1007/978-3-030-28247-9_6

Frassa, M. V., Ceolín, M., Dreon, M. S., & Heras, H. (2010). Structure and stability of the neurotoxin PV2 from the eggs of the apple snail Pomacea canaliculata. *Biochimica et Biophysica Acta - Proteins and Proteomics*, *1804*(7), 1492–1499. https://doi.org/10.1016/j.bbapap.2010.02.013

Gershenson, A., Gierasch, L. M., Pastore, A., & Radford, S. E. (2014). Energy landscapes of functional proteins are inherently risky. *Nature Chemical Biology*, *10*(11), 884–891. https://doi.org/10.1038/NCHEMBIO.1670

Giglio, M. L., Ituarte, S., Ibañez, A. E., Dreon, M. S., Prieto, E., Fernández, P. E., & Heras, H. (2020). Novel Role for Animal Innate Immune Molecules: Enterotoxic Activity of a Snail Egg MACPF-Toxin. *Frontiers in Immunology*, *11*(March), 1–14. https://doi.org/10.3389/fimmu.2020.00428

Giglio, M. L., Ituarte, S., Pasquevich, M. Y., & Heras, H. (2016). *The eggs of the apple snail Pomacea maculata are defended by indigestible polysaccharides and toxic*

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.28.538759; this version posted April 29, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

proteins. *785*(September), 777–785.

Hayes, K. A., Cowie, R. H., Albrecht, C., & Thiengo, S. C. (2009). Molluscan models in evolutionary biology: Apple snails (Gastropoda:Ampullariidae) as a system for addressing fundamental questions. *American Malacological Bulletin*, *27*(47), 58.

Heras, H., Dreon, M. S., Ituarte, S., & Pollero, R. J. (2007). Egg carotenoproteins in neotropical Ampullariidae (Gastropoda: Architaenioglossa). *Comp.Biochem.Physiol.*, *C 146*, 158–167. http://www.ncbi.nlm.nih.gov/pubmed/17320485

Heras, H., Frassa, M. V., Fernández, P. E., Galosi, C. M., Gimeno, E. J., & Dreon, M. S. (2008). First egg protein with a neurotoxic effect on mice. *Toxicon*, *52*(3), 481–488. https://doi.org/10.1016/j.toxicon.2008.06.022

Heras, H., Garin, C. F., & Pollero, R. J. (1998). Biochemical composition and energy sources during embryo development and in early juveniles of the snail Pomacea canaliculata (Mollusca: Gastropoda). *Journal of Experimental Zoology*, *280*(6), 375–383. https://doi.org/10.1002/(SICI)1097-010X(19980415)280:6<375::AID-JEZ1>3.0.CO;2-K

Ip, J. C. H., Mu, H., Zhang, Y., Sun, ., Heras, H., Chu, K. H., & Qiu, J. W. (2019). Understanding the transition from water to land: Insights from multi-omic analyses of the perivitelline fluid of apple snail eggs. *J Proteomics*, *194*, 79–88.

Ituarte, S., Brola, T. R., Fernández, P. E., Mu, H., Qiu, J.-W., Heras, H., & Dreon, M. S. (2018). A lectin of a non-invasive apple snail as an egg defense against predation alters the rat gut morphophysiology. *PLoS ONE*, *13*(6). https://doi.org/10.1371/journal.pone.0198361

Ituarte, S., Dreon, M. S., Ceolin, M., & Heras, H. (2012). Agglutinating Activity and Structural Characterization of Scalarin, the Major Egg Protein of the Snail Pomacea scalaris (d'Orbigny, 1832). *PLoS ONE*, *7*(11). https://doi.org/10.1371/journal.pone.0050115

Ituarte, S., Dreon, M. S., Ceolín, M., & Heras, H. (2008). Isolation and characterization of a novel perivitellin from the eggs of Pomacea scalaris (Mollusca, Ampullariidae). *Molecular Reproduction and Development*, *75*(9), 1441–1448. https://doi.org/10.1002/mrd.20880

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.28.538759; this version posted April 29, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Ituarte, S., Dreon, M. S., Pasquevich, M. Y., Fernández, P. E., & Heras, H. (2010). Carbohydrates and glycoforms of the major egg perivitellins from Pomacea apple snails (Architaenioglossa: Ampullariidae). *Comparative Biochemistry and Physiology - B Biochemistry and Molecular Biology*, *157*(1). https://doi.org/10.1016/j.cbpb.2010.05.004

Ituarte, S., Dreon, M. S., Pollero, R. J., & Heras, H. (2008). Isolation and partial characterization of a new lipo-glyco-carotenoprotein from Pomacea scalaris (Gastropoda: Ampullariidae). *Mol.Reprod.Dev.*, *75*(9), 1441–1448. http://www.ncbi.nlm.nih.gov/pubmed/18213678

Kim, B. M., Kim, H., Raines, R. T., & Lee, Y. (2004). Glycosylation of onconase increases its conformational stability and toxicity for cancer cells. *Biochemical and Biophysical Research Communications*, *315*(4), 976–983. https://doi.org/10.1016/J.BBRC.2004.01.153

Laemmli, U. K. (1970). Cleavage of Structural Proteins during the Assembly of the Head of Bacteriophage T4. *Nature Publishing Group*, *227*, 680–685. http://www.mendeley.com/research/discreteness-conductance-chnge-n-bimolecular-lipid-membrane-presence-certin-antibiotics/

Larkin, M. A., Blackshields, G., Brown, N. P., Chenna, R., Mcgettigan, P. A., McWilliam, H., Valentin, F., Wallace, I. M., Wilm, A., Lopez, R., Thompson, J. D., Gibson, T. J., & Higgins, D. G. (2007). Clustal W and Clustal X version 2.0. *Bioinformatics*, *23*(21), 2947–2948. https://doi.org/10.1093/bioinformatics/btm404

Manning, M., & Colón, W. (2004a). Structural basis of protein kinetic stability: Resistance to sodium dodecyl sulfate suggests a central role for rigidity and a bias toward β-sheet structure. *Biochemistry*, *43*(35), 11248–11254. https://doi.org/10.1021/bi0491898

Manning, M., & Colón, W. (2004b). Structural basis of protein kinetic stability: Resistance to sodium dodecyl sulfate suggests a central role for rigidity and a bias toward β-sheet structure. *Biochemistry*, *43*(35), 11248–11254.

Mehta, A. Y., & Cummings, R. D. (2019). GLAD: GLycan Array Dashboard, a visual analytics tool for glycan microarrays. *Bioinformatics*, *35*(18), 3536–3537. https://doi.org/10.1093/bioinformatics/btz075

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.28.538759; this version posted April 29, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Pasquevich, M. Y., Dreon, M. S., & Heras, H. (2014). The major egg reserve protein from the invasive apple snail Pomacea maculata is a complex carotenoprotein related to those of Pomacea canaliculata and Pomacea scalaris. *Comparative Biochemistry and Physiology - B Biochemistry and Molecular Biology*, *169*(1). https://doi.org/10.1016/j.cbpb.2013.11.008

Pasquevich, M. Y., Dreon, M. S., Qiu, J.-W., Mu, H., & Heras, H. (2017). Convergent evolution of plant and animal embryo defences by hyperstable non-digestible storage proteins. *Scientific Reports*, *7*(15848). https://doi.org/10.1038/s41598-017-16185-9

Pasquevich, M. Y., & Heras, H. (2020). Apple snail egg perivitellin coloration, as a taxonomic character for invasive Pomacea maculata and P. canaliculata, determined by a simple method. *Biological Invasions*, *22*(7), 2299–2307. https://doi.org/10.1007/s10530-020-02255-z

Sun, J., Mu, H., Ip, J. C. H., Li, R., Xu, T., Accorsi, A., Alvarado, A. S., Ross, E., Lan, Y., Sun, Y., Castro-Vazquez, A., Vega, I. A., Heras, H., Ituarte, S., Van Bocxlaer, B., Hayes, K. A., Cowie, R. H., Zhao, Z., Zhang, Y., … Qiu, J. W. (2019). Signatures of divergence, invasiveness, and terrestrialization revealed by four apple snail genomes. *Molecular Biology and Evolution*, *36*(7), 1507–1520. https://doi.org/10.1093/molbev/msz084

Tokuriki, N., & Tawfik, D. S. (2009a). Special sectionProtein Dynamism and Evolvability. *Science (New York, N.Y.)*, *324*(April), 203–207.

Tokuriki, N., & Tawfik, D. S. (2009b). Stability effects of mutations and protein evolvability. *Current Opinion in Structural Biology*, *19*(5), 596–604. https://doi.org/10.1016/j.sbi.2009.08.003

Zeldovich, K. B., Chen, P., & Shakhnovich, E. I. (2007). Protein stability imposes limits on organism complexity and speed of molecular evolution. *Proceedings of the National Academy of Sciences of the United States of America*, *104*(41), 16152–16157. https://doi.org/10.1073/pnas.0705366104

Zheng, J., Guo, N., & Wagner, A. (2020). Selection enhances protein evolvability by increasing mutational robustness and foldability. *Science*, *370*(6521). https://doi.org/10.1126/science.abb5962

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.28.538759; this version posted April 29, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Table 1. Size and molecular weight estimation of PV1s

	MW (kDa)	Diameter Size (nm)
PpaPV1	267.5	12.2
PsSC	270.3	13.0
PcOvo	269.8	13.4
PmPV1	265.6	12.4

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.28.538759; this version posted April 29, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Table 2. Thermodynamic parameters of *Pomacea* apple snail perivitellins unfolding induced by chemical treatment.

Clade	**Flagellata**	**Bridgesii**		**Canaliculata**	
	P. patula	*P. scalaris*	*P. diffusa*	*P. maculata*	*P. canaliculata*
PV1	**PpaPV1**	**PsSC**	PdPV1[£]	**PmPV1**	PcOVO[¥]
ΔG^0H_2O (kcal/mol)	**6.40 ± 0.32**	**1.43 ± 0.44**	1.14	**3.32 ± 0.19**	3.48 ± 0.06
Cm (M)	**5.4**	**4.1**	3.4	**2.8**	ND

[¥] Data from Dreon et al. 2007; [£] Data from Brola et al. 2020; ND not determined; In bold: data from this study (See Material and Methods for experimental details).

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.28.538759; this version posted April 29, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Table 3. Comparison of the Half-Lives and Unfolding Rate Constants of snail PV1s and proteins resistant and not resistant to SDS. Proteins were sorted according to their half-life.

Protein*	Unfolding Half-life	k_{unf} (sec-1) in 6.6 M GdnHCl	k_{unf} (sec-1) in 0 M GdnHCl
SDS-Resistant			
AVD	270 years	2.1×10^{-3}	8.1×10^{-11}
TTR	244 years	3.2×10^{-3}	9.0×10^{-11}
PAP	165 years	1.5×10^{-2}	1.3×10^{-10}
PpaPV1	**51 years**	**5.6×10^{-3} ¥**	**4.3×10^{-10}**
TSP	13 years	1.0×10^{-2}	1.6×10^{-9}
SOD	3.7 years	1.7×10^{-3}	6.0×10^{-9}
CPAP	2.5 years	1.6×10^{-4}	8.8×10^{-9}
SVD	318 days	1.0×10^{-4}	2.5×10^{-8}
SAP	79 days	4.0×10^{-2}	1.0×10^{-7}
PsSC	**7.1 days**	**3.5×10^{-4}**	**1.14×10^{-6}**
PmPV1	**2.3 days**	**2.3×10^{-2} ¥**	**3.5×10^{-6}**
SDS Non-resistant			
ADH	19 h	Unobservable	1.0×10^{-5}
TIM	15 h	Unobservable	2.8×10^{-5}
BLA	12 h	Unobservable	2.8×10^{-5}
β2M	24 min	Unobservable	4.9×10^{-4}
ConA	22 min	Unobservable	5.3×10^{-4}
GAPDH	14 min	Unobservable	8.2×10^{-4}

*ADH, yeast alcohol dehydrogenase; AVD, avidin; β2M, β2-microglobulin; BLA, bovine R-lactalbumin; ConA, concanavalinA; CPAP, chymopapain; GAPDH, glyceraldehyde 3-phosphate dehydrogenase; PAP, papain; SAP, serum amyloid P; SOD, copper/zinc superoxide dismutase; SVD, streptavidin; TSP, P22 tailspike protein; TIM, triosephosphate isomerase from porcine muscle; TTR, transthyretin. ¥K *unf* was meassured at 6.5 GdnHCl M. Although all PV1s are SDS resistant, Flagellata clade PV1 (PpaPV1) is much more stable than Bridgesii (PsSC) and Canaliculata (PmPV1) and is listed among the most stable proteins so far reported in nature. Taken from Manning & Colón, 2004 and modified with data from this study (in bold).

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.28.538759; this version posted April 29, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Table 4. Main glycan structures recognized by PpaPV1

Rank	Oligosaccharide Structure	Average RFU	%CV
1	GalNAca1-3(Fuca1-2)Galb1-4GlcNAcb1-2Mana1-6(GalNAca1-3(Fuca1-2)Galb1-4GlcNAcb1-2Mana1-3)Manb1-4GlcNAcb1-4GlcNAcb-Sp20	1081 ± 73	7
2	GalNAca1-3(Fuca1-2)Galb1-4GlcNAcb1-3Galb1-4GlcNAcb-Sp0	325 ± 40	12
3	Galb1-4GalNAca1-3(Fuca1-2)Galb1-4GlcNAcb-Sp8	318 ± 41	13
4*	Galb1-3GlcNAcb1-3Galb1-3GlcNAcb-Sp0	309 ± 3	1
5	Galb1-3GalNAca1-3(Fuca1-2)Galb1-4GlcNAc-Sp0	239 ± 28	12
6	GalNAca1-3(Fuca1-2)Galb1-4GlcNAcb1-2Mana-Sp0	204 ± 20	10
7*	(3S)Galb1-4(Fuca1-3)(6S)Glc-Sp0	163 ± 14	9

Binding intensities are expressed as the mean of relative fluorescence units (RFU) ± 1SD, N=4. %CV = 100 x SD. Full data of PpaPV1 glycan specificity is available as Supporting Information. *not PpaPV1 concentration-dependent binding.

bioRxiv preprint doi: https://doi.org/10.1101/2023.04.28.538759; this version posted April 29, 2023. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Figure 1




Figure 1.

Apple snail egg PV1s have similar size and number of subunits, but different electrophoretic migration, charge surface and immune cross-reactivity in a clade-related fashion.

A. Particle size analysis as determined by DLS indicate that all PV1s have similar size and MW. **B.** Oligomers of PV1s in a native PAGGE (top panel). Arrows indicate the relative mobility of PV1s. Blue arrow highlight the increase in mobility along clades. Subunits of PV1s in a SDS-PAGE (lower panel). **C**. Western blot analysis using Anti PsSC antiserum and Anti PcOvo antiserum. Line 1, molecular weight marker. Line 2, PpaPV1. Line 3, PsSC. Line 4, PmPV1. Line 5, PcOvo. *Fl: Flagellata; Br: Bridgesii; Ca: Canaliculata.* PV1 were purified from*:* PpaPV1 from *P. patula*; PsSC from *P. scalaris*; PmPV1 from *P. maculata*; PcOvo from *P. canaliculata*.

Figure 2




Figure 2. *Pomacea* eggs and PV1 absorption spectra shifts toward red in most derived clades. **A.** Egg extract (perivitelline fluid) **B.** purified egg carotenoproteins. Spectra are ordered from basal (top) to derived (bottom) clades. Arrows indicate the maximum of each spectrum to highlight the red-shift from basal to derived clades. Spectra were normalized to easy comparison. Data of *P. scalaris* egg carotenoprotein taken from Ituarte *et. al.* 2008.

Figure 3




Figure 3. Structural stability of PV1 decrease in a clade-related fashion.
Stability of PpaPV1, PsSC and PmPV1 was evaluated by the unfold induced by GdnHCl. **A.** Unfolded population of PV1s in the equilibrium. **B** Dependence of the unfolding free energy (ΔG^0) with GndHCl concentration. ΔG^0H_2O was calculated from the ordinate intercept. C_m : GndHCl concentration at $\Delta G^0=0$ (midpoint of the denaturing transition). PmPV1 data taken from Pasquevich *et al.* 2017.

Figure 4




Figure 4. The PV1s of basal clades are hyperstable proteins extremely resistant to detergent treatment and chemical denaturation. A. SDS-PAGE of PV1s previously unheated (U) or boiled (B) in the presence of SDS detergent for 10 min and immediately loaded into the gel. PpaPV1 is more resistant to detergent treatment than those PV1s from other clades. A comparison with other hyperstable proteins in nature is given in table 3. Fl: Flagellata; Br: Bridgesii; Ca: Canaliculata; **B.** Unfolding rates of PpaPV1 and PmPV1 under native-like conditions are shown by extrapolating the unfolding rate determined at different concentrations of GdnHCl to 0 M. The *y-intercept* of each extrapolation curve indicates the unfolding rate of the native protein. PpaPV1 has an unfolding kinetics much slower than the orthologue of the most derived clade.

Figure 5




Figure 5. **Lectin capacity of Flagellata PpaPV1 is not as strong as Bridgesii PV1s and has a narrower glycan binding motifs. A.** Microplate well showing the hemagglutinating activity of *P. patula* purified PpaPV1 and PmPV1 from *P. maculata.* Circled well correspond to the last dilution that hemagglutinate. The last well of each row corresponds to PBS. PpaPV1 has a moderate hemagglutinating activity while PV1 belonging to the derived canaliculata clade lacks this capacity. **B.** Main glycan structures recognized by PpaPV1 highlighting a common recognizing pattern: *GalNAc*a1-3*(Fuca1-2)Gal*b1-4*GlcNAc* (rectangle). Glycan structure plot was taken from the Consortium for Functional Glycomics (http://www.functionalglycomics.org). See Table 4 for more details.

Figure 6




Figure 6. **Evolution of structure, stability and functional features of PV1 carotenoproteins in *Pomacea* genus.** *This study. Dots indicate proteins studied for the trait. *Pomacea* phylogeny was based on Hayes *et al.* 2009.
Data was taken from Brola et al. 2020; Dreon et al. 2004a, 2004b; Ituarte *et. al.* 2008, 2010, 2012; Pasquevich et al., 2014, 2017.