Designing allosteric regulators of thrombin. Exosite 2 features multiple subsites that can...

Search within citing articles

On the process of discovering leads that target the heparin-binding site of neutrophil elastase in the sputum of cystic fibrosis patients

S Morla, NV Sankaranarayanan… - Journal of medicinal …, 2019 - pmc.ncbi.nlm.nih.gov

Cystic fibrosis (CF) is a disease of dysregulated salt and fluid homeostasis that results in massive accumulation of neutrophil elastase resulting in lung degradation and death. The …

☆ Save 99 Cite Cited by 20 Related articles All 7 versions

Targeting the GPIbα binding site of thrombin to simultaneously induce dual anticoagulant and antiplatelet effects

AY Mehta, JN Thakkar, BM Mohammed… - Journal of medicinal …, 2014 - ACS Publications

Exosite 2 of human thrombin contributes to two opposing pathways, the anticoagulant pathway and the platelet aggregation pathway. We reasoned that an exosite 2 directed …

☆ Save 99 Cite Cited by 31 Related articles All 7 versions

Development of orally active thrombin inhibitors for the treatment of thrombotic disorder diseases

LW He, WC Dai, NG Li - Molecules, 2015 - pmc.ncbi.nlm.nih.gov

Thrombotic disorders represent the major share of the various cardiovascular diseases, and significant progress has been made in the development of synthetic thrombin inhibitors as …

☆ Save 99 Cite Cited by 21 Related articles All 11 versions

Plasmin regulation through allosteric, sulfated, small molecules

RA Al-Horani, R Karuturi, DT White, UR Desai - Molecules, 2015 - mdpi.com

Plasmin, a key serine protease, plays a major role in clot lysis and extracellular matrix remodeling. Heparin, a natural polydisperse sulfated glycosaminoglycan, is known to …

☆ Save 99 Cite Cited by 28 Related articles All 16 versions »

De novo synthesis of benzofurans *via* trifluoroacetic acid catalyzed cyclization/oxidative aromatization cascade reaction of 2-hydroxy-1,4-diones

Q Sha, H Liu - Organic & Biomolecular Chemistry, 2019 - pubs.rsc.org

A facile, one-pot method for the de novo synthesis of benzofurans from 2-hydroxy-1, 4-diones is described. Using trifluoroacetic acid (TFA) as the catalyst and N-bromobutanimide …

☆ Save 99 Cite Cited by 17 Related articles All 6 versions

Novel heparin mimetics reveal cooperativity between exosite 2 and sodium-binding site of thrombin

MHA Aziz, UR Desai - Thrombosis research, 2018 - Elsevier

Introduction Thrombin is a primary target of most anticoagulants. Yet, thrombin's dual and opposing role in pro-as well as anti-coagulant processes imposes considerable challenges …

☆ Save 99 Cite Cited by 12 Related articles All 8 versions

Synthesis of glycosaminoglycan mimetics through sulfation of polyphenols

RA Al-Horani, R Karuturi, S Verespy III… - … : Chemistry and Biology, 2014 - Springer

In nearly all cases of biological activity of sulfated GAGs, the sulfate group (s) are critical for interacting with target proteins. A growing paradigm is that appropriate small, sulfated …

☆ Save 99 Cite Cited by 17 Related articles All 6 versions

Synthesis and biological evaluation of some dibenzofuran-piperazine derivatives

L Yurttaş, U Abu Mohsen, Y Ozkan… - Journal of Enzyme …, 2016 - Taylor & Francis

In the present paper, a novel series of dibenzofuran-piperazine derivatives were synthesized via the treatment of N-(2-methoxy-3-dibenzofuranyl)-2-chloroacetamide with …

☆ Save 99 Cite Cited by 15 Related articles All 12 versions

Merging cultures and disciplines to create a drug discovery ecosystem at Virginia commonwealth university: Medicinal chemistry, structural biology, molecular …

GE Kellogg, Y Cen, M Dukat, KC Ellis, Y Guo, J Li… - SLAS Discovery, 2023 - Elsevier

Abstract The Department of Medicinal Chemistry, together with the Institute for Structural Biology, Drug Discovery and Development, at Virginia Commonwealth University (VCU) has …

☆ Save 99 Cite Cited by 5 Related articles All 5 versions

Polyglycine hydrolases secreted by Pleosporineae fungi that target the linker region of plant class IV chitinases

TA Naumann, DT Wicklow, NPJ Price - Biochemical Journal, 2014 - portlandpress.com

Cmps (chitinase-modifying proteins) are fungal proteases that truncate plant class IV chitinases by cleaving near their N-termini. We previously described Fv-cmp, a fungalysin …

☆ Save 99 Cite Cited by 15 Related articles All 8 versions