Targeting the GPIbα binding site of thrombin to simultaneously induce dual anticoagulant...

Search within citing articles

Evaluating prodrug characteristics of a novel anticoagulant fusion protein neorudin, a prodrug targeting release of hirudin variant 2-Lys47 at the thrombosis site, by …

X Dong, R Gu, X Zhu, H Gan, J Liu, J Jin, Z Meng… - European Journal of …, 2018 - Elsevier

Recombinant neorudin (EPR-hirudin, EH), a low-bleeding anticoagulant fusion protein, is an inactive prodrug designed to be converted to the active metabolite, hirudin variant 2-Lys47 …

☆ Save 99 Cite Cited by 17 Related articles All 4 versions

Development of orally active thrombin inhibitors for the treatment of thrombotic disorder diseases

LW He, WC Dai, NG Li - Molecules, 2015 - pmc.ncbi.nlm.nih.gov

Thrombotic disorders represent the major share of the various cardiovascular diseases, and significant progress has been made in the development of synthetic thrombin inhibitors as …

☆ Save 99 Cite Cited by 21 Related articles All 11 versions

Detection of α-thrombin with platelet glycoprotein ibα (GP1bα) for the development of a coagulation marker

D Oh, YJ Chae, JY Teoh, B Yim, D Yoo, Y Park… - ACS …, 2024 - pmc.ncbi.nlm.nih.gov

The detection of prothrombotic markers is crucial for understanding thromboembolism and assessing the effectiveness of anticoagulant drugs. α-Thrombin is a marker that plays a …

☆ Save 99 Cite Cited by 3 Related articles All 8 versions

The progress in the research of antiplatelet agents (1995–2017)

O Benimana, L Zhao, Y Kong, Z Li… - Future Medicinal …, 2017 - Taylor & Francis

Antiplatelet therapy displays a critical role in the treatment and prevention of antithrombotic disorders. Many new antiplatelet agents have been developed following the emergence of …

☆ Save 99 Cite Cited by 18 Related articles All 3 versions

Merging cultures and disciplines to create a drug discovery ecosystem at Virginia commonwealth university: Medicinal chemistry, structural biology, molecular …

GE Kellogg, Y Cen, M Dukat, KC Ellis, Y Guo, J Li… - SLAS Discovery, 2023 - Elsevier

Abstract The Department of Medicinal Chemistry, together with the Institute for Structural Biology, Drug Discovery and Development, at Virginia Commonwealth University (VCU) has …

☆ Save 99 Cite Cited by 5 Related articles All 5 versions

Structure–activity relationship studies for multitarget antithrombotic drugs

AR Neves, M Correia-da-Silva, E Sousa… - Future medicinal …, 2016 - Taylor & Francis

Thrombosis is a complex process involving multiple pathways. Currently, therapy relies on the combination of two or more antithrombotic drugs, showing that inhibiting more than one …

☆ Save 99 Cite Cited by 13 Related articles All 3 versions

Thrombin Exosite Maturation and Ligand Binding at ABE II Help Stabilize PAR Binding Competent Conformation at ABE I

R Billur, TM Sabo, MC Maurer - Biochemistry, 2019 - pmc.ncbi.nlm.nih.gov

Thrombin, derived from zymogen prothrombin (ProT), is a serine protease involved in procoagulation, anticoagulation, and platelet activation. Thrombin's actions are regulated …

☆ Save 99 Cite Cited by 10 Related articles All 7 versions

Present and future prospective of lignin-based materials in biomedical fields

W Gao, F Kong, J Chen, P Fatehi - Lignin-Based Materials for Biomedical …, 2021 - Elsevier

Lignin is a bio-based material with tremendous applications in different fields. In this chapter, the potential use of lignin in the medical field is discussed. This chapter discusses the status …

☆ Save 99 Cite Cited by 8 Related articles All 4 versions

Deciphering conformational changes associated with the maturation of thrombin anion binding exosite I

R Billur, D Ban, TM Sabo, MC Maurer - Biochemistry, 2017 - pmc.ncbi.nlm.nih.gov

Thrombin participates in procoagulation, anticoagulation, and platelet activation. This enzyme contains anion binding exosites, ABE I and ABE II, which attract regulatory …

☆ Save 99 Cite Cited by 6 Related articles All 5 versions

Substantial non-electrostatic forces are needed to induce allosteric disruption of thrombin's active site through exosite 2

AY Mehta, UR Desai - Biochemical and biophysical research …, 2014 - Elsevier

Sulfated β-O4 lignin (SbO4L), a non-saccharide glycosaminoglycan mimetic, was recently disclosed as a novel exosite 2-directed thrombin inhibitor with the capability of mimicking …

☆ Save 99 Cite Cited by 9 Related articles All 10 versions