Recognition of highly branched N-glycans of the porcine whipworm by the immune system

Search within citing articles

N-glycoproteomic profiling of the hookworm Ancylostoma ceylanicum reveals heterogeneity and phosphorylcholine-modified glycans in larval and adult parasites.

CL Paraggio, V Chang, KE Mahoney, N Chung… - bioRxiv, 2026 - biorxiv.org

Glycans of parasitic worms (helminths) have been previously described with roles in immunomodulation and the recognition of helminth proteins by the host immune system …

☆ Save 99 Cite Related articles »

Application of Post-Glycosylation Modifying Enzymes for Mass Spectrometry Imaging of Modified N-Glycans In Situ

JE Dreifus, L Chuzel, EE Escobar… - Journal of the …, 2026 - pmc.ncbi.nlm.nih.gov

Glycans are essential components of cells involved in numerous biological processes, and changes in glycan profiles are often correlated with disease progression. Glycans are …

☆ Save 99 Cite Cited by 1 Related articles All 12 versions

Bioinformatic, Enzymatic, and Structural Characterization of Trichuris suis Hexosaminidase HEX-2

Z Dutkiewicz, A Varrot, KJ Breese, KA Stubbs… - …, 2024 - pmc.ncbi.nlm.nih.gov

Hexosaminidases are key enzymes in glycoconjugate metabolism and occur in all kingdoms of life. Here, we have investigated the phylogeny of the GH20 glycosyl hydrolase family in …

☆ Save 99 Cite Related articles All 13 versions

Discovering and Characterizing Conserved Anthelmintic Targets in the Early Parasitic Stages of Gastrointestinal Nematodes

H Jung - 2024 - search.proquest.com

Gastrointestinal nematodes (GINs) collectively infect 1.5 billion people, leading to mortality and various forms of morbidity in humans and hindering normal development in children …

☆ Save 99 Cite Related articles All 3 versions

Core Tri-fucosylation of Nematode N-glycans Requires Golgi α-mannosidase III Activity that Impacts Animal Growth and Behaviours

J Kendler, F Wöls, S Thapliyal, E Arcalis, H Gabriel… - bioRxiv, 2024 - biorxiv.org

Many nematodes possess N-glycans with complex core chitobiose modifications, which is a feature observed in various free-living and parasitic nematodes but is absent in mammals …

☆ Save 99 Cite Related articles All 2 versions »

Ad ancing helminth gl comics: structural specificit and immunogenicit of schistosomal and filarial gl cans

LMC Petralia - scholarlypublications …

Helminths are parasitic macroorganisms belonging to the two major phyla of nematodes (roundworms) or platyhelminths (flatworms) 1. The latter include trematodes (flukes) and …

☆ Save 99 Cite Related articles All 2 versions »