Chemokine binding to PSGL-1 is controlled by O-glycosylation and tyrosine sulfation

Search within citing articles

Truncated O-glycosylation in metastatic triple-negative breast cancer reveals a gene expression signature associated with extracellular matrix and proteolysis

MF Festari, E Jara, M Costa, A Iriarte, T Freire - Scientific Reports, 2024 - academia.edu

Breast cancer (BC) is the leading cause of death by cancer in women worldwide. Triple-negative (TN) BC constitutes aggressive and highly metastatic tumors associated with …

☆ Save 99 Cite Related articles All 4 versions »

Charge Matters: How flanking substrate charge and silicates modulate O-glycan biosynthesis

CJ Ballard - 2025 - search.proquest.com

Mucin type O-glycosylation is a ubiquitous and essential form of posttranslational modification of proteins initiated by family of Golgi resident transferases known as the …

☆ Save 99 Cite Related articles All 2 versions

Parallel Synthesis of Glycopeptide Libraries Using Glyco-SPOT Synthesis

AY Mehta, J Heimburg-Molinaro… - … Libraries: Methods and …, 2025 - Springer

Synthesis of glycopeptide libraries is often limited by the cost and availability of the glycoamino acids. To circumvent this, we have devised a method of using SPOT synthesis …

☆ Save 99 Cite Related articles All 5 versions

CCR5 Sulfopeptide Library and O-Glycosylated Amino Acids as Chemical Tools for Probing Chemokine Receptor Recognition

MMA Eldeen - 2026 - search.proquest.com

Chemokine-receptor interactions play a central role in immune regulation and disease pathogenesis, particularly through CCR5, a receptor engaged by chemokines such as …

☆ Save 99 Cite Related articles

An in vitro one-pot synthetic biology approach to simulating diverging Golgi O-glycosylation of tumor-associated MUC1 from normal tissue MUC1

A Nashed, K Dilsook, T Senapathi, K Naidoo - 2024 - chemrxiv.org

Peptide O-glycosylation is a non-template-driven process that relies on the coordinated action of glycosyltransferases (GTs) within the endoplasmic reticulum (ER) and Golgi …

☆ Save 99 Cite Related articles All 6 versions

GALNT4 promotes the endothelial cell inflammatory response via the NF-κB signaling pathway

LW Guo, L Zhou, P Wei, S Li, D Li - 2024 - assets-eu.researchsquare.com

Background Atherosclerosis (AS) is a chronic in ammatory disease caused by dysfunction of vascular endothelial cells (ECs). Polypeptide N-acetylgalactosaminyl-transferase 4 …

☆ Save 99 Cite Related articles All 3 versions »

Abdullateef Nashed 1, 2 Kyllen Dilsook 1, 2 Tharindu Senapathi 1, 3 AA

KJ Naidoo - assets-eu.researchsquare.com

Peptide O-glycosylation relies on the coordinated action of glycosyltransferases (GTs) within the endoplasmic reticulum (ER) and Golgi apparatus. An in vitro one-pot synthetic biology …

☆ Save 99 Cite Related articles »

糖基化修饰对蛋白质和肽性质的影响及修饰方法

郑小菊， 刘长松， 王凤山 - 生命的化学 - sciengine.com

糖基化修饰在生物体内扮演着重要的角色, 广泛参与细胞黏附, 分子运输, 信号传导, 受体激活等各种生理过程, 同时也与癌症, 感染性疾病, 糖尿病等多种疾病的发生发展密切相关 …

☆ Save 99 Cite Related articles All 2 versions »