Allosteric partial inhibition of monomeric proteases. Sulfated coumarins induce regulation,...

Search within citing articles

A synthetic heparin mimetic that allosterically inhibits factor XIa and reduces thrombosis in vivo without enhanced risk of bleeding

RA Al-Horani, EI Abdelfadiel, DK Afosah… - … of Thrombosis and …, 2019 - Wiley Online Library

Background Human factor XI a (FXI a) is an actively pursued target for development of safer anticoagulants. Our long-standing hypothesis has been that allosterism originating from …

☆ Save 99 Cite Cited by 38 Related articles All 11 versions »

Phytochemical and Analytical Characterization of Novel Sulfated Coumarins in the Marine Green Macroalga *Dasycladus vermicularis* (Scopoli) Krasser

A Hartmann, M Ganzera, U Karsten, A Skhirtladze… - Molecules, 2018 - mdpi.com

The siphonous green algae form a morphologically diverse group of marine macroalgae which include two sister orders (Bryopsidales and Dasycladales) which share a unique …

☆ Save 99 Cite Cited by 36 Related articles All 11 versions »

Characteristics of allosteric proteins, sites, and modulators

X He, D Ni, S Lu, J Zhang - Protein Allostery in Drug Discovery, 2019 - Springer

Allostery is considered one of the most direct and efficient ways to regulate biological macromolecule functions. Allostery is increasingly receiving attention in the field of drug …

☆ Save 99 Cite Cited by 29 Related articles All 6 versions

Discovery of benzyl tetraphosphonate derivative as inhibitor of human factor Xia

S Kar, M Mottamal, RA Al-Horani - ChemistryOpen, 2020 - Wiley Online Library

The inhibition of factor XIa (FXIa) is a trending paradigm for the development of new generations of anticoagulants without a substantial risk of bleeding. In this report, we present …

☆ Save 99 Cite Cited by 23 Related articles All 11 versions

Thermodynamics, kinetics and mechanisms of noncompetitive allosteric inhibition of chymotrypsin by dihydrolipoic acid-coated gold nanoclusters

WQ Chen, MM Yin, PJ Song, XH He, Y Liu, FL Jiang - Langmuir, 2020 - ACS Publications

Enzymes are an important class of biomacromolecules which catalyze many metabolic processes in living systems. Nanomaterials can be synthesized with tailored sizes as well as …

☆ Save 99 Cite Cited by 26 Related articles All 6 versions

Quantitative proteomics analysis reveals core and variable tick salivary proteins at the tick-vertebrate host interface

C Bensaoud, S Tenzer, A Poplawski… - Molecular …, 2022 - Wiley Online Library

Few studies have examined tick proteomes, how they adapt to their environment, and their roles in the parasite–host interactions that drive tick infestation and pathogen transmission …

☆ Save 99 Cite Cited by 18 Related articles All 9 versions

Potent, selective, allosteric inhibition of human plasmin by sulfated non-saccharide glycosaminoglycan mimetics

DK Afosah, RA Al-Horani… - Journal of medicinal …, 2017 - pmc.ncbi.nlm.nih.gov

Although plasmin inhibitors could be used in multiple disorders, their use has been restricted to preventing blood loss in hemostatic dysregulation because of poor efficacy and …

☆ Save 99 Cite Cited by 39 Related articles All 4 versions

Mechanistic insights into ultrasmall gold nanoparticle–protein interactions through measurement of binding kinetics

RS Ferreira, AL Lira, RJS Torquato… - The Journal of …, 2019 - ACS Publications

Gold nanoparticles (NPs) in the ultrasmall size regime provide a new paradigm in the way that nanomaterials can be used to regulate protein structure and function. However, the …

☆ Save 99 Cite Cited by 26 Related articles All 6 versions

Sulfated non-saccharide glycosaminoglycan mimetics as novel drug discovery platform for various pathologies

DK Afosah, RA Al-Horani - Current medicinal chemistry, 2020 - benthamdirect.com

Glycosaminoglycans (GAGs) are very complex, natural anionic polysaccharides. They are polymers of repeating disaccharide units of uronic acid and hexosamine residues. Owing to …

☆ Save 99 Cite Cited by 25 Related articles All 7 versions

Synthesis and study of new 2*H*-pyranoquinolin-2-one-based inhibitors of blood coagulation factors Xa and XIa

AY Potapov, BV Paponov, NA Podoplelova… - Russian Chemical …, 2021 - Springer

Abstract Condensation of 7-hydroxy-1, 2, 2, 4-tetramethylhydroquinoline-6-carbaldehydes with various hetaryl-substituted 3-oxo esters afforded a series of hybrid hydro-6, 8, 8, 9 …

☆ Save 99 Cite Cited by 18 Related articles All 4 versions