Reusable glycan microarrays using a microwave assisted wet-erase (MAWE) process

Search within citing articles

Glycan-mediated interactions of infectious hypodermal and hematopoietic necrosis virus (IHHNV) with host tissues in Pacific whiteleg shrimp (*Penaeus vannamei)*

C Sombutkayasith, A Watthammawut… - …, 2026 - academic.oup.com

Infectious hypodermal and hematopoietic necrosis virus (IHHNV) poses a significant threat to global shrimp aquaculture, causing substantial economic losses due to Runt Deformity …

☆ Save 99 Cite Related articles All 3 versions

Untersuchung von Anti-Glykan-Antikörpern und Erforschung ihres medizinischen Potenzials mithilfe synthetischer Glykane

F Weber, A Valenti, J Ledvinka, O Moscovitz… - Angewandte …, 2026 - Wiley Online Library

ZUSAMMENFASSUNG Die dynamische Glykan-Schicht, die menschliche und mikrobielle Zellen umgibt, spielt eine wesentliche Rolle für die Fähigkeit des Immunsystems …

☆ Save 99 Cite Related articles

Glycan-reactive antibodies isolated from human HIV-1 vaccine trial participants show broad pathogen cross-reactivity

PJ Jamieson, X Shen, AA Abu-Shmais… - Journal of …, 2025 - journals.asm.org

ABSTRACT HIV-1 continues to pose a significant global health challenge, requiring ongoing research into effective prevention and treatment strategies. Understanding the B-cell …

☆ Save 99 Cite Cited by 2 Related articles All 10 versions

Establishing “Glycoantigens” as a cross-field concept in Glycobiology, transfusion, transplant, and immunology

WJ Lane - Human Immunology, 2026 - Elsevier

Cell-surface glycans (carbohydrate structures) serve as critical antigens in transfusion and transplantation, yet their study remains fragmented across disciplines. The ABO blood group …

☆ Save 99 Cite Related articles All 2 versions

STRUCTURE-FUNCTION STUDIES OF AN RTX-ADHESIN FROM VIBRIO VULNIFICUS AND ITS ROLE IN BINDING HOST CELL SURFACES

BP Soares - 2026 - queensu.scholaris.ca

Numerous Gram-negative pathogenic bacteria employ repeat-in-toxin-adhesins to colonize a niche and initiate biofilm formation. Surface adhesion can occur via specific ligand-binding …

☆ Save 99 Cite Related articles All 2 versions