Journal Pre-proof



An Evolutionary Distinct Nipah Virus N-Glycosylation Site Provides Stability for Receptor Engagement

Tia E. Hawkins, Valeria Calvaresi, Sean A. Burnap, Yana Demyanenko, Liang Wu, Weston B. Struwe

PII: S1535-9476(26)00027-7

DOI: https://doi.org/10.1016/j.mcpro.2026.101531

Reference: MCPRO 101531

To appear in: *Molecular & Cellular Proteomics*

Received Date: 3 July 2025

Revised Date: 9 February 2026

Accepted Date: 15 February 2026

Please cite this article as: Hawkins TE, Calvaresi V, Burnap SA, Demyanenko Y, Wu L, Struwe WB, An Evolutionary Distinct Nipah Virus N-Glycosylation Site Provides Stability for Receptor Engagement, *Molecular & Cellular Proteomics* (2026), doi: https://doi.org/10.1016/j.mcpro.2026.101531.

This is a PDF of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability. This version will undergo additional copyediting, typesetting and review before it is published in its final form. As such, this version is no longer the Accepted Manuscript, but it is not yet the definitive Version of Record; we are providing this early version to give early visibility of the article. Please note that Elsevier's sharing policy for the Published Journal Article applies to this version, see: https://www.elsevier.com/about/policies-and-standards/sharing#4-published-journal-article. Please also note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2026 THE AUTHORS. Published by Elsevier Inc on behalf of American Society for Biochemistry and Molecular Biology.

Journal Pre-proof

An Evolutionary Distinct Nipah Virus N-Glycosylation Site Provides Stability for Receptor Engagement

Authors

Tia É. Hawkins[a,b,c], Valeria Calvaresi[b,d], Sean A. Burnap[b,d], Yana Demyanenko[a], Liang Wu[a,e*], Weston B. Struwe[b,d*]

Addresses

a. The Rosalind Franklin Institute, Harwell Science & Innovation Campus, Harwell OX11 0FA, U.K.

b. Kavli Institute for Nanoscience Discovery, University of Oxford, OX1 3QU, U.K.

c. Department of Chemistry, University of Oxford, Oxford OX1 3TA, U.K.

d. Department of Biochemistry, University of Oxford, Oxford, OX1 3QU, U.K.

e. Division of Structural Biology, Nuffield Department of Medicine, University of Oxford, Oxford, OX3 7BN, U.K.

Correspondents

*liang.wu@rfi.ac.uk, *weston.struwe@bioch.ox.ac.uk

Journal Pre-proof

Abbreviations:

2-AA, 2-anthranilic acid

AmBic, ammonium bicarbonate

BioNJ, biological neighbour joining

BirA, *E. coli* biotin ligase

BLI, biolayer interferometry

BP, beta propellers

DC-SIGN, dendritic cell-specific intercellular adhesion molecule-3-grabbing non-integrin

EB2, ephrin B2

Endo-H, endoglycosidase H

Env, envelope protein

EtHCD, electron-transfer high energy collision induced dissociation

F/F_M, Nipah virus fusion glycoprotein (subscript M denotes Malaysia strain)

FA, formic acid

FDR, false discovery rate

G/G_M, Nipah virus attachment glycoprotein (subscript M denotes Malaysia strain)

HCD, high energy collision induced dissociation

HDX, hydrogen deuterium exchange

HILIC-HPLC, hydrophilic interaction liquid chromatography-high performance liquid chromatography

Journal Pre-proof

IM-MS, ion mobility mass spectrometry

JTT, Jones-Taylor-Thornton

K_d, dissociation constant

LB, Luria broth

LFQ, label free quantification

L-SIGN, liver/lymph node-specific intercellular adhesion molecule-3-grabbing non-integrin

Man, mannose

MeCN, acetonitrile

MP, mass photometry

MWCO, molecular weight cut off

OST, oligosaccharyltransferase complex

PES, polyethersulfone

pH/D, estimated pH value calculated for a deuterated sample

PSM, peptide spectral match

PNGase-F, peptide:endoglycosidase F

PNGase-Rc, peptide:endoglycosidase *Rudaea cellulosilytica*

RT, room temperature

RefSeq, reference sequence

SEC, size exclusion chromatography

SP3, single-pot, solid-phase-enhanced sample-preparation

Journal Pre-proof

Journal Pre-proof

TCEP, tris(2-carboxyethyl)phosphine

TWIMS-TOF, travelling wave ion mobility spectrometry - time of flight

Abstract:

Nipah virus is a deadly paramyxovirus with 40-75% mortality and >750 cases since 1998. Currently there are no clinically approved vaccines or therapeutics to treat infection. Nipah is an enveloped virus with two surface glycoproteins, the trimeric fusion glycoprotein (F), and the tetrameric attachment glycoprotein (G), which is responsible for cellular attachment via binding to the host ephrin B2/B3 receptor. Glycosylation can substantially affect immunogenicity, receptor binding and structural conformations for virus glycoproteins, but its effects on Nipah G receptor engagement have not been studied. Here, phylogenetic and mass spectrometry analysis of the Nipah G Malaysia strain reveal how N-glycosylation has evolved since the appearance of the virus in 1998. We discovered that the N481 sequon is not conserved and the threonine/serine in the glycosylation site is critical for maintaining long-range stability of G subunits that facilitates ephrin B2 binding affinity. Together, these data reveal plasticity of N-glycosylation sites across Nipah species and the presence of hydrogen bonding networks that contribute to G stability and host engagement – results that are valuable for understanding virus attachment/entry mechanisms and the rationale design of structure-based vaccines.

Journal Pre-proof

Introduction:

Nipah virus is a zoonotic paramyxovirus first reported in 1998 following outbreaks of severe encephalitis in Malaysia and Singapore[1-3], with near-yearly outbreaks currently occurring in Bangladesh[4-6]. Nipah disease progression begins with fever, myalgia, nausea and coughing before rapid onset of encephalitis, seizures, coma and death[1,6]. The difficulty of Nipah diagnosis in the early stages of infection, coupled with a lack of approved therapeutics or vaccines, contributes to its high mortality rate (40-75%)[5]. Consequently, Nipah has been placed on the WHO R&D blueprint for emerging pathogens and epidemics[7].

Nipah virus entry and infection are not well understood, and the lack of effective treatment regimens reflect this. Nipah virus harbours two glycosylated surface proteins, the tetrameric attachment glycoprotein[8] (G) that binds to the host receptor ephrin B2 or B3 via its globular head domains[9-11] and the trimeric fusion glycoprotein[12] (F) that mediates viral fusion[13,14] (**Fig. 1a**). While G does not display neuraminidase activity, its head domains adopt a classical neuraminidase structure, with a six-bladed beta-propeller fold[3,15,16]. F is a class I fusion protein, activated by cathepsin L and B cleavage in the endosome[17]. G binds ephrin B2 and B3 with nanomolar affinity for each head domain[10], and picomolar affinity for the tetramer[18] due to avidity effects. Recent data suggests the extension of a G protein head outward towards the F protein could induce the F conformational changes that lead to fusion pore formation[19]. Despite several studies of host cell attachment and entry, there remains a general lack of understanding of structure-function relationships involving G and F glycoproteins, which can inform effective treatment regimens.

Currently, there are four candidate Nipah vaccines in phase I trials, including a live attenuated (PHV02)[20], mRNA (mRNA-1215)[21], adenoviral vector (ChAdOx-1 NiV)[22] and recombinant protein (HeV-G-sV)[23] vaccines[24]. While these candidates use different routes to elicit an immune response, vaccine-mediated immunity is acquired through exposure to Nipah G and F glycoproteins. Furthermore, the only biotherapeutic currently in clinical trials is the monoclonal

Journal Pre-proof

antibody m102.4, which targets the Nipah G protein[25]. Defining Nipah G and F glycosylation, as well as conserved areas of vulnerability, will aid therapeutic and vaccine development.

The Malaysia strain of Nipah G (G_M) has 28 predicted N-glycan sites per tetramer, and F (F_M) has 15 per trimer. N-glycan sites are genetically encoded through an N-X-S/T sequon that are subject to both positive and negative selection pressures[26-28]. Despite their prevalence, there is limited research into the roles of glycosylation for Nipah virus protein function[16,29]. Glycosylation of G has been suggested to be important for viral fusion[29-31], but a site-specific characterisation of glycan structure and the associated effects on receptor engagement has not been performed. Here, we examine the effects of Nipah G N-glycosylation on receptor binding using mass spectrometry (glycomics, glycoproteomics and hydrogen deuterium exchange (HDX)), mass photometry (MP) and biolayer interferometry (BLI) with phylogenetic analysis to uncover how a single glycan site governs structural dynamics and stability of a receptor binding complex.

Journal Pre-proof

Experimental Procedures:

Phylogenetic analysis of Nipah G isolates

81 Nipah G isolates and the Hendra G RefSeq (AAC83193, NP_047112) were exported as fasta sequences from the NCBI Virus database (https://www.ncbi.nlm.nih.gov/labs/virus/vssi/#/, accessed 31/03/2025) (Supplemental Tables S1-S3, Supplemental Data File 1). Evolutionary analyses were conducted in MEGA11[32] and the evolutionary history was inferred by using the Maximum Likelihood method and JTT matrix-based model[33]. Initial tree(s) for the heuristic search were obtained automatically by applying Neighbour-Join and BioNJ algorithms to a matrix of pairwise distances estimated using the JTT model, and then selecting the topology with superior log likelihood value. There were a total of 604 positions in the final dataset[32].

Journal Pre-proof

DNA production and purification

NiVG 71-602, NiVG 183-602, ephrin B2 28-165 and BirA constructs in pOPIN vectors for mammalian expression were kindly gifted by Raymond Owens and Joseph Thrush (Supplemental Table S4). DNA was transformed into DH5α NEB competent cells (New England Bioscience), and purified at scale using the Qiagen Gigaprep kit, following manufacturers' protocols. The coding sequence of each plasmid was verified by Sanger sequencing prior to use.

Site directed mutagenesis and 3C site cleavage insertion

Site directed mutagenesis and insertion of a 3C cleavage site was carried out using PCR with Q5 polymerase via the manufacturer's instructions (New England Biolabs). Primers were designed so codon mutations were in the forward primer only to generate linear DNA with non-overlapping primers (Supplemental Table S5). Subsequent linearised plasmids were ligated using KLD enzyme mix following manufacturer's instructions (New England Biolabs). The 3C cleavage site was inserted by separately linearising the NiVG 71-602 vector backbone and CDS using overlapping primers where one contained the 3C sequence (Supplemental Table S6). The linearised insert and vector were purified by gel extraction (Qiagen) and ligated using the In-Fusion enzyme kit (Takara Biosciences) as per the manufacturer's instructions. Constructs were transformed into chemically competent DH5α cells. DNA was purified and verified as described above.

Protein Expression and Purification

Expi293F™(Gibco) suspension cells cultured in Expi293™ expression medium (Gibco) were seeded to 1 x10^6 cells per mL and placed at 37°C, 5% CO_2, with shaking, 24 hours prior to transfection. For transfection, a solution amounting to 10% of the final transfection culture volume of OptiMem (Gibco), DNA (1 μg/mL final concentration) and PEI 40kDa (5.4 μg/mL final concentration) (Polysciences) was added to the culture. 18 hours post transfection, valproic acid

(5.9 mM final concentration), sodium propionate (6.8 mM final concentration) and glucose (50 mM final concentration) were added (biotin (840 µM final concentration) was also added to cultures containing BirA) before incubating for a further 6 days. Conditioned media was harvested (6000 *g,* 30 min, 4°C) and the supernatant loaded onto a Histrap Excel 1 mL (Cytiva). Proteins were washed in PBS pH 7.4, 20 mM imidazole before eluting with 1 M imidazole in PBS pH 7.4 with a gradient elution of 0-100% over 15 column volumes. Ephrin B2 purification was further supplemented with a high salt wash of PBS, pH 7.4, 20 mM imidazole, 300 mM NaCl prior to elution. Fractions containing proteins of interest were pooled and concentrated in 10 kDa (ephrin B2), 30 kDa (Nipah G head and mutants) or 50 kDa (Nipah G ectodomain) MWCO PES concentrators (Vivaspin) to 600 µL (4000 *g*, 4°C). The Nipah G 3C construct was incubated with hexahistidine tagged HRV 3C protease for 18 hrs, 4°C, at a ratio of 10:1 mg Nipah G: 3C. The concentrate was loaded onto a Superdex 200 increase 10/300 GL (Cytiva) pre-equilibrated and run in PBS pH 7.4 at 0.75 mL/min. For the 3C cleaved Nipah G protein, a 1 mL Histrap Excel was connected in tandem with the SEC column. Desired peak fractions were analysed via SDS-PAGE and concentrated as described above, and snap frozen in liquid nitrogen before storing at –80°C. Herein, Nipah G protein which has been 3C cleaved is described as "3C cleaved Nipah G_M ectodomain".

Mass Photometry

Mass Photometry measurements were carried out on a OneMP™ (Refeyn Ltd). Sample chambers were assembled by placing CultureWell™ gaskets (3 mm x 1mm, Grace Bio-Labs) on glass coverslips (High Precision No. 1.5H, Marienfeld Superior). 18 µL of PBS pH 7.4 was placed in a gasket to find focus, before addition of 2 µL of sample. Data were collected using Acquire MP (2023 R1.1, Refeyn Ltd). Nipah G 71-602 was measured at a final concentration of 5 nM with a medium field of view, exposure time = 1.91 ms, frame rate = 509.5 Hz, frame binning = 3, pixel binning = 4. Nipah 183-602 was measured at a final concentration of 10 nM with a small field of

view, exposure time = 0.95 ms, frame rate = 999.0 Hz, frame binning = 10, pixel binning = 4. Movies were recorded for 60s. Mass calibrations were performed for each field of view using an in-house protein standard, with peaks at 90, 180, 360 and 540 kDa. Data were analysed in Discover MP (v2023 R1.2, Refeyn Ltd), including gaussian fits and histogram generation. Data was exported and plotted in GraphPad Prism v10.2.3.

HPLC of N-glycans

10 µg Nipah G_M ectodomain was run on a 4-12% Bis-Tris SDS PAGE gel (Invitrogen) in biological quintuplicate. Gel bands were extracted and de-stained in 50 mM ammonium bicarbonate (AmBic) 50% acetonitrile (MeCN), incubated at 37°C for 30 min and washed in 50 mM AmBic 50% MeCN. Gel bands were dehydrated in 100% MeCN before covering with 700 µL containing PNGaseF at 12.5 ng/µL (in house). Digests were incubated at 4°C for 1 h, before moving to 37°C for a further 18 h.

Following digest, the PNGaseF reaction mixture was pooled with a subsequent three washes of gel bands with 100 µL of HPLC grade water (10 min, 37°C). N-glycans were dried and resuspended in 30 µL of HPLC grade water. 80 µL of freshly prepared 2-AA reaction mix (30 mg/mL 2-anthracillic acid, 45 mg/mL $NaBH_3CN$, 4% w/v $CH_3COONa{\cdot}3H_2O$, 2% w/v $B(OH)_3$; in MeOH) was added to glycans and incubated at 80°C for 1 h. Samples were cooled before clean-up with Spe-ed amide 2 cartridges. Samples were loaded with 1 mL 97% ACN, washed twice with 1 mL 95% ACN, then eluted twice with 750 µL HPLC grade water and lyophilised for 18 h.

Labelled glycans were resuspended in 20 µL HPLC grade water, and half of the glycans were further treated with 500 NEB units of Endo-H in 1X GlycoBuffer 3 (New England Bioscience), before incubating at 37°C for 22 h. The Endo-H reaction mix was cleaned up with 10 kDa MWCO cartridges (Omega), before drying in a vacuum concentrator and resuspending in HPLC grade water.

MeCN was added to labelled glycans to a final concentration of 70% MeCN. 5 µL of glycans were separated using an ACQUITY UPLC BEH Amide column (1.7 µM, 2.1x150 mm) coupled to a 1290 Infinity II UHPLC (Agilent) (Supplemental Table S7). Dextran, Man_5-$GlcNAc_2$ and Man_9-$GlcNAc_2$ glycan standards (Ludger) were run to help identify N-glycan peaks. Glycan peak identifications and quantification can be found in Supplemental Data File 2.

IM-MS/MS of N-glycans

40 µg of Nipah G was run on a 4-12% Bis-Tris SDS PAGE gel (Invitrogen) in biological triplicate. Gel bands were extracted and de-stained in 50 mM ammonium bicarbonate (AmBic) 50% acetonitrile (MeCN), incubated at 37°C for 30 min and washed in 50 mM AmBic 50% MeCN. Gel bands were dehydrated in 100% MeCN before covering with 700 µL 50 mM AmBic containing PNGaseF at 12.5 ng/µL (made in-house). Reaction was incubated at 4°C for 1 h, before moving to 37°C for a further 18 h. Released N-glycans were desalted using C18 zip tips (Merck), manually overlayed with porous graphite carbon (extracted from a Hypercarb™ porous graphite carbon column), then dried down and resuspended in 50:50 MeOH:H_2O.

Approximately 2 µL of N-glycan sample material was injected by static nanoelectrospray into a Synapt XS (Waters) using gold-coated borosilicate glass capillaries (prepared in-house). Data acquisition settings are presented in Supplemental Table S8. Data acquisition and processing were carried out using Waters Driftscope (version 2.8) software and MassLynx™ (version 4.1). IM-MS peak lists can be found in Supplemental Data File 3, while MS assignments and glycan compositions are listed in Supplemental Tables S9-11.

N-linked LFQ Glycoproteomics

3 µg of Nipah G_M ectodomain was run on a 4-12% Bis-Tris SDS PAGE gel (Invitrogen) in biological sextuplicate. 3 µg of Nipah G_M head domain, N481D and T483A mutants were also run on an SDS-PAGE gel in biological triplicate. Gel bands of interest were extracted and de-stained in 50

mM AmBic 50% MeCN and incubated at 37°C for 30 min, before washing in 50 mM AmBic 50% MeCN. Gel bands were dehydrated in 100% MeCN before adding 10 mM dithiothreitol in 100 mM AmBic and incubating for 30 min at 56°C. Gel bands were then dehydrated in 80% MeCN for 10 min at RT. Dithiothreitol solution was replaced with 50 mM iodoacetamide in 100 mM AmBic and the immersed gel bands were incubated in the dark for 30 min. Iodoacetamide solution was removed, and gel bands were washed sequentially with 100% MeCN and 50 mM AmBic. Gel bands were dehydrated 100% MeCN before addition of 12.5 ng/µL of the relevant protease (trypsin (Promega), chymotrypsin (Promega) or alpha-lytic (New England Bioscience)) in 50 mM AmBic. Gel bands were incubated with the protease solution at 37°C for 1 h before replacing the solution with fresh 50 mM AmBic and incubating the bands at 37°C for a further 18 h. The resulting peptide solution was pooled with sequential washes of 5% (v/v) FA in H_2O and MeCN. Samples were dried in a vacuum concentrator and reconstituted in 2% MeCN, 0.05% TFA for analysis by LC-MS/MS.

Samples were analysed using an Ultimate 3000 RSLCnano system coupled to an Orbitrap Q Exactive mass spectrometer (Thermo Fisher Scientific). Peptides were loaded onto a 75 µm × 2 cm pre-column and separated on a 75 µm × 15 cm Pepmap C18 analytical column (Thermo Fisher Scientific). Buffer A was 0.1% FA in H_2O and buffer B was 0.1% FA in 80% MeCN with 20% H_2O. A 100-minute linear gradient (Supplemental Table S12) and universal HCD identification method (Supplemental Table S13) was used.

O-linked Glycoproteomics

10 µg of 3C cleaved Nipah G_M ectodomain was run on a 4-12% Bis-Tris SDS PAGE gel (Invitrogen) in biological triplicate. Samples were then prepared for LC-MS/MS analysis in the same way as the N-linked glycoproteomics samples (described above).

Journal Pre-proof

Journal Pre-proof

Samples were analysed using an Ultimate 3000 RSLCnano system coupled to an Orbitrap Ascend mass spectrometer (Thermo Fisher Scientific). Peptides were loaded onto a C18 PepMap100 trap column (300 µm x 5 mm, 100 Å, Thermo Fisher Scientific) and separated on an in-house packed Reprosil-Gold C18 analytical column (50 µm × 50 cm, 1.9 µm particle size, Dr Maisch). Buffer A was 0.1% FA, 5% DMSO in H_2O, buffer B was 0.1% FA, 5% DMSO in MeCN and the loading solvent was 0.1% FA in H_2O. A 90-minute linear gradient (Supplemental Table S14) and oxonium ion triggered EtHCD method was used (Supplemental Table S15 and 16).

Nipah G_M Head, N481D and T483A samples were analysed using an Ultimate 3000 RSLCnano system coupled to an Orbitrap Eclipse mass spectrometer (Thermo Fisher Scientific). Peptides were loaded onto a 75 µm × 2 cm pre-column and separated on a 75 µm × 15 cm Pepmap C18 analytical column (Thermo Fisher Scientific). Buffer A was 0.1% FA in H_2O and buffer B was 0.1% FA in 80% MeCN with 20% H_2O. A 60-minute linear gradient (Supplemental Table S17) and oxonium ion triggered EtHCD (Supplemental Table S18 and 19) method was used.

Glycoproteomics Data Analysis

Peak lists were generated using XCalibur™ (Thermo Fisher Scientific, version 4.0.27.19) and data were analysed in Byos (Protein Metrics, version 5.4.52). The database queried consisted of the FASTA sequence for the Nipah G protein sequence of interest, alongside Byonics 71 common contaminants and decoys. Digestion was set to RK, FYWML and TASV for trypsin, chymotrypsin and alpha-lytic protease digests respectively with maximum of two missed cleavages. Fixed modifications were carbamidomethylation (57.02 Da) and variable modifications were methionine oxidation (15.99 Da) and deamidation (N or Q) (0.98 Da). Byos common human N-linked (132 glycans, which covered the structures identified in our IM-MS/MS described below) and O-linked (6 glycans for N-linked analysis, 9 glycans for O-linked analysis) glycan libraries were used. FDR

was set to 0.01 and mass tolerances for precursor and fragment ions set to 6.00 ppm and 20.00 ppm respectively. For analysis, a minimum Byos threshold score of 100 was used for O-glycopeptide filtering and 300 for N-glycopeptide filtering, with the exception of peptides covering site N72 for which a score of 100 was used due to limited detection and fragmentation data arising from the hexahistidine tag, leading to lower scoring. Glycopeptides were manually validated, with O-glycan site localisation confirmed through c/z and intact b/y fragmentation products, as well as oxonium ion presence. For N-glycan quantification, one peptide sequence per glycosite was chosen as representative for each site (Supplemental Table S20) and the extracted ion chromatogram intensities for each glycopeptide were summed and plotted relative to the total intensity for each glycosite (Supplemental Data File 4). O-glycopeptide site identifications are listed in Supplemental Data File 5.

Differential Endoglycosidase Digestion with ^{18}O labelling

The method by Cao *et al*[34] was adapted with the following modifications; 5 µg of Nipah G_M ectodomain was denatured in 4M Urea, 1% SDS for 1hr at RT. Protein was reduced in 10 mM TCEP (pH 7.5) for 30 min RT and labelled with 50 mM iodoacetamide in 100 mM AmBic for 30 min in the dark, RT. Protein solution was added to 250 µg of 50:50 mix of Sera-Mag™ speed bead magnetic carboxylate modified particles (Cytiva). MeCN was added to a final concentration of ~90% and the solution was incubated for 30 min, 37°C, with shaking. A magnetic plate was used to immobilise the beads, allowing aspiration of the supernatant. Beads were washed sequentially with of 80% EtOH (x2) and 100% MeCN. 0.2 µg of trypsin/LysC (Promega) in 50 mM AmBic was added to the beads, before incubation overnight at 37°C. The subsequent peptide solution was aspirated and kept, before 5% DMSO H_2O was added for 2 min, and pooled with the peptides. Half of the samples were further treated with 0.2 µg of GluC (NEB) in 50 mM AmBic for 5.5 h at

37°C, after which all peptides were dried in a vacuum concentrator. Experiment was carried out in biological triplicate for both trypsin/LysC only and trypsin/LysC + GluC conditions.

Peptides were resuspended in 50 mM ammonium acetate (pH 5.5) (37°C with 10 min shaking) before addition of 313 NEB units of Endo-H (New England Bioscience) per sample. Samples were incubated for 1 h at 37°C before drying and sealing. PNGase F (New England Bioscience) was lyophilised and resuspended in 50 mM AmBic in $H_2{}^{18}O$ before adding to peptides, with a 160 NEB units per sample. Samples were incubated for 30 min 37°C before heat inactivation (10 min, 100°C) and drying in a vacuum concentrator. Labelled peptides were reconstituted in 2% MeCN 0.05% TFA and were run on an LC-MS/MS using the same method as for glycoproteomics.

Peak lists were generated by XCalibur™ (Thermo Fisher Scientific, version 4.0.27.19) and data were analysed in Proteome Discoverer (Thermo Fisher Scientific, version 3.0). Digestion was set to RK, and RKED for trypsin or trypsin/LysC + GluC protease digests respectively with maximum two missed cleavages. Static modifications were carbamidomethylation (57.02 Da) and dynamic modifications were methionine oxidation (15.99 Da) and ^{18}O deamidation (N only) (2.98 Da). Strict FDR was set to 0.01, and mass tolerances for precursor and fragment ions were set to 10 ppm and 0.6 Da respectively. The library searched consisted of the Nipah G_M sequence lacking a signal sequence, supplemented with the CRAPome[35]. The same peptide sequences were used for quantification, with the exception of N529 (Supplemental Table S20). This was due low intensities of the matching peptide (<3000) and subsequent missing values for multiple replicates (Supplemental Data File 6). A shorter, more abundant, peptide spanning N529 from the trypsin/LysC-GluC digestion was used instead. For quantification, the extracted ion chromatogram intensities for each peptide of that sequence were summed and plotted relative to the total intensity for each glycosite, listed in Supplemental Data File 6.

Biolayer Interferometry

Journal Pre-proof

Octet® streptavidin biosensors (Sartorius) were soaked in 200 µL PBS for 30 min prior to each run. Nipah G mutants and ephrin B2 were diluted in PBS with 0.01% Tween 20. Blocking buffer was PBS with 0.01% Tween 20. Runs were carried out in technical quintuplicate using an Octet® R8 system (Supplemental Table S21 for run settings). Concentrations of mutants were titrated in a 2-fold dilution series starting at 200, 100 or 44 nM. Timings and concentrations were determined prior such that the association phase would reach steady state, and the dilution series covered both above and below the determined K_D. Analysis was carried out in Octet® Analysis Software, where baselines were subtracted prior to fitting data to a 1:1 binding model. The R_{max} of each curve was fit to a steady state model to give the K_D values. Students unpaired, two-tailed t-test was used to calculate statistical significance for K_D values.

HDX-MS

Nipah G WT head, T483A and N481D were diluted to 5.9 µM, and deuterated PBS (pD 7.4) was added to a deuterium fraction of 88% to initiate the exchange reaction. At selected time intervals (15 s, 150 s, 1500 s, 15000 s, 24 h), a fraction of reaction mixture containing the equivalent of 23 pmol of protein sample was withdrawn and quenched with ice-cold buffer containing 45 mM phosphate,2 M urea and 40 mM TCEP, which dropped the pH/D to 2.3 and the deuterium fraction to 50%. Quenched samples were kept for 30 s on ice then flash frozen in liquid nitrogen and stored at -80°C before LC-MS analysis. For WT-ephrin B2 complexing, Nipah G WT head was incubated in a 1:1.2 molar ratio with ephrin B2 (28-165), or an equivalent volume of PBS. Deuterium labelling was carried out as specified above, and the equivalent of 46 pmol of protein samples were quenched. Maximally labelled samples (MaxD) were produced by labelling Nipah G WT head, T483A and N481D at 37°C in the presence of 4 M deuterated Urea in D_2O and 2.4 mM TCEP, keeping the deuterium fraction at 88%. The maximally labelled samples were quenched after 18 h with ice-cold buffer containing 45 mM phosphate, 1.4% FA, in a similar manner as for the other labelled samples and stored in the same fashion. Timepoints were

repeated and recorded in technical triplicate, with the exception of the following timepoints: WT (mutant analysis) 15 s, WT (ephrin B2 analysis) 1500 s, N481D 150 s, T483A 150 s, which were recorded in duplicate

For LC-MS, samples were quickly thawed and injected into an Acquity UPLC M-Class System with HDX Technology (Waters), on-line digested at 20°C into a hand packed pepsin column, prior to online deglycosylation with a PNGAse Rc column[36] (Affipro), which removed N-glycans through enzymatic deamidation of N-glycosylated asparagines into aspartates. Peptides were trapped/desalted with solvent A (0.23% formic acid in water) for 4 min at 120 µL/min and at 1°C through an Acquity BEH C18 VanGuard pre-column (1.7 µm, 2.1 mm × 5 mm, Waters). Peptides were eluted from the trap column into an Acquity UPLC BEH C18 analytical column (1.7 µm, 1 mm × 100 mm, Waters) with a 7 min-linear gradient raising from 8 to 35% of solvent B (0.23% formic acid in acetonitrile) at a flow rate of 40 µL/min and at 1°C (Supplemental Table S22 for LC instrument settings). Eluted peptides went through electrospray ionization in positive mode and underwent MS analysis onto a Synapt-G2 Si TWIMS-TOF (Waters) with ion mobility separation enabled (Supplemental Table S23 for IM-MS instrument settings). For peptide identification, non-deuterated protein samples were run with the same method as for deuterated samples, but underwent fragmentation using an MS^E method, applying collision energy ramping from 20 to 30 kV (Supplemental Table S24 for MS^E instrument settings)

Peak lists were generated by MassLynx™ (version 4.1, Waters) and MS^E runs were analysed with ProteinLynx Global Server (PLGS) (version 3.0, Waters) and peptides identified in 3 out of 4 replicates, with at least 0.2 fragments per amino acid (at least 2 fragments in total), with mass error below 10 ppm were selected in DynamX (version 3.0, Waters). Using both retention time and drift time, spectra of deuterated peptides were matched to the peptide map and manually annotated to calculate peptide deuterium incorporation. HDX-MS uptake plots and individual uptake values can be found in Supplemental Data Files 7 and 8 respectively.

Statistical Rationale

For the glycoproteomics experiments, FDR was set to >0.01. Students unpaired, two-tailed t-test was used to test for statistical significance between complex glycan containing glycopeptides and endo-H resistant HPLC glycan species. For testing statistical significance between complex glycan containing glycopeptides and ^{18}O labelled peptides at each glycosite, a one-way ANOVA with post-hoc Šidák multiple pairwise comparison correction was applied to control for type-I error inflation when conducting multiple pairwise comparisons between paired means. This test was also used for the comparison between ectodomain glycopeptides and head domain glycopeptides. Identical glycopeptides and ^{18}O labelled peptides were compared, except for N529, where the more abundant trypsin/LysC-GluC peptide was used for ^{18}O labelled peptides (see "Differential endoglycosidase treatment with ^{18}O labelling" methods section for more detail). A one-way ANOVA with post-hoc Dunnett's multiple comparison correction was used to test for statistical significance between the K_d of Nipah G_M Head and glycosite mutants, and between Nipah G_M head ± kifunensine. This test and correction were chosen to control for type-I error inflation when conducting pairwise comparisons between a control and multiple experimental means. Statistics and graphing for these analyses were conducted in GraphPad Prism v10.2.3. Statistical reporting for HPLC, proteomics and BLI can be found in Supplemental Data File 9.

The threshold for a statistically significant difference in HDX between two states was established based on an approach described earlier[37]. Briefly, a confidence interval (CI) was calculated based on the average standard deviation (SD) of peptide deuterium uptake values for time points performed in triplicates, according to equation 1:

Equation 1. $$SD_{state} = \frac{\sum {SD_i}^2}{N}$$

where N is the number of peptides considered, multiplied by the number of time points performed in triplicate. The CI at the significance level of 99% between two states was calculated based on

a zero-centered average difference in deuterium content, considering a two-tailed distribution with two degrees of freedom (n = 3) (equation 2)

Equation 2. $$CI = 9.925 \times \frac{\sqrt{SD_{stateA}+SD_{stateB}}}{\sqrt{n}}$$

The deuterium uptake of peptides carrying T483A were compared to the correspondent WT peptides by normalization with their respective MaxD uptake values (equation 3)[38]:

Equation 3. $$\Delta HDX = D_{max}\ WT\ peptide\left(\frac{Uptake\ T483A\ peptide}{Uptake\ T483A\ D_{max}\ peptide}\right) - Uptake\ WT\ peptide$$

HDX-MS reporting forms, including statistical outcomes, are reported in Supplemental Data Files 10 and 11.

Results

Phylogenetic analysis of Nipah G N-glycosylation

Most research on the G protein has used the primary sequence isolated from the 1998 outbreak[3], the so-called “Malaysia” strain. However, evolutionary analyses have shown Nipah has diverged to a second strain, the Bangladesh strain[39,40]. To identify potential differences in N-glycosylation sites between strains, we aligned 81 complete sequences from the NCBI virus database using MEGA11 (Supplemental Figs. S1 and S2), including viruses isolated from four different host species, across five countries. In agreement with other studies, we identified distinct groupings corresponding to the Malaysia and Bangladesh strains (clade I and II respectively in our analyses). These clades were further classified into a and b subclades, with IIa consisting of the perennial Bangladesh outbreaks, and IIb the more recent outbreaks in India. The location of sample collection shifted from Malaysia to Bangladesh and India over time (Supplemental Tables S1-S3), where most infections are now located. Notably, we found six of the seven predicted N-

glycan sites were conserved between all clades (**Fig. 1b**), however, a specific mutation (N481D) resulted in the loss of an N-glycan site. This was either introduced during the split from Hendra virus or occurred independently in both clade II and clade Ib isolates (**Fig. 1c**). This mutation event was also independent of host and country of origin, as there were no differences in host and origin within clades themselves (Supplemental Figs. S1 and 2).

Site-specific glycosylation of Nipah G

The role of the Nipah G N481 glycan is unknown but it could contribute to receptor binding affinity, infectivity and/or immune detection/evasion. Critically, there is no comprehensive site-specific glycan analysis of Nipah G, which is a first step for exploring glycan structure-function relationships, including N481. We used the soluble ectodomain of Nipah G_M (**Fig. 2**) for site-specific glycan characterisation, which was recombinantly expressed and purified from human embryonic kidney (HEK) Expi-293 cells to best mimic human glycosylation *in vitro*[41]. Purified SEC fractions of Nipah G_M ectodomain and head domain (Supplemental Fig. S3a and b) were measured by MP, showing that soluble G_M formed the expected stable tetramer (Supplemental Fig. S3d and e)[8]. Using BLI, we determined soluble G_M tetrameric ectodomain bound with picomolar affinity to recombinantly expressed soluble ephrin B2 binding domain (Supplemental Figs. S3c and S4). The G_M monomeric head domain interacted with ephrin B2 with nanomolar affinity in a 1:1 binding model (Supplemental Fig. S4b).

To identify all N-glycan structures present on G_M, we performed IM-MS/MS on three biological replicates. A previous study used MALDI-MS to investigate glycosylation, and identified a range of hybrid and complex type N-glycans, in addition to $Man_5GlcNAc_2$ for the monomeric G_M head domain[16]. However, our data showed a shift towards oligomannose structures with the tetrameric G_M ectodomain, with $Man_{5\text{-}8}GlcNAc_2$ ($Man_{5\text{-}8}$) present as major ion species (**Fig. 2a**). Three

biantennary complex type glycans were also detected as major ion species (*m/z* 1183, 1576 and 2077), as well as two bisected complex type glycans (*m/z* 1779 and 1941) (Supplemental Figs. S5 and 6). Additional complex-type N-glycans, specifically tri- and tetra-antennary species, were identified at low levels from ion-mobility extracted doubly and triply charged species (Supplemental Fig. S5). Oligomeric assembly has been implicated in influencing glycosylation by affecting protein conformation and glycosite accessibility for glycan trimming by mannosidases, and extension by glyosyltransferases[42-44]. Therefore, the abundance of oligomannose glycans is likely due to differences in glycan processing for the G_M tetrameric ectodomain vs monomeric head domain, as previously seen for HIV Env trimers and gp120 monomers[44].

HPLC of 2-AA labelled N-glycans confirmed the relative abundance of G_M structures by type (**Fig. 2b**). Treatment of N-glycans with Endo-H identifies oligomannose and hybrid-type glycans (i.e. Endo-H sensitive) from complex-type structures (i.e. Endo-H resistant). Overall, IM-MS/MS and HPLC data both indicated Endo-H sensitive glycans accounted for just under 50% of the total N-glycan pool (**Fig. 2c** and Supplemental Fig. S7), with ~70% present as Man_5 (32.6 ± 6.1%) Man_7 (23.8 ± 4.2%) and Man_8 (14.7 ± 4.1%) (Supplemental Data File 2). Conversely, multiple peaks were identified as Endo-H resistant glycans across the chromatogram, indicating complex-type N-glycans are structurally diverse, but individually present at lower levels compared to oligomannose structures.

We mapped the site-specific attachment and abundances of N-glycans on G_M from six biological replicates using LFQ glycoproteomics, (**Figs. 2d**). Four of the seven N-glycan sites were predominantly complex-type: N72 (72.3 ± 1.9%) and head sites N378 (84.9 ± 4.1%), N417 (80.7 ± 4.1%) and N529 (83.1 ± 6.3%). N378 and N529 glycopeptides were also predominantly fucosylated (82.3% and 82.9% respectively) and, along with N72, contained extensively sialyated N-glycans (36.6%, 43.0% and 35.1% respectively) (Supplemental Fig. S8), consistent with the complex N-glycans identified by IM-MS/MS (Supplemental Fig. S5). N417 was minimally

Journal Pre-proof

fucosylated (4.7 ± 1.6%) and had notably lower levels of sialyation (19.3 ± 5.6%). N306 had a mixture of complex (56.4 ± 7.6%), oligomannose (36.7 ± 9.6%) and hybrid-type (6.9 ± 2.6%) glycans. The N159 glycan, present in the neck of G_M, contained primarily oligomannose glycans (61.4 ± 7.4%), while the evolutionary divergent site, N481, was oligomannose rich (74.2 ± 6.6%) and was the only site with negligible amounts of complex N-glycans (<1%). Overall, results between the three methods were in high agreement, with no significant difference between HPLC and glycoproteomics quantifications (Supplemental Fig. S9). We also compared the relative abundance of glycopeptides between the Nipah G_M tetrameric ectodomain and the head domain. Sites N306, N378, N417 and N529 in the head domain displayed a significantly greater proportion of complex-type glycans than in the tetrameric ectodomain, ranging from an 11.4% increase in complex type glycans for N378 (ectodomain 80.1 % ± 5.6%, head domain 91.5 ±1.2%) to a 26.1% increase for N306 (ectodomain 56.4 ±7.6%, head domain 82.5 ± 1.8%). These data, along with our IM-MS/MS data, suggest that the glycan differences between the monomeric head domain and tetrameric ectodomain are a result of G tetramer oligomerisation (Supplemental Fig. S10).

To further confirm G_M N-glycosylation site occupancy as determined by LFQ glycoproteomics, we employed differential glycopeptide digestion, with sequential treatment of Endo-H, followed by PNGase-F digestion in the presence of $H_2{}^{18}O$, from three biological replicates (Supplemental Fig. S11)[34]. As PNGase-F requires a minimum of a chitobiose core, this method enables identification and quantification of glycopeptides through different mass shifts: *+203* (+GlcNAc) for glycopeptides containing Endo-H sensitive glycans whose core GlcNAc are not removed by PNGase-F, *+3* for glycopeptides with Endo-H resistant glycans that undergo enzymatic deamidation by PNGase-F in the presence of ^{18}O (Asn→Asp(^{18}O)), and *+0* for unoccupied N-glycan sites. We showed that our PNGase-F had negligible activity on the remaining GlcNAc following Endo-H digestion, as kifunensine treated G protein had >99% Endo-H sensitive peptides, which were not modified by further PNGase-F treatment (μ = 99.3 %, σ = 0.81). Due to

using different proteolytic digest methods (in-gel digestion vs SP3), we were unable to cover site N306 (see further description in the methods section). Five of the six sites were ~100% occupied, including N481 (μ = 100.0%, σ = 0.03), and only N529 was not fully occupied with 46.9% occupancy observed (σ = 1.8, n = 2) (**Figs. 2e** and **f**). To validate our glycoproteomics data, we compared the relative abundances observed for complex glycopeptides and Endo-H resistant glycopeptides using the LFQ and ^{18}O datasets respectively. We compared peptides with identical sequences, except for N529 (see Methods) (Supplemental Table S20) and samples from the same biological replicate. There were no significant differences between these two methods (Supplemental Fig. S11b), suggesting ionisation efficiency between glycopeptides did not affect quantification among N-glycan types.

Previous research into Hendra virus showed the stalk domain its G protein is highly populated with O-glycans that are important for viral fusion, and some of these sites are conserved with Nipah[30,31]. However, Nipah site-specific O-glycosylation has not been investigated. Using three biological replicates, we identified and localised eight O-glycan sites on Nipah G_M and two glycosite regions (Supplemental Data File 5, Supplemental Figure S12-13). O-glycans and their sites were assigned using oxonium ion triggered EtHCD, with sufficiency of c/z fragmentation affecting whether a site could be localised to a residue or a region. Seven sites were conserved between Nipah and Hendra (T103, T117, T119, S132, S134, T135, S137) including the two ST-rich regions S110-T119 and S132-S137 (Supplemental Fig. S12a). The first ST-rich region, S110-T119, located in the lower stalk region between the bottom two heads of Nipah (Supplemental Fig. S12b), contained both short and extended mono- and di-sialyated core-1 and core-2 type O-glycans. In particular, we could localise T117 displaying a single HexNAc, and T119 showing HexNAc, $HexNAc_1Hex_1$ or $HexNAc_2Hex_2NeuAc_1$ O-glycans. On the other hand, S132-S137, the other ST-rich region sitting on the upper stalk region, presented less extended mono- and di-sialyated core-2 type O glycans than S110-T119. We saw reduced extension amongst S132 and

Journal Pre-proof

S137, both displaying a single HexNAc and S132 a $HexNAc_1Hex_1$, while S134 and T135 showed evidence of more extended core-1 and core-2 structures of $HexNAc_2Hex_2NeuAc_1$ and $HexNAc_2Hex_2NeuAc_2$. Overall, the lesser global extent of sialylation and extension in S132-S137 suggests that it is less accessible for O-glycosyltransferase enzymes compared to S110-S119. This is expected as S110-T119 resides at the base of the tetrameric ectodomain, where the stalk helices are further apart from one another, while S132-S137 sit above the lower head regions on the stalk, where it is packed more tightly. The other conserved site, T103, which flanks S110-S119, was mainly populated with extended mono/di-sialyated core-1 and core-2 O-glycans, as was observed for Hendra[31].

The other O-glycosylation site, which was not present on Hendra G, was T149. Only a single HexNAc was present on T149, which is located in the neck domain. The model of the Nipah G_M ectodomain, based on the cryo-EM structure[8], has the side chains of T149 facing outward (Supplemental Fig. S12b), and therefore we would potentially expect further extension of these glycans. The lack of extended, larger structures at this site suggests this O-glycan is not accessible to glycosylation enzymes in the same way as the other sites. Our findings reveal that N-glycans are present near the ephrin B2 binding site while O-glycans are localised to the stalk. Based on known structures of Nipah G, they could have direct or allosteric implications for receptor binding.

Receptor binding of Nipah G_M glycosite mutants

Glycans attached to viral glycoproteins can influence receptor binding via direct interactions[45] and by altering protein structural conformations[46-48]. For Nipah G protein, the direct biophysical effects of glycan loss upon ephrin B2 binding have not been investigated, with previous studies focussing on phenotypic outcomes (viral fusion and cellular infectivity)[29]. To examine binding, we generated

individual N-glycosite mutants of G_M; T308A, S380A, S417A, T483A, T513A plus the evolutionary mutant N481D (Supplemental Fig. S14). Each N-glycan site was mutated via Ser/Thr to Ala in its respective sequon (N-X-S/T→N-X-A) to prevent potential O-glycosylation of Ser/Thr upon Asn mutation[49]. The binding affinity between ephrin B2 and the head domain of each mutant was measured by BLI. The head domain was chosen over the tetramer in order to study the interaction in a 1:1 binding model (**Fig. 3** and Supplemental Fig. S15). Furthermore, the picomolar affinity to the tetramer is at the concentration limit for BLI[50] (Supplemental Fig. S4). We did not detect any statistically significant differences in K_d values between G_M WT and T308A, S380A, S419A, and N481D mutants, indicating that these N-glycans do not directly contribute to ephrin B2 binding. However, a significant 4.7-fold decrease in binding affinity was observed with T531A (K_d = 95.5 ± 23.7 nM) compared to the WT (K_d = 20.7 ± 8.6 nM). The sidechain of T531 was previously suggested to interact through hydrophobic interactions with ephrin B2[9], therefore, a reduction in affinity for this mutant cannot be solely attributed to N-glycan (or O-glycan) loss. T483A also exhibited a significant 3.7-fold reduction in binding affinity (K_d = 76.8 ± 35.6 nM) for ephrin B2 compared to WT. Both the evolutionary mutant N481D and T483A result in N-glycan loss at N481, but T483A elicited 2.1-fold weaker binding than N481D (K_d = 36.7 ± 9.7 nM). Affinity for ephrin B2 did not change with kifunensine treated G_M, indicating that the presence of solely $GlcNAc_2Man_{8/9}$ glycoforms does not affect binding[51]. Furthermore, because the presence of an O-glycan at T483 was not observed for the WT and N481D, O glycosylation, or lack of, at this site did not affect binding (Supplemental File 5). Collectively these data suggest the reduction of ephrin B2 affinity observed for T483A was not due to loss of the associated N481 glycan but was directly related to the amino acid change from threonine to alanine (**Fig. 3**).

Structural dynamics of Nipah G_M and ephrin B2 complex

Journal Pre-proof

Loss of binding affinity for T483A implies a structural rearrangement of the binding site involving residues near the N481-X-T483 sequon. We used HDX MS to investigate structural dynamics associated with the N481D and T483A mutations compared to WT G_M head domains with and without ephrin B2. We followed the HDX of 139 peptides from 15 s to 24 h, covering 87% of G_M head domain sequence and 4 out of 5 glycosylation sites, including N481 (Supplemental Fig. S16).

A significant decrease in HDX was measured across overlapping peptides in multiple regions for WT G_M head upon ephrin B2 binding (**Fig. 4a** and Supplemental Fig. S17) mostly in agreement with the co-crystal structure[10] and previously published HDX-MS data for G_M ectodomain-ephrin B2 complex[9]. Peptide regions that displayed significant HDX changes were localised to the ephrin B2 binding site (Supplemental Fig. S17) and distributed across the oligomerisation interface, with amino acids Y231-L234, Q455-S459, E505-Y508 and L526-T538 displaying greatest protection from HDX in the bound G_M form. Both Y231-L234 and Q455-S459 do not form direct contacts with ephrin B2 (Supplemental Fig. S18), thus decrease in HDX is indicative of allosteric effects of rigidification in this region.

Compared to the WT, both N481D and T483A mutants exhibited a significant increase in HDX localised to the mutation sites (**Fig. 4a** and Supplemental Figs. S18b and c), in the region comprising residues T471-X481 (**Fig. 4b**). T483A also showed a clear increase in HDX across multiple time points in the longer peptide spanning T471-F496, which was only seen in a single time point for N481D (**Fig. 4c**). This indicates the effect of deprotection for the T483A mutant is not only localised to the mutation site but is transmitted to the adjacent segment N482-F496 (the region not covered by peptide T471-X481). Interestingly, the receptor-bound complex did not display a significant change in HDX for peptide T471-X481 (**Fig. 4d**), but did exhibit decreased HDX for T471-F496 (**Fig. 4e**). This shows protection for residues N482-F496 (**Fig. 4f**), which is the same region impacted by T483A mutant (and minimally by N481D mutant). Crucially, the

residues N482-F496 span a large loop in the ephrin B2 binding site (**Fig. 4f, left**). As the deuterium uptake of residues T471-F496 increases over time with the unbound WT (**Fig. 4e**), its slower deuterium exchange indicates the N482-F496 loop has a secondary structure. Ephrin B2 binding results in HDX protection at this loop, and the differences observed in HDX between the two mutants correlates to their different binding affinities for ephrin B2. The T483A mutation has a greater HDX compared to N481D, highlighting that this mutation has a greater impact on the dynamics of this loop – this is further reflected lower affinity for ephrin B2 observed for the T483A mutant compared to N481D, and the WT (**Fig. 3**). Of the other N-glycan sites covered, peptides spanning the two adjacent glycosites, N306 and N529 displayed lower HDX in the presence of ephrin B2. However, this is likely due to solvent occlusion from the glycan or protein backbone interactions with the receptor, as a loss of N306 through T308 doesn't change ephrin B2 affinity, and both regions are within the ephrin B2 binding site.

Discussion:

Here, we have identified a conserved region of vulnerability in Nipah G involving an N-glycan sequon, which substantially impacts receptor engagement. During Nipah virus evolution, the N481 residue in this sequon evolved to D481, but the following sixteen amino acids, including T483 (S483 in Hendra), are 100% conserved across all clades of Nipah and the primary strain of Hendra (Supplemental Fig. S2). These residues form a structured loop that becomes stabilised upon ephrin B2 binding, with our HDX data revealing stabilisation is localised within residues N482-F496. We have also shown that the naturally occurring N481D mutation has minimal impact on the conformational dynamics of this loop and on receptor binding, which in principle does not impact infectivity potential. However, mutating T483 to A483 significantly decreases receptor

Journal Pre-proof

binding affinity by negatively impacting dynamics of this loop. The crystal structure of the Nipah G_M-ephrin B2 complex (PDB: 2VSM) shows the T483 hydroxyl engaging in a hydrogen bond with the carbonyl of the N543 sidechain (**Fig. 5**) and mutation of 483 to alanine would disengage this bonding, leading to conformational rearrangement of the loop and subsequent decrease in ephrin B2 binding affinity. Taken together, our data suggest that the residues spanning 483-496 are a good candidate for therapeutic targeting, as many of the other sites displaying increased protection upon receptor engagement are buried within the oligomerisation interface, and therefore potentially much less accessible to therapeutic monoclonal antibodies or antivirals.

To our knowledge, this is the first site-specific glycan analysis of Nipah G_M. We identified that most sites are fully occupied with the exception of N529. However, we hypothesise that occupancy of N72 in a full-length G_M (i.e. on a Nipah virion) would likely be low due to the close predicted proximity of this residue to the viral membrane preventing entry to the OST complex active site[52,53]. We found N159 and N481 N-glycans were oligomannose rich, and oligomannose-rich sites have been recently linked to immune escape for enveloped viruses, due to their ability to sample a large conformational space and be recognised by innate immune C-type lectin receptors DC-SIGN, L-SIGN, dectin-2 and langerin[54,55]. The evolutionary variable N481 site had the greatest oligomannose occupancy, and so loss of this glycan could reduce immune pressure and clearance of Nipah. However, this does not correlate with the mortality rates seen for the two Nipah strains, where the mortality rate for the Bangladesh strain, which lacks this N-glycan, is considerably higher (~70-90%) than the Malaysia strain (~40%)[56]. Previous work suggests that loss of the 481 glycan through N481D increases fusogenicity in a pseudoviral model[29]. Therefore, an explanation for the non-conserved glycosite could be in response to increasing viral fusion, suggesting this region could be involved in F-protein engagement.

Acknowledgments

We thank Raymond Owens, Joseph Thrush, Kelly Parker and Leah McCaffrey for the NiVG 71-602 (full-length ectodomain), NiVG 183-602 (head domain), ephrin B2 28-165, BirA DNA constructs and HRV 3C protease. We also thank Shabaz Mohammed for useful discussion on SP3 digestion, ^{18}O labelling and proteomic analysis and Nikita Levin for useful discussion on O-glycan fragmentation. This work was supported by funding from the UK Research and Innovation Future Leaders Fellowship (MR/V02213*X*/1 to WBS) and the Wellcome Trust Sir Henry Dale Fellowship (218579/Z/19/Z to LW). The Rosalind Franklin Institute is funded by UK Research and Innovation through the Engineering and Physical Sciences Research Council (EPSRC).

CRediT author statement

Tia É. Hawkins: Conceptualisation, Methodology, Validation, Formal Analysis, Investigation, Data Curation, Writing – Original Draft, Writing – Review & Editing, Visualisation. Valeria Calvaresi: Methodology, Validation, Formal Analysis, Investigation, Data Curation, Writing – Review & Editing. Sean A. Burnap: Methodology, Validation, Investigation, Data Curation, Writing – Review & Editing. Yana Demyanenko: Methodology, Validation, Investigation. Liang Wu: Conceptualisation, Validation, Resources, Writing – Review & Editing, Supervision, Project Administration, Funding Acquisition. Weston B. Struwe: Conceptualisation, Validation, Resources, Writing – Review & Editing, Supervision, Project Administration, Funding Acquisition.

Data availability

Glycoproteomics, IM-MS and hydrogen deuterium exchange data are deposited to ProteomeXchange Consortium (http://proteomecentral.proteomexchange.org) via the PRIDE partner repository[57] with the dataset identifier PXD062381. HPLC and BLI raw data are deposited at Zenodo (doi:10.5281/zenodo.15569940). HPLC peak identifications (Supplemental Data File 2), IM-MS peak lists (Supplemental Data File 3) glycoproteomics peptide identifications (Supplemental Data 4-6) and statistic outputs for HPLC, glycoproteomics and BLI (Supplemental

Data File 9) are also attached as Supplementary files. In line with hydrogen deuterium mass spectrometry reporting guidelines[58], HDX-MS uptake plots, uptake values and reporting forms are attached as supplemental data files (Supplemental Data Files 7, 8, 10 and 11). Annotated MS/MS spectra for O-glycan site localisation is provided in Supplemental Data File 12. All other data available upon request.

Supplemental Data:

Supplemental Data File 1. Supporting figures and tables

Supplemental Data File 2. Released glycan HPLC identifications

Supplemental Data File 3. IM-MS peak lists

Supplemental Data File 4. LFQ N-glycoproteomics (glyco)peptide identifications

Supplemental Data File 5. O-glycopeptide identifications

Supplemental Data File 6. ^{18}O-labelled N-glycosite peptide identifications

Supplemental Data File 7. HDX-MS uptake plots

Supplemental Data File 8. HDX-MS individual deuterium uptake values

Supplemental Data File 9. Statistics Reporting for HPLC, proteomics and BLI experiments

Supplemental Data File 10. HDX-MS reporting form (mutants)

Supplemental Data File 11. HDX-MS reporting form (ephrin B2 complex)

Supplemental Data File 12. Annotated MS/MS Spectra for O-glycan site localisation

Competing interests:

WBS is a shareholder of Refeyn Ltd.

Journal Pre-proof

References:

(1) Chua, K. B.; Goh, K. J.; Wong, K. T.; Kamarulzaman, A.; Tan, P. S.; Ksiazek, T. G.; Zaki, S. R.; Paul, G.; Lam, S. K.; Tan, C. T. Fatal encephalitis due to Nipah virus among pig-farmers in Malaysia. *Lancet* **1999**, *354* (9186), 1257-1259. DOI: 10.1016/S0140-6736(99)04299-3

(2) Harcourt, B. H.; Tamin, A.; Ksiazek, T. G.; Rollin, P. E.; Anderson, L. J.; Bellini, W. J.; Rota, P. A. Molecular characterization of Nipah virus, a newly emergent paramyxovirus. *Virology* **2000**, *271* (2), 334-349. DOI: 10.1006/viro.2000.0340

(3) Chua, K. B.; Bellini, W. J.; Rota, P. A.; Harcourt, B. H.; Tamin, A.; Lam, S. K.; Ksiazek, T. G.; Rollin, P. E.; Zaki, S. R.; Shieh, W.; et al. Nipah virus: a recently emergent deadly paramyxovirus. *Science* **2000**, *288* (5470), 1432-1435. DOI: 10.1126/science.288.5470.1432

(4) Lo, M. K.; Lowe, L.; Hummel, K. B.; Sazzad, H. M. S.; Gurley, E. S.; Hossain, M. J.; Luby, S. P.; Miller, D. M.; Comer, J. A.; Rollin, P. E.; et al. Characterization of Nipah virus from outbreaks in Bangladesh, 2008-2010. *Emerg Infect Dis* **2012**, *18* (2), 248-255. DOI: 10.3201/eid1802.111492

(5) Nazmunnahar; Ahmed, I.; Roknuzzaman, A. S. M.; Islam, M. R. The recent Nipah virus outbreak in Bangladesh could be a threat for global public health: A brief report. *Health Science Reports* **2023**, *6* (7), e1423. DOI: 10.1002/hsr2.1423

(6) Hsu, V. P.; Hossain, M. J.; Parashar, U. D.; Ali, M. M.; Ksiazek, T. G.; Kuzmin, I.; Niezgoda, M.; Rupprecht, C.; Bresee, J.; Breiman, R. F. Nipah virus encephalitis reemergence, Bangladesh. *Emerg Infect Dis* **2004**, *10* (12), 2082-2087. DOI: 10.3201/eid1012.040701

(7) Sweileh, W. M. Global research trends of World Health Organization's top eight emerging pathogens. *Global Health* **2017**, *13* (1), 9. DOI: 10.1186/s12992-017-0233-9

(8) Wang, Z.; Amaya, M.; Addetia, A.; Dang, H. V.; Reggiano, G.; Yan, L.; Hickey, A. C.; DiMaio, F.; Broder, C. C.; Veesler, D. Architecture and antigenicity of the Nipah virus attachment

glycoprotein. *Science* **2022**, *375* (6587), 1373-1378. DOI: 10.1126/science.abm5561 From PubMed.

(9) Wong, J. J. W.; Young, T. A.; Zhang, J.; Liu, S.; Leser, G. P.; Komives, E. A.; Lamb, R. A.; Zhou, Z. H.; Salafsky, J.; Jardetzky, T. S. Monomeric ephrinB2 binding induces allosteric changes in Nipah virus G that precede its full activation. *Nat Commun* **2017**, *8* (1), 781. DOI: 10.1038/s41467-017-00863-3 From PubMed.

(10) Bowden, T. A.; Aricescu, A. R.; Gilbert, R. J. C.; Grimes, J. M.; Jones, E. Y.; Stuart, D. I. Structural basis of Nipah and Hendra virus attachment to their cell-surface receptor ephrin-B2. *Nat Struct Mol Biol* **2008**, *15* (6), 567-572. DOI: 10.1038/nsmb.1435 From PubMed.

(11) Bonaparte, M. I.; Dimitrov, A. S.; Bossart, K. N.; Crameri, G.; Mungall, B. A.; Bishop, K. A.; Choudhry, V.; Dimitrov, D. S.; Wang, L.-F.; Eaton, B. T.; et al. Ephrin-B2 ligand is a functional receptor for Hendra virus and Nipah virus. *Proceedings of the National Academy of Sciences* **2005**, *102* (30), 10652-10657. DOI: 10.1073/pnas.0504887102 (acccessed 2023/09/18/10:33:09). From pnas.org (Atypon).

(12) Xu, K.; Chan, Y.-P.; Bradel-Tretheway, B.; Akyol-Ataman, Z.; Zhu, Y.; Dutta, S.; Yan, L.; Feng, Y.; Wang, L.-F.; Skiniotis, G.; et al. Crystal Structure of the Pre-fusion Nipah Virus Fusion Glycoprotein Reveals a Novel Hexamer-of-Trimers Assembly. *PLOS Pathogens* **2015**, *11* (12), e1005322. DOI: 10.1371/journal.ppat.1005322 (acccessed 2023/09/18/09:58:57). From PLoS Journals.

(13) Aguilar, H. C.; Aspericueta, V.; Robinson, L. R.; Aanensen, K. E.; Lee, B. A Quantitative and Kinetic Fusion Protein-Triggering Assay Can Discern Distinct Steps in the Nipah Virus Membrane Fusion Cascade. *Journal of Virology* **2010**, *84* (16), 8033-8041. DOI: 10.1128/jvi.00469-10 (acccessed 2024/11/11/15:14:21). From journals.asm.org (Atypon).

(14) Liu, Q.; Stone, J. A.; Bradel-Tretheway, B.; Dabundo, J.; Benavides Montano, J. A.; Santos-Montanez, J.; Biering, S. B.; Nicola, A. V.; Iorio, R. M.; Lu, X.; et al. Unraveling a three-step

spatiotemporal mechanism of triggering of receptor-induced Nipah virus fusion and cell entry. *PLOS Pathogens* **2013**, *9* (11), e1003770. DOI: 10.1371/journal.ppat.1003770

(15) Murray, K.; Selleck, P.; Hooper, P.; Hyatt, A.; Gould, A.; Gleeson, L.; Westbury, H.; Hiley, L.; Selvey, L.; Rodwell, B. A morbillivirus that caused fatal disease in horses and humans - PubMed. *Science* **1995**, *268* (5207). DOI: 10.1126/science.7701348

(16) Bowden, T. A.; Crispin, M.; Harvey, D. J.; Aricescu, A. R.; Grimes, J. M.; Jones, E. Y.; Stuart, D. I. Crystal structure and carbohydrate analysis of Nipah virus attachment glycoprotein: a template for antiviral and vaccine design. *Journal of Virology* **2008**, *82* (23), 11628-11636. DOI: 10.1128/JVI.01344-08

(17) Wang, Q.; Liu, J.; Luo, Y.; Kliemke, V.; Matta, G. L.; Wang, J.; Liu, Q. The nanoscale organization of the Nipah virus fusion protein informs new membrane fusion mechanisms. *eLife* **2024**, *13*. DOI: 10.7554/eLife.97017.2

(18) Negrete, O. A.; Chu, D.; Aguilar, H. C.; Lee, B. Single Amino Acid Changes in the Nipah and Hendra Virus Attachment Glycoproteins Distinguish EphrinB2 from EphrinB3 Usage. *Journal of Virology* **2007**, *81* (19), 10804-10814. DOI: 10.1128/JVI.00999-07

(19) Fan, P.; Sun, M.; Zhang, X.; Zhang, H.; Liu, Y.; Yao, Y.; Li, M.; Fang, T.; Sun, B.; Chen, Z.; et al. A potent Henipavirus cross-neutralizing antibody reveals a dynamic fusion-triggering pattern of the G-tetramer. *Nat Commun* **2024**, *15* (1), 4330. DOI: 10.1038/s41467-024-48601-w

(20) Monath, T. P.; Nichols, R.; Feldmann, F.; Griffin, A.; Haddock, E.; Callison, J.; Meade-White, K.; Okumura, A.; Lovaglio, J.; Hanley, P. W.; et al. Immunological correlates of protection afforded by PHV02 live, attenuated recombinant vesicular stomatitis virus vector vaccine against Nipah virus disease. *Front Immunol* **2023**, *14*. DOI: 10.3389/fimmu.2023.1216225

(21) Loomis, R. J.; DiPiazza, A. T.; Falcone, S.; Ruckwardt, T. J.; Morabito, K. M.; Abiona, O. M.; Chang, L. A.; Caringal, R. T.; Presnyak, V.; Narayanan, E.; et al. Chimeric Fusion (F) and Attachment (G) Glycoprotein Antigen Delivery by mRNA as a Candidate Nipah Vaccine. *Front Immunol* **2021**, *12*. DOI: 10.3389/fimmu.2021.772864

Journal Pre-proof

(22) Doremalen, N. v.; Avanzato, V. A.; Goldin, K.; Feldmann, F.; Schulz, J. E.; Haddock, E.; Okumura, A.; Lovaglio, J.; Hanley, P. W.; Cordova, K.; et al. ChAdOx1 NiV vaccination protects against lethal Nipah Bangladesh virus infection in African green monkeys. *NPJ Vaccines* **2022**, *7* (1). DOI: 10.1038/s41541-022-00592-9

(23) Eldridge, J. H.; Egan, M. A.; Matassov, D.; Hamm, S.; Hermida, L.; Chen, T.; Tremblay, M.; Sciotto-Brown, S.; Xu, R.; Dimitrov, A.; et al. A Brighton Collaboration standardized template with key considerations for a benefit/risk assessment for a soluble glycoprotein vaccine to prevent disease caused by Nipah or Hendra viruses. *Vaccine* **2021**, *39* (38). DOI: 10.1016/j.vaccine.2021.07.030

(24) Rodrigue, V.; Gravagna, K.; Yao, J.; Nafade, V.; Basta, N. E. Current progress towards prevention of Nipah and Hendra disease in humans: A scoping review of vaccine and monoclonal antibody candidates being evaluated in clinical trials. *Tropical Medicine & International Health* **2024**, *29* (5). DOI: 10.1111/tmi.13979

(25) Playford, E. G.; Munro, T.; Mahler, S. M.; Elliott, S.; Gerometta, M.; Hoger, K. L.; Jones, M. L.; Griffin, P.; Lynch, K. D.; Carroll, H.; et al. Safety, tolerability, pharmacokinetics, and immunogenicity of a human monoclonal antibody targeting the G glycoprotein of henipaviruses in healthy adults: a first-in-human, randomised, controlled, phase 1 study - PubMed. *The Lancet. Infectious diseases* **2020**, *20* (4). DOI: 10.1016/S1473-3099(19)30634-6

(26) Alves, I.; Fernandes, Â.; Santos-Pereira, B.; Azevedo, C. M.; Pinho, S. S. Glycans as a key factor in self and nonself discrimination: impact on the breach of immune tolerance. *FEBS Lett* **2022**, *596* (12), 1485-1502. DOI: 10.1002/1873-3468.14347

(27) Miller, N. L.; Clark, T.; Raman, R.; Sasisekharan, R. Glycans in Virus-Host Interactions: A Structural Perspective. *Frontiers in Molecular Biosciences* **2021**, *8*, 666756. DOI: 10.3389/fmolb.2021.666756 (acccessed 2024/11/29/13:13:50). From PubMed Central.

Journal Pre-proof

(28) Kobayashi, Y.; Suzuki, Y. Evidence for N-Glycan Shielding of Antigenic Sites during Evolution of Human Influenza A Virus Hemagglutinin. *Journal of Virology* **2012**, *86* (7), 3446-3451. DOI: 10.1128/jvi.06147-11

(29) Biering, S. B.; Huang, A.; Vu, A. T.; Robinson, L. R.; Bradel-Tretheway, B.; Choi, E.; Lee, B.; Aguilar, H. C. N-Glycans on the Nipah Virus Attachment Glycoprotein Modulate Fusion and Viral Entry as They Protect against Antibody Neutralization. *Journal of Virology* **2012**, *86* (22), 11991. DOI: 10.1128/JVI.01304-12

(30) Stone, J. A.; Nicola, A. V.; Baum, L. G.; Aguilar, H. C. Multiple Novel Functions of Henipavirus O-glycans: The First O-glycan Functions Identified in the Paramyxovirus Family. *PLOS Pathogens* **2016**, *12* (2). DOI: 10.1371/journal.ppat.1005445

(31) Colgrave, M. L.; Snelling, H. J.; Shiell, B. J.; Feng, Y-R.; Chan, Y-P.; Bossart, K. N.; Xu, K.; Nikolov, D. B.; Broder, C. C.; Michalski, W. P. Site occupancy and glycan compositional analysis of two soluble recombinant forms of the attachment glycoprotein of Hendra virus. *Glycobiology* **2011**, 22 (4). DOI: 10.1093/glycob/cwr180

(32) Jones, D. T.; Taylor, W. R.; Thornton, J. M. The rapid generation of mutation data matrices from protein sequences. *Comput Appl Biosci* **1992**, *8* (3), 275-282. DOI: 10.1093/bioinformatics/8.3.275

(33) Tamura, K.; Stecher, G.; Kumar, S. MEGA11: Molecular Evolutionary Genetics Analysis Version 11. *Mol Biol Evol* **2021**, *38* (7), 3022-3027. DOI: 10.1093/molbev/msab120

(34) Cao, L.; Diedrich, J. K.; Ma, Y.; Wang, N.; Pauthner, M.; Park, S.-K. R.; Delahunty, C. M.; McLellan, J. S.; Burton, D. R.; Yates, J. R.; et al. Global site-specific analysis of glycoprotein N-glycan processing. *Nat Protoc* **2018**, *13* (6), 1196-1212. DOI: 10.1038/nprot.2018.024

(35) Mellacheruvu, D.; Wright, Z.; Couzens, A. L.; Lambert, J.-P.; St-Denis, N. A.; Li, T.; Miteva, Y. V.; Hauri, S.; Sardiu, M. E.; Low, T. Y.; et al. The CRAPome: a contaminant repository for affinity purification–mass spectrometry data. *Nature Methods 2013 10:8* **2013**, *10* (8). DOI: 10.1038/nmeth.2557

Journal Pre-proof

(36) Lambert, T.; Gramlich, M.; Stutzke, L.; Smith, L.; Deng, D.; Kaiser, P. D.; Rothbauer, U.; Benesch, J. L. P.; Wagner, C.; Koenig, M.; et al. Development of a PNGase Rc Column for Online Deglycosylation of Complex Glycoproteins during HDX-MS. *J Am Soc Mass Spectrom* **2023**, *34* (11). DOI: 10.1021/jasms.3c00268

(37) Houde, D.; Berkowitz, S. A.; Engen, J. R. The Utility of Hydrogen/Deuterium Exchange Mass Spectrometry in Biopharmaceutical Comparability Studies. *Journal of pharmaceutical sciences* **2010**, *100* (6). DOI: 10.1002/jps.22432

(38) Calvaresi, V.; Wrobel, A. G.; Toporowska, J.; Hammerschmid, D.; Doores, K. J.; Bradshaw, R. T.; Parsons, R. B.; Benton, D. J.; Roustan, C.; Reading, E.; et al. Structural dynamics in the evolution of SARS-CoV-2 spike glycoprotein. *Nature Communications 2023 14:1* **2023**, *14* (1). DOI: 10.1038/s41467-023-36745-0

(39) de Campos, G. M.; Cella, E.; Kashima, S.; Alcântara, L. C. J.; Sampaio, S. C.; Elias, M. C.; Giovanetti, M.; Slavov, S. N. Updated Insights into the Phylogenetics, Phylodynamics, and Genetic Diversity of Nipah Virus (NiV). *Viruses* **2024**, *16* (2), 171. DOI: 10.3390/v16020171

(40) Shi, J.; Sun, J.; Hu, N.; Hu, Y. Phylogenetic and genetic analyses of the emerging Nipah virus from bats to humans. *Infect Genet Evol* **2020**, *85*, 104442. DOI: 10.1016/j.meegid.2020.104442

(41) Goh, J. B.; Ng, S. K. Impact of host cell line choice on glycan profile. *Critical Reviews in Biotechnology* **2018**, *38* (6). DOI: 10.1080/07388551.2017.1416577

(42) Rudd, P. M.; Dwek, R. A. Glycosylation: Heterogeneity and the 3D Structure of Proteins. *Critical Reviews in Biochemistry and Molecular Biology* **1997**, *32* (1). DOI: 10.3109/10409239709085144

(43) Fercher, C.; Zacchi, L. F. Resolving the TorsinA Oligomerization Conundrum: The Glycan Hypothesis. *Frontiers in Molecular Biosciences* **2020**, *7*. DOI: 10.3389/fmolb.2020.585643

(44) Struwe, W. B.; Chertova, E.; Allen, J. D.; Seabright, G. E.; Watanabe, Y.; Harvey, D. J.; Medina-Ramirez, M.; Roser, J. D.; Smith, R.; Westcott, D.; et al. Site-Specific Glycosylation of

Virion-Derived HIV-1 Env Is Mimicked by a Soluble Trimeric Immunogen. *Cell Rep* **2018**, *24* (8), 1958-1966.e1955. DOI: 10.1016/j.celrep.2018.07.080

(45) St-Pierre, C.; Manya, H.; Ouellet, M.; Clark, G. F.; Endo, T.; Tremblay, M. J.; Sato, S. Host-Soluble Galectin-1 Promotes HIV-1 Replication through a Direct Interaction with Glycans of Viral gp120 and Host CD4. *Journal of Virology* **2011**, *85* (22). DOI: 10.1128/jvi.05351-11

(46) Casalino, L.; Gaieb, Z.; Goldsmith, J. A.; Hjorth, C. K.; Dommer, A. C.; Harbison, A. M.; Fogarty, C. A.; Barros, E. P.; Taylor, B. C.; McLellan, J. S.; et al. Beyond Shielding: The Roles of Glycans in the SARS-CoV-2 Spike Protein. *ACS Cent Sci* **2020**, *6* (10), 1722-1734. DOI: 10.1021/acscentsci.0c01056

(47) Li, Y.; Luo, L.; Rasool, N.; Kang, C. Y. Glycosylation is necessary for the correct folding of human immunodeficiency virus gp120 in CD4 binding. *Journal of Virology* **1993**, *67* (1). DOI: 10.1128/jvi.67.1.584-588.1993

(48) Mathys, L.; François, K. O.; Quandte, M.; Braakman, I.; Balzarini, J. Deletion of the Highly Conserved N-Glycan at Asn260 of HIV-1 gp120 Affects Folding and Lysosomal Degradation of gp120, and Results in Loss of Viral Infectivity. *PLoS ONE* **2014**, *9* (6). DOI: 10.1371/journal.pone.0101181

(49) Trinidad, J. C.; Schoepfer, R.; Burlingame, A. L.; Medzihradszky, K. F. N- and O-Glycosylation in the Murine Synaptosome. *Molecular & Cellular Proteomics* **2013**, *12* (12). DOI: 10.1074/mcp.M113.030007

(50) Kamat, V.; Rafique, A. Designing binding kinetic assay on the bio-layer interferometry (BLI) biosensor to characterize antibody-antigen interactions. *Analytical Biochemistry* **2017**, *536*. DOI: 10.1016/j.ab.2017.08.002

(51) Elbein, A. D.; Tropea, J. E.; Mitchell, M.; Kaushal, G. P. Kifunensine, a potent inhibitor of the glycoprotein processing mannosidase I. *The Journal of Biological Chemistry* **1990**, *265* (26), 15599-15605

Journal Pre-proof

(52) Alcorlo, M.; Dik, D. A.; De Benedetti, S.; Mahasenan, K. V.; Lee, M.; Domínguez-Gil, T.; Hesek, D.; Lastochkin, E.; López, D.; Boggess, B.; et al. Structural basis of denuded glycan recognition by SPOR domains in bacterial cell division. *Nat Commun* **2019**, *10* (1), 5567. DOI: 10.1038/s41467-019-13354-4

(53) Bañó-Polo, M.; Baldin, F.; Tamborero, S.; Marti-Renom, M. A.; Mingarro, I. N-glycosylation efficiency is determined by the distance to the C-terminus and the amino acid preceding an Asn-Ser-Thr sequon. *Protein Science : A Publication of the Protein Society* **2010**, *20* (1). DOI: 10.1002/pro.551

(54) Gao, C.; Stavenhagen, K.; Eckmair, B.; McKitrick, T. R.; Mehta, A. Y.; Matsumoto, Y.; McQuillan, A. M.; Hanes, M. S.; Eris, D.; Baker, K. J.; et al. Differential recognition of oligomannose isomers by glycan-binding proteins involved in innate and adaptive immunity. *Science Advances* **2021**, *7* (24). DOI: 10.1126/sciadv.abf6834

(55) Fogarty, C. A.; Fadda, E. Oligomannose N-Glycans 3D Architecture and Its Response to the FcγRIIIa Structural Landscape. *The Journal of Physical Chemistry B* **2021**, *125* (10). DOI: 10.1021/acs.jpcb.1c00304

(56) Tan, F. H.; Sukri, A.; Idris, N.; Ong, K. C.; Schee, J. P.; Tan, C. T.; Tan, S. H.; Wong, K. T.; Wong, L. P.; Tee, K. K.; et al. A systematic review on Nipah virus: global molecular epidemiology and medical countermeasures development - PubMed. *Virus evolution* **2024**, *10* (1). DOI: 10.1093/ve/veae048

(57) Perez-Riverol, Y.; Bandla, C.; Kundu, Deepti J.; Kamatchinathan, S.; Bai, J.; Hewapathirana, S.; John, Nithu S.; Prakash, A.; Walzer, M.; Wang, S.; et al. The PRIDE database at 20 years: 2025 update. *Nucleic Acids Res* **2025**, *53* (D1). DOI: 10.1093/nar/gkae1011

(58) Masson, G. R.; Burke, J. E.; Ahn, N. G.; Anand, G. S.; Borchers, C.; Brier, S.; Bou-Assaf, G. M.; Engen, J. R.; Englander, S. W.; Faber, J.; et al. Recommendations for performing, interpreting and reporting hydrogen deuterium exchange mass spectrometry (HDX-MS) experiments. *Nature Methods 2019 16:7* **2019**, *16* (7). DOI: 10.1038/s41592-019-0459-y

Journal Pre-proof

Legends:

Figure 1. Nipah virus evolved a divergent N-glycosylation site. a) Schematic representation of Nipah virus binding to host cells through ephrinB2/B3 interactions with the G protein, leading to fusion through F protein rearrangement. **b**) Overview of Nipah G_M domain structure and predicted N-glycans sites. **c**) Distribution of clades across location of isolates (left). Most prevalent clade shown for each country, identified through phylogenetic analysis (Supplemental Fig. S1). clade Ia: yellow, clade Ib: green, clade IIa: purple, clade IIb blue. Sequence alignment of clades and Hendra outgroup for residues 479-489 (right). Red amino acids denote a loss of N-glycosylation.

Figure 2. Nipah G_M ectodomain N-glycosylation. a) Total ion spectrum of released N-glycans by IM-MS scaled to the most abundant ion, Man_5-$GlcNAc_2$ (*m/z* 1347). **b**) HPLC of 2-AA labelled N-glycans with peaks labelled as endo-H sensitive (i.e. oligomannose and hybrid) in green and endo-H resistant (i.e. complex) in purple. **c**) Mean abundance of glycan subtype (coloured as above). **d**) Glycan subtype specific for each site determined by label-free quantitative (LFQ) glycoproteomics, (n=6 and n=5 for N72). Glycans types are labelled green for oligomannose, yellow for hybrid and purple for complex. **e**) Occupancy of N-glycan sites measured by ^{18}O labelled proteomics (n=3, except N592 where n=2). **f**) Location of glycans on Nipah G tetrameric ectodomain (PDB: 7TXZ & 7TY0) modelled using GLYCAM (the most abundant glycan species per site is shown).

Journal Pre-proof

Figure 3. Effects of Nipah G_M glycan knockout mutants on ephrin B2 binding. Location of glycans on Nipah G_M in relation to the ephrin B2 binding interface (top). N-glycan chitobiose or singular GlcNAcs are shown in blue. G_M and ephrin B2 are yellow and green respectively (Image produced in PyMOL using PDB: 2VSM, 7TY0). Box and whisker plots of steady state K_d of G_M mutants binding to ephrin B2 (bottom). Boxes represent the 25th-75th percentile, with whiskers extending from the minimum to maximum datapoints. Annotated values are mean K_D values ± 1 standard deviation.

Figure 4. HDX-MS of Nipah G_M mutants +/- ephrin B2. a) Differential HDX plot of Nipah G_M head compared to N481D, T483A, and plus ephrin B2 (+EB2). Alignment of the Nipah G_M structure to the HDX plot (top), with the beta propellers (BP) highlighted. b) Normalised uptake plot for peptide T471-X481. c) Normalised uptake plot for peptide T471-F496. d) Normalised uptake plot for peptide T471-X481 for incubation with ephrin B2. e) Normalised uptake plot for peptide T471-F496 for incubation with ephrin B2. f) Magnitude of HDX change across the Nipah G_M structure upon ephrin B2 binding is superimposed to PDB: 2VSM & 7TY0; peptides T471-X481 (light purple) and T471-F496 (indigo) are highlighted. Ephrin B2 in green, uncovered regions in grey.


Figure 5. Hydrogen bonding network around N481. N482-F496 is shown in light pink, ephrin B2 is green, T483 is dark magenta. Image produced from PDB: 2VSM (distances are in Å).

Journal Pre-proof


Journal Pre-proof

Evolutionary Glycan Site Stabilizes Nipah G Engagement


Journal Pre-proof




Journal Pre-proof




Journal Pre-proof







Journal Pre-proof



Journal Pre-proof



Journal Pre-proof



Journal Pre-proof

Journal Pre-proof

Highlights

- First site-specific glycan characterisation of the Nipah virus glycoprotein (G) tetramer
- Phylogenetic analysis reveals the N481 N-glycan site is not evolutionary conserved
- The absence of the N481 glycan does not contribute to ephrin B2 receptor binding
- However, the Ser/Thr in the sequon is conserved and maintains Nipah G stability

Journal Pre-proof

Journal Pre-proof

In Brief Statement:

Nipah virus is a deadly pathogen and the roles of glycosylation in its infection are poorly understood. Here, we investigated the evolution of the Nipah virus attachment glycoprotein (G) and characterised its site-specific glycosylation. We determined key allosteric residues in a non-conserved N-glycan site that are critical for receptor binding. Our data provides structural and dynamic evidence for Nipah glycoprotein stabilisation within the N481 sequon, offering insights for therapeutic development.

Journal Pre-proof

Journal Pre-proof

Declaration of Interest Statement

☐ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☐ The author is an Editorial Board Member/Editor-in-Chief/Associate Editor/Guest Editor for this journal and was not involved in the editorial review or the decision to publish this article.

☒ The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Weston Struwe is a shareholder of Refeyn Ltd who manufacture mass photometry instrumentation used in this study.

Journal Pre-proof