Galectin-3 decreases 4-1BBL bioactivity by crosslinking soluble and membrane expressed 4-1BB

Search within citing articles

CD137 (4-1BB) requires physically associated cIAPs for signal transduction and antitumor effects

J Glez-Vaz, A Azpilikueta, MC Ochoa, I Olivera… - Science …, 2023 - science.org

CD137 (4-1BB) is a member of the TNFR family that mediates potent T cell costimulatory signals upon ligation by CD137L or agonist monoclonal antibodies (mAbs). CD137 agonists …

☆ Save 99 Cite Cited by 17 Related articles All 9 versions

The role of galectin-3 in retinal degeneration and other ocular diseases: a potential novel biomarker and therapeutic target

Z Zhou, Z Feng, X Sun, Y Wang, G Dou - International Journal of …, 2023 - mdpi.com

Galectin-3 is the most studied member of the Galectin family, with a large range of mediation in biological activities such as cell growth, proliferation, apoptosis, differentiation, cell …

☆ Save 99 Cite Cited by 12 Related articles All 8 versions »

Large soluble CD18 complexes with exclusive ICAM-1-binding properties are shed during immune cell migration in inflammation

A Ferapontov, A Mellemkjær, HM McGettrick… - Journal of translational …, 2025 - Elsevier

The family of heterodimeric CD11/CD18 integrins facilitate leukocyte adhesion and migration in a wide range of normal physiologic responses, as well as in the pathology of …

☆ Save 99 Cite Cited by 5 Related articles All 8 versions

Protein–protein interactions in immune disorders and inflammation

KM Poluri, K Gulati, DK Tripathi, N Nagar - Protein-Protein Interactions …, 2023 - Springer

Immune system disorders arise when the immune system is either over-activated or depressed. Autoimmune disorders that affect millions of people worldwide, set in once the …

☆ Save 99 Cite Cited by 8 Related articles

Polyarticular juvenile idiopathic arthritis has a distinct co-inhibitor receptor profile

E Sag, Z Balik, S Demir, U Akca Kaya, S Sener… - …, 2025 - academic.oup.com

Objectives JIA is the most common rheumatic disease of childhood; the pathogenesis is associated with T-cell activation. T-cell activation can be counterbalanced by signals …

☆ Save 99 Cite Cited by 1 Related articles All 9 versions

CD137 (4-1BB) requires physically associated cIAPs for signal transduction and antitumor effects

J González-Vaz, A Azpilikueta, MC Ochoa, I Olivera… - 2023 - digital.csic.es

CD137 (4-1BB) is a member of the TNFR family that mediates potent T cell costimulatory signals upon ligation by CD137L or agonist monoclonal antibodies (mAbs). CD137 agonists …

☆ Save 99 Cite Related articles »

Impacto da inibição da galectina-3 na reversão da esteato-hepatite não alcoólica em modelo experimental

ARM Santos, JPG Camporez - 2025 - repositorio.usp.br

Padrões de vida sedentários, associados ao consumo excessivo de dietas com alto teor lipídico e calórico, estão diretamente relacionados ao desenvolvimento de obesidade …

☆ Save 99 Cite Related articles All 2 versions »