N-glycosylation of mannose receptor (CD206) regulates glycan binding by C-type lectin domains

Search within citing articles

Microbial Rhamnolipid-Stabilized mRNA Nanovaccines Enhance Adaptive Immunity via Dendritic Cell Targeting

W Kong, Z Dong, Y Wei, W Liu, S Li… - Advanced Functional …, 2026 - Wiley Online Library

ABSTRACT Poly (ethylene glycol)(PEG)-lipids endow lipid nanoparticles (LNPs) with colloidal stability but elicit anti-PEG immunity upon repeat dosing. Here it is replaced PEG …

Save Cite Cited by 2 Related articles All 2 versions

The role of *ladderlectin* in spermatogenesis and ovarian sperm storage in the black rockfish (*Sebastes schlegelii*)

R Yang, N Wang, W Song, F Zhang, X Gao… - Communications …, 2025 - nature.com

Ladderlectin, a teleost-specific C-type lectin, has been primarily associated with innate immune defense. However, this study unveils an important role of ladderlectin in the …

Save Cite Cited by 2 Related articles All 5 versions

Spatiotemporal targeting of miR-125b-5p regulatory hub via dual-ligand nanocarriers to polarize tumor-associated macrophages towards tumoricidal phenotype in …

G Sharma, AR Sharma, V Dewaker, S Kim… - Chemical Engineering …, 2026 - Elsevier

Reprogramming tumor-associated macrophages (TAMs) from an immunosuppressive M2 state toward a tumoricidal M1 phenotype is a promising strategy for cancer immunotherapy …

Save Cite Related articles All 2 versions

A Mannose Receptor from *Litopenaeus vannamei* Involved in Innate Immunity by Pathogen Recognition and Inflammation Regulation

N Guo, Y Liu, Q Hao, M Sun, F Li - International journal of molecular …, 2023 - mdpi.com

Mannose receptor, as a member of the C-type lectin superfamily, is a non-canonical pattern recognition receptor that can internalize pathogen-associated ligands and activate …

Save Cite Cited by 10 Related articles All 9 versions

Deglycosylation induces a novel distal conformation in the mannose receptor CD206

G Alvarez, S Di Lella, M Pickholz, P Scodeller… - Scientific reports, 2026 - nature.com

Deglycosylation can disrupt protein structure and dynamics in ways that compromise function. In this study, we examine the structural and functional consequences of N-glycan …

Save Cite Cited by 2 Related articles All 7 versions

Nanotherapies for Atherosclerosis: Targeting, Catalysis, and Energy Transduction

Y Yang, J Qu, T Xiao, X Han… - Advanced healthcare …, 2026 - Wiley Online Library

Atherosclerosis (AS), the pathological cornerstone of global cardiovascular mortality, remains inadequately managed by current therapies due to low bioavailability, off-target …

Save Cite Cited by 4 Related articles All 6 versions

A macrophage mannose receptor 1 with dual CTLDs played a crucial role in defense against Ecytonucleospora hepatopenaei in Litopenaeus vannamei

L Zhang, Y Ma, Z Li, Q Zeng, M Wang - Fish & Shellfish Immunology, 2026 - Elsevier

Mannose receptor (MR), as a crucial member of the C-type lectins (CTLs), plays an important role in the innate immune responses of invertebrates. In Litopenaeus vannamei …

Save Cite Related articles All 4 versions

Mannose receptor contributes to preferential oligomannosidic glycoform clearance in a murine pharmacokinetics model

D Falck, S Gijze, AM Wasynczuk, S Malik, A Tridon… - Mabs, 2026 - Taylor & Francis

The pharmacokinetic profile of therapeutic antibodies, which affects the dosing frequency, efficacy, and cost of the products, is influenced by several molecular features, including N …

Save Cite Related articles All 4 versions

Associations of polymorphisms in the myeloid immunoregulatory receptor gene MRC1 with expression and prognosis in acute myeloid leukemia

M Li, F Yin, Z Shi, T Sun, C Ji, M Li - Frontiers in Immunology, 2026 - frontiersin.org

Background Acute myeloid leukemia (AML) shows substantial biological and clinical heterogeneity that extends beyond leukemia-intrinsic genetic alterations. MRC1, which …

Save Cite Related articles All 5 versions

Pinellia ternata lectin induces mitochondrial damage and inflammation in RAW264. 7 cells via mannose receptor-mediated endocytosis

Y Huang, J Li, J Cao, X Wang, H Wu, M Shen… - Toxicology and Applied …, 2026 - Elsevier

A lectin from Pinellia ternata tubers (PTL, 12 kDa) with potent pro-inflammatory activity binds to the plasma membrane and translocates into the cytoplasm. However, its cellular entry …

Save Cite Related articles All 2 versions