Recognition of highly branched N-glycans of the porcine whipworm by the immune system

Search within citing articles

Insights into glycobiology and the protein-glycan interactome using glycan microarray technologies

J Heimburg-Molinaro, AY Mehta, CA Tilton… - Molecular & Cellular …, 2024 - mcponline.org

Glycans linked to proteins and lipids and also occurring in free forms have many functions, and these are partly elicited through specific interactions with glycan-binding proteins …

☆ Save 99 Cite Cited by 33 Related articles All 6 versions

Distribution, biosynthesis, recognition, identification and biological functions of LacdiNAc structures on N-and O-glycans: A review

Y Zhang, L Wu, X Liu, X Chen, J Li - International Journal of Biological …, 2025 - Elsevier

Abstract LacdiNAc (GalNAcβ1→ 4GlcNAc), a disaccharide structure on N-and O-glycans of glycoproteins, represents an important form of glycosylation modification. Several reviews …

☆ Save 99 Cite Cited by 2 Related articles All 3 versions

Glycoproteomic and single-protein glycomic analyses reveal zwitterionic N-glycans on natural and recombinant proteins derived from insect cells

S Yan, J Vanbeselaere, C Ives, D Stenitzer… - Molecular & Cellular …, 2025 - mcponline.org

Insect cells are a convenient cell factory to produce recombinant glycoproteins. Their glycosylation potential is believed to be simple, needing primarily addition of …

☆ Save 99 Cite Cited by 10 Related articles All 12 versions

Diversity-Oriented Enzymatic Modular Synthesis of *N*-Glycans Bearing LacdiNAc-Related Epitopes

X Yin, X Zhang, K Zhong, H Wang, Z Hu, CC Liu… - Organic …, 2025 - ACS Publications

The disaccharide unit GalNAcβ1, 4GlcNAc (LacdiNAc or LDN) and its α1, 3-fucosylated derivative LDNF and α2, 6-sialylated derivative 6′-SLDN are common epitopes of various …

☆ Save 99 Cite Cited by 3 Related articles All 4 versions

Evolutionary analyses of the animal glycosyltransferase family 54 reveals two β1, 4-N-acetylglucosaminyltransferase families

A Morigo, RE Teppa, M Nagae, H Yagi, Y Kizuka… - Iscience, 2025 - cell.com

N-acetylglucosaminyltransferases involved in branched N-glycans synthesis, a major post-translational modification, are gathered in the CAZy glycosyltransferase family 54. To date …

☆ Save 99 Cite Cited by 3 Related articles All 11 versions

Dual roles of *Anisakis pegreffii* proteins in macrophage immune dynamics

M Zeng, S Alsobaie, X Wang, S Li… - Frontiers in …, 2025 - frontiersin.org

Background Anisakis infections have become a significant public health concern primarily caused by consuming raw or undercooked seafood. This in is due to a shift in global eating …

☆ Save 99 Cite Cited by 3 Related articles All 9 versions »

N-glycan core tri-fucosylation requires Golgi α-mannosidase III activity that impacts nematode growth and behavior

J Kendler, F Wöls, S Thapliyal, E Arcalis… - Journal of Biological …, 2024 - jbc.org

N-glycans with complex core chitobiose modifications are observed in various free-living and parasitic nematodes but are absent in mammals. Using Caenorhabditis elegans as a …

☆ Save 99 Cite Cited by 6 Related articles All 13 versions

Evolution and diversity of glycomolecules from unicellular organisms to humans

RD Cummings - BioCosmos, 2024 - reference-global.com

The emergence of glycomolecules in nature coincided with the dawning of the ribonucleic acid (RNA) and deoxyribonucleic acid (DNA) world, along with the development of a …

☆ Save 99 Cite Cited by 10 Related articles All 3 versions »

Glycans of parasitic nematodes–from glycomes to novel diagnostic tools and vaccines

S Yan - Carbohydrate Research, 2025 - Elsevier

Nematodes, commonly known as roundworms, are among the most prevalent and diverse multicellular organisms on Earth, belonging to the large phylum Nematoda. In addition to …

☆ Save 99 Cite Cited by 7 Related articles All 7 versions

Integrative Quantitative Proteomics and N-Glycoproteomics Reveal Age-Processing of N-Glycosylation in *Caenorhabditis elegans*

X Hu, W Xiang, S Qin, X Zhang, Z Tian - Proteomics, 2026 - Wiley Online Library

ABSTRACT N-glycosylation, a highly diverse post-translational modification, critically regulates organismal aging. The Caenorhabditis elegans (C. elegans) model is one of the …

☆ Save 99 Cite Cited by 1 Related articles All 3 versions