Sulfated pentagalloylglucoside is a potent, allosteric, and selective inhibitor of factor XIa

Search within citing articles

Studies on fragment-based design of allosteric inhibitors of human factor XIa

RS Boothello, NV Sankaranarayanan… - Bioorganic & medicinal …, 2020 - Elsevier

Human factor XIa (hFXIa) has emerged as an attractive target for development of new anticoagulants that promise higher level of safety. Different strategies have been adopted so …

☆ Save Cite Cited by 9 Related articles All 6 versions

Chemical Synthesis and Biochemical Evaluation of Decasulfated Galloyl-glucopyranose Heparin Mimetic

RA Al-Horani - … & hematological agents in medicinal chemistry, 2026 - benthamdirect.com

Introduction Thrombosis remains a major global health burden, necessitating safer and more effective antithrombotic strategies. Current therapies, including vitamin K antagonists …

☆ Save Cite Related articles All 5 versions

Murine models in the evaluation of heparan sulfate-based anticoagulants

D Gailani, Q Cheng, IS Ivanov - Glycosaminoglycans: Chemistry and …, 2014 - Springer

Evaluating anticoagulants in animal thrombosis models is a standard component of preclinical drug testing. Mice are frequently used for these initial evaluations because a …

☆ Save Cite Cited by 11 Related articles All 5 versions

Enantiomeric separation of thiourea derivatives of naringenin on amylose and cellulose polymeric chromatographic chiral columns

I Ali, R Nabti, N Belboukhari, K Sekkoum, MEA Zaid… - Chirality, 2024 - Wiley Online Library

Due to a great demand for amylose and cellulose polymeric chromatographic chiral columns, the enantiomeric separation of thiourea derivatives of naringenin was achieved on …

☆ Save Cite Cited by 1 Related articles All 7 versions

Efforts to total synthesis of philinopside E: convergent synthesis of the sulfated lanostane-type tetraglycoside

S Bai, Z Wu, Q Huang, L Zhang, P Chen, C Wang… - RSC Advances, 2016 - pubs.rsc.org

As an important step to total synthesis of philinopside E with important antitumor activities (Ed50 0.75–3.50 μg mL− 1), we described herein convergent synthesis of a triterpene …

☆ Save Cite Cited by 9 Related articles All 4 versions

Revisiting the effect of cholesteryl sulfate on clotting and fibrinolysis: Inhibition of human thrombin and other human blood proteases

RA Al-Horani - Heliyon, 2024 - cell.com

Cholesteryl sulfate (CS) was quantitatively synthesized by microwave-assisted sulfonation of cholesterol followed by sodium exchange chromatography. In vitro effects of CS on human …

☆ Save Cite Cited by 3 Related articles All 7 versions

Orally bioavailable amine-linked macrocyclic inhibitors of factor XIa

T Fang, JR Corte, PJ Gilligan, Y Jeon, H Osuna… - Bioorganic & Medicinal …, 2020 - Elsevier

The discovery of orally bioavailable FXIa inhibitors has been a challenge. Herein, we describe our efforts to address this challenge by optimization of our imidazole-based …

☆ Save Cite Cited by 8 Related articles All 2 versions

Identifying new clotting factor XIa inhibitors in virtual high-throughput screens using PCA-GA-SVM models and signature

JJ Chen, LN Schmucker, DP Visco Jr - Biotechnology Progress, 2018 - Wiley Online Library

Blood Clotting Factor XI is an important actor in the clotting mechanism: it activates downstream zymogen involved in the clotting process. It can be targeted for activation or …

☆ Save Cite Cited by 3 Related articles All 6 versions

Thiocarbamate-Linked Polysulfonate–Peptide Conjugates As Selective Hepatocyte Growth Factor Receptor Binders

S Besret, J Vicogne, F Dahmani, V Fafeur… - Bioconjugate …, 2014 - pmc.ncbi.nlm.nih.gov

The capacity of many proteins to interact with natural or synthetic polyanions has been exploited for modulating their biological action. However, the polydispersity of these …

☆ Save Cite Cited by 2 Related articles All 6 versions

ASSESSING DRUG DEVELOPMENT PROPORTIES OF ANTITHROMBOTIC AND ANTI-CANCER AGENTS

El Abdelfadiel - 2023 - scholarscompass.vcu.edu

The current GAG anticoagulants such as heparin, heparin derivatives, and vitamin K antagonists, such as warfarin continue to be the backbone of anticoagulant therapy. These …

☆ Save Cite Related articles All 2 versions »